

# BEST AVAILABLE COPY

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
28 August 2003 (28.08.2003)

PCT


(10) International Publication Number  
WO 03/070910 A2

(51) International Patent Classification <sup>7</sup> :	C12N	60/440,129	15 January 2003 (15.01.2003)	US
(21) International Application Number:	PCT/US03/05022	(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier applications:		
(22) International Filing Date:	20 February 2003 (20.02.2003)	US	60/393,796 (CON)	
(25) Filing Language:	English	Filed on	3 July 2002 (03.07.2002)	
(26) Publication Language:	English	US	60/399,348 (CON)	
		Filed on	29 July 2002 (29.07.2002)	
		US	10/306,747 (CON)	
		Filed on	27 November 2002 (27.11.2002)	
(30) Priority Data:		US	10/287,949 (CON)	
60/358,580	20 February 2002 (20.02.2002)	US	Filed on	4 November 2002 (04.11.2002)
60/363,124	11 March 2002 (11.03.2002)	US	US	PCT/US02/17674 (CON)
PCT/US02/17674	29 May 2002 (29.05.2002)	US	Filed on	29 May 2002 (29.05.2002)
60/386,782	6 June 2002 (06.06.2002)	US	US	60/358,580 (CON)
60/393,796	3 July 2002 (03.07.2002)	US	Filed on	20 February 2002 (20.02.2002)
60/399,348	29 July 2002 (29.07.2002)	US	US	60/363,124 (CON)
60/406,784	29 August 2002 (29.08.2002)	US	Filed on	11 March 2002 (11.03.2002)
60/408,378	5 September 2002 (05.09.2002)	US	US	60/386,782 (CON)
60/409,293	9 September 2002 (09.09.2002)	US	Filed on	6 June 2002 (06.06.2002)
10/287,949	4 November 2002 (04.11.2002)	US	US	60/406,784 (CON)
10/306,747	27 November 2002 (27.11.2002)	US	Filed on	29 August 2002 (29.08.2002)

[Continued on next page]

(54) Title: RNA INTERFERENCE MEDIATED INHIBITION OF VASCULAR ENDOTHELIAL GROWTH FACTOR AND VASCULAR ENDOTHELIAL GROWTH FACTOR RECEPTOR GENE EXPRESSION USING SHORT INTERFERING NUCLEIC ACID (siNA)

## A375 24h 36B4 VEGFR1 mRNA Expression



(57) Abstract: The present invention concerns methods and reagents useful in modulating vascular endothelial growth factor (VEGF, VEGF-B, VEGF-C, VEGF-D) and/or vascular endothelial growth factor receptor (e.g., VEGFR1, VEGFR2, and/or VEGFR3) gene expression in a variety of applications, including use in therapeutic, diagnostic, target validation, and genomic discovery applications. Specifically, the invention relates to small nucleic acid molecules, such as short interfering nucleic acid (siNA), short interfering RNA (siRNA), double-stranded RNA (dsRNA), micro-RNA (miRNA), and short hairpin RNA (shRNA) molecules capable of mediating RNA interference (RNAi) against VEGF and/or VEGFR gene expression and/or activity. The small nucleic acid molecules are useful in the diagnosis and treatment of cancer, proliferative diseases, and any other disease or condition that responds to modulation of VEGF and/or VEGFR expression or activity.

WO 03/070910 A2



US 60/408,378 (CON)  
 Filed on 5 September 2002 (05.09.2002)  
 US 60/409,293 (CON)  
 Filed on 9 September 2002 (09.09.2002)  
 US 60/440,129 (CON)  
 Filed on 15 January 2003 (15.01.2003)

(71) **Applicant (for all designated States except US):** RI-BOZYME PHARMACEUTICALS, INCORPORATED [US/US]; 2950 Wilderness Place, Boulder, CO 80301 (US).

(72) **Inventors; and**

(75) **Inventors/Applicants (for US only):** McSWIGGEN, James [US/US]; 4866 Franklin Drive, Boulder, CO 80301 (US). BEIGELMAN, Leonid [US/US]; 5530 Colt Drive, Longmont, CO 80503 (US). PAVCO, Pamela [US/US]; 705 Barberry Circle, Lafayette, CO 80026 (US).

(74) **Agent:** TERPSTRA, Anita, J.; McDonnell Boehnen Hulbert & Berghoff, Suite 3200, 300 South Wacker Drive, Chicago, IL 60606 (US).

(81) **Designated States (national):** AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

**RNA INTERFERENCE MEDIATED INHIBITION OF VASCULAR ENDOTHELIAL  
GROWTH FACTOR AND VASCULAR ENDOTHELIAL GROWTH FACTOR  
RECEPTOR GENE EXPRESSION USING SHORT INTERFERING NUCLEIC ACID  
(siNA)**

5        This invention claims the benefit of McSwiggen, USSN 60/393,796 filed July 3, 2002,  
of McSwiggen, USSN 60/399,348 filed July 29, 2002, of Pavco, USSN 10/306,747, filed  
November 27, 2002, which claims the benefit of Pavco USSN 60/334461, filed November  
30, 2001, of Pavco, USSN 10/287,949 filed November 4, 2002, of Pavco, PCT/US02/17674  
10        filed May 29, 2002, of Beigelman USSN 60/358,580 filed February 20, 2002, of Beigelman  
USSN 60/363,124 filed March 11, 2002, of Beigelman USSN 60/386,782 filed June 6, 2002,  
of Beigelman USSN 60/406,784 filed August 29, 2002, of Beigelman USSN 60/408,378  
filed September 5, 2002, of Beigelman USSN 60/409,293 filed September 9, 2002, and of  
Beigelman USSN 60/440,129 filed January 15, 2003. These applications are hereby  
incorporated by reference herein in their entireties, including the drawings.

15

**Field Of The Invention**

      The present invention concerns compounds, compositions, and methods for the study,  
diagnosis, and treatment of conditions and diseases that respond to the modulation of  
vascular endothelial growth factor (VEGF) and/or vascular endothelial growth factor  
receptor (e.g., VEGFr1, VEGFr2 and/or VEGFr3) gene expression and/or activity. The  
20        present invention also concerns compounds, compositions, and methods relating to  
conditions and diseases that respond to the modulation of expression and/or activity of genes  
involved in VEGF and VEGF receptor pathways. Specifically, the invention relates to small  
nucleic acid molecules, such as short interfering nucleic acid (siNA), short interfering RNA  
(siRNA), double-stranded RNA (dsRNA), micro-RNA (miRNA), and short hairpin RNA  
25        (shRNA) molecules capable of mediating RNA interference (RNAi) against VEGF and  
VEGF receptor gene expression.

### Background Of The Invention

The following is a discussion of relevant art pertaining to RNAi. The discussion is provided only for understanding of the invention that follows. The summary is not an admission that any of the work described below is prior art to the claimed invention.

- 5 RNA interference refers to the process of sequence-specific post-transcriptional gene silencing in animals mediated by short interfering RNAs (siRNAs) (Fire *et al.*, 1998, *Nature*, 391, 806). The corresponding process in plants is commonly referred to as post-transcriptional gene silencing or RNA silencing and is also referred to as quelling in fungi. The process of post-transcriptional gene silencing is thought to be an evolutionarily-  
10 conserved cellular defense mechanism used to prevent the expression of foreign genes and is commonly shared by diverse flora and phyla (Fire *et al.*, 1999, *Trends Genet.*, 15, 358). Such protection from foreign gene expression may have evolved in response to the production of double-stranded RNAs (dsRNAs) derived from viral infection or from the random integration of transposon elements into a host genome via a cellular response that  
15 specifically destroys homologous single-stranded RNA or viral genomic RNA. The presence of dsRNA in cells triggers the RNAi response through a mechanism that has yet to be fully characterized. This mechanism appears to be different from the interferon response that results from dsRNA-mediated activation of protein kinase PKR and 2',5'-oligoadenylate synthetase resulting in non-specific cleavage of mRNA by ribonuclease L.
- 20 The presence of long dsRNAs in cells stimulates the activity of a ribonuclease III enzyme referred to as dicer. Dicer is involved in the processing of the dsRNA into short pieces of dsRNA known as short interfering RNAs (siRNAs) (Berstein *et al.*, 2001, *Nature*, 409, 363). Short interfering RNAs derived from dicer activity are typically about 21 to about 23 nucleotides in length and comprise about 19 base pair duplexes (Elbashir *et al.*,  
25 2001, *Genes Dev.*, 15, 188). Dicer has also been implicated in the excision of 21- and 22-nucleotide small temporal RNAs (stRNAs) from precursor RNA of conserved structure that are implicated in translational control (Hutvagner *et al.*, 2001, *Science*, 293, 834). The RNAi response also features an endonuclease complex, commonly referred to as an RNA-induced silencing complex (RISC), which mediates cleavage of single-stranded RNA having



sequence complementary to the antisense strand of the siRNA duplex. Cleavage of the target RNA takes place in the middle of the region complementary to the antisense strand of the siRNA duplex (Elbashir *et al.*, 2001, *Genes Dev.*, 15, 188).

RNAi has been studied in a variety of systems. Fire *et al.*, 1998, *Nature*, 391, 806, were the first to observe RNAi in *C. elegans*. Wianny and Goetz, 1999, *Nature Cell Biol.*, 2, 70, describe RNAi mediated by dsRNA in mouse embryos. Hammond *et al.*, 2000, *Nature*, 404, 293, describe RNAi in *Drosophila* cells transfected with dsRNA. Elbashir *et al.*, 2001, *Nature*, 411, 494, describe RNAi induced by introduction of duplexes of synthetic 21-nucleotide RNAs in cultured mammalian cells including human embryonic kidney and HeLa cells. Recent work in *Drosophila* embryonic lysates (Elbashir *et al.*, 2001, *EMBO J.*, 20, 6877) has revealed certain requirements for siRNA length, structure, chemical composition, and sequence that are essential to mediate efficient RNAi activity. These studies have shown that 21-nucleotide siRNA duplexes are most active when containing 3'-terminal dinucleotide overhangs. Furthermore, complete substitution of one or both siRNA strands with 2'-deoxy (2'-H) or 2'-O-methyl nucleotides abolishes RNAi activity, whereas substitution of the 3'-terminal siRNA overhang nucleotides with 2'-deoxy nucleotides (2'-H) was shown to be tolerated. Single mismatch sequences in the center of the siRNA duplex were also shown to abolish RNAi activity. In addition, these studies also indicate that the position of the cleavage site in the target RNA is defined by the 5'-end of the siRNA guide sequence rather than the 3'-end of the guide sequence (Elbashir *et al.*, 2001, *EMBO J.*, 20, 6877). Other studies have indicated that a 5'-phosphate on the target-complementary strand of a siRNA duplex is required for siRNA activity and that ATP is utilized to maintain the 5'-phosphate moiety on the siRNA (Nykanen *et al.*, 2001, *Cell*, 107, 309).

Studies have shown that replacing the 3'-terminal nucleotide overhanging segments of a 21-mer siRNA duplex having two -nucleotide 3'-overhangs with deoxyribonucleotides does not have an adverse effect on RNAi activity. Replacing up to four nucleotides on each end of the siRNA with deoxyribonucleotides has been reported to be well tolerated, whereas complete substitution with deoxyribonucleotides results in no RNAi activity (Elbashir *et al.*, 2001, *EMBO J.*, 20, 6877). In addition, Elbashir *et al.*, *supra*, also report that substitution of siRNA with 2'-O-methyl nucleotides completely abolishes RNAi activity. Li *et al.*,

International PCT Publication No. WO 00/44914, and Beach *et al.*, International PCT Publication No. WO 01/68836 preliminarily suggest that siRNA may include modifications to either the phosphate-sugar backbone or the nucleoside to include at least one of a nitrogen or sulfur heteroatom, however, neither application postulates to what extent such  
5 modifications would be tolerated in siRNA molecules, nor provides any further guidance or examples of such modified siRNA. Kreutzer *et al.*, Canadian Patent Application No. 2,359,180, also describe certain chemical modifications for use in dsRNA constructs in order to counteract activation of double-stranded RNA-dependent protein kinase PKR, specifically 2'-amino or 2'-O-methyl nucleotides, and nucleotides containing a 2'-O or 4'-C  
10 methylene bridge. However, Kreutzer *et al.* similarly fails to provide examples or guidance as to what extent these modifications would be tolerated in siRNA molecules.

Parrish *et al.*, 2000, *Molecular Cell*, 6, 1977-1087, tested certain chemical modifications targeting the unc-22 gene in *C. elegans* using long (>25 nt) siRNA transcripts. The authors describe the introduction of thiophosphate residues into these siRNA transcripts  
15 by incorporating thiophosphate nucleotide analogs with T7 and T3 RNA polymerase and observed that RNAs with two phosphorothioate modified bases also had substantial decreases in effectiveness as RNAi. Further, Parrish *et al.* reported that phosphorothioate modification of more than two residues greatly destabilized the RNAs *in vitro* such that interference activities could not be assayed. *Id.* at 1081. The authors also tested certain  
20 modifications at the 2'-position of the nucleotide sugar in the long siRNA transcripts and found that substituting deoxynucleotides for ribonucleotides produced a substantial decrease in interference activity, especially in the case of Uridine to Thymidine and/or Cytidine to deoxy-Cytidine substitutions. *Id.* In addition, the authors tested certain base modifications, including substituting, in sense and antisense strands of the siRNA, 4-thiouracil, 5-  
25 bromouracil, 5-iodouracil, and 3-(aminoallyl)uracil for uracil, and inosine for guanosine. Whereas 4-thiouracil and 5-bromouracil substitution appeared to be tolerated, Parrish reported that inosine produced a substantial decrease in interference activity when incorporated in either strand. Parrish also reported that incorporation of 5-iodouracil and 3-(aminoallyl)uracil in the antisense strand resulted in a substantial decrease in RNAi activity  
30 as well.

The use of longer dsRNA has been described. For example, Beach *et al.*, International PCT Publication No. WO 01/68836, describes specific methods for attenuating gene expression using endogenously-derived dsRNA. Tuschl *et al.*, International PCT Publication No. WO 01/75164, describe a *Drosophila in vitro* RNAi system and the use of specific siRNA molecules for certain functional genomic and certain therapeutic applications; although Tuschl, 2001, *Chem. Biochem.*, 2, 239-245, doubts that RNAi can be used to cure genetic diseases or viral infection due to the danger of activating interferon response. Li *et al.*, International PCT Publication No. WO 00/44914, describe the use of specific dsRNAs for attenuating the expression of certain target genes. Zernicka-Goetz *et al.*, International PCT Publication No. WO 01/36646, describe certain methods for inhibiting the expression of particular genes in mammalian cells using certain dsRNA molecules. Fire *et al.*, International PCT Publication No. WO 99/32619, describe particular methods for introducing certain dsRNA molecules into cells for use in inhibiting gene expression. Plaetinck *et al.*, International PCT Publication No. WO 00/01846, describe certain methods for identifying specific genes responsible for conferring a particular phenotype in a cell using specific dsRNA molecules. Mello *et al.*, International PCT Publication No. WO 01/29058, describe the identification of specific genes involved in dsRNA-mediated RNAi. Deschamps Depaillette *et al.*, International PCT Publication No. WO 99/07409, describe specific compositions consisting of particular dsRNA molecules combined with certain anti-viral agents. Waterhouse *et al.*, International PCT Publication No. 99/53050, describe certain methods for decreasing the phenotypic expression of a nucleic acid in plant cells using certain dsRNAs. Driscoll *et al.*, International PCT Publication No. WO 01/49844, describe specific DNA constructs for use in facilitating gene silencing in targeted organisms.

Others have reported on various RNAi and gene-silencing systems. For example, Parrish *et al.*, 2000, *Molecular Cell*, 6, 1977-1087, describe specific chemically-modified siRNA constructs targeting the unc-22 gene of *C. elegans*. Grossniklaus, International PCT Publication No. WO 01/38551, describes certain methods for regulating polycomb gene expression in plants using certain dsRNAs. Churikov *et al.*, International PCT Publication No. WO 01/42443, describe certain methods for modifying genetic characteristics of an organism using certain dsRNAs. Cogoni *et al.*, International PCT Publication No. WO

01/53475, describe certain methods for isolating a *Neurospora* silencing gene and uses thereof. Reed *et al.*, International PCT Publication No. WO 01/68836, describe certain methods for gene silencing in plants. Honer *et al.*, International PCT Publication No. WO 01/70944, describe certain methods of drug screening using transgenic nematodes as  
5 Parkinson's Disease models using certain dsRNAs. Deak *et al.*, International PCT Publication No. WO 01/72774, describe certain *Drosophila*-derived gene products that may be related to RNAi in *Drosophila*. Arndt *et al.*, International PCT Publication No. WO 01/92513 describe certain methods for mediating gene suppression by using factors that enhance RNAi. Tuschl *et al.*, International PCT Publication No. WO 02/44321, describe  
10 certain synthetic siRNA constructs. Pachuk *et al.*, International PCT Publication No. WO 00/63364, and Satishchandran *et al.*, International PCT Publication No. WO 01/04313, describe certain methods and compositions for inhibiting the function of certain polynucleotide sequences using certain dsRNAs. Echeverri *et al.*, International PCT Publication No. WO 02/38805, describe certain *C. elegans* genes identified via RNAi.  
15 Kreutzer *et al.*, International PCT Publications Nos. WO 02/055692, WO 02/055693, and EP 1144623 B1 describes certain methods for inhibiting gene expression using RNAi. Graham *et al.*, International PCT Publications Nos. WO 99/49029 and WO 01/70949, and AU 4037501 describe certain vector expressed siRNA molecules. Fire *et al.*, US 6,506,559, describe certain methods for inhibiting gene expression in vitro using certain long dsRNA  
20 (greater than 25 nucleotide) constructs that mediate RNAi.

### SUMMARY OF THE INVENTION

This invention relates to compounds, compositions, and methods useful for modulating the expression of genes, such as those genes associated with angiogenesis and proliferation using short interfering nucleic acid (siNA) molecules. This invention also  
25 relates to compounds, compositions, and methods useful for modulating the expression and activity of vascular endothelial growth factor (VEGF) and/or vascular endothelial growth factor receptor (e.g., VEGFr1, VEGFr2, VEGFr3) genes, or genes involved in VEGF and/or VEGFr pathways of gene expression and/or VEGF activity by RNA interference (RNAi) using small nucleic acid molecules, such as short interfering nucleic acid (siNA), short

interfering RNA (siRNA), double-stranded RNA (dsRNA), micro-RNA (miRNA), and short hairpin RNA (shRNA) molecules. In particular, the instant invention features small nucleic acid molecules, such as short interfering nucleic acid (siNA), short interfering RNA (siRNA), double-stranded RNA (dsRNA), micro-RNA (miRNA), and short hairpin RNA (shRNA) molecules and methods used to modulate the expression of VEGF and/or VEGFr genes. A siNA of the invention can be unmodified or chemically-modified. A siNA of the instant invention can be chemically synthesized, expressed from a vector or enzymatically synthesized. The instant invention also features various chemically-modified synthetic short interfering nucleic acid (siNA) molecules capable of modulating VEGF and/or VEGFr gene expression or activity in cells by RNA interference (RNAi). The use of chemically-modified siNA improves various properties of native siNA molecules through increased resistance to nuclease degradation *in vivo* and/or through improved cellular uptake. Further, contrary to earlier published studies, siNA having multiple chemical modifications retains its RNAi activity. The siNA molecules of the instant invention provide useful reagents and methods for a variety of therapeutic, diagnostic, target validation, genomic discovery, genetic engineering, and pharmacogenomic applications.

In one embodiment, the invention features one or more siNA molecules and methods that independently or in combination modulate the expression of gene(s) encoding proteins, such as vascular endothelial growth factor (VEGF) and/or vascular endothelial growth factor receptors (e.g., VEGFr1, VEGFr2, VEGFr3), associated with the maintenance and/or development of cancer and other proliferative diseases, such as genes encoding sequences comprising those sequences referred to by GenBank Accession Nos. shown in Table I, referred to herein generally as VEGF and/or VEGFr. The description below of the various aspects and embodiments of the invention is provided with reference to the exemplary VEGF and VEGFr (e.g., VEGFr1, VEGFr2, VEGFr3) genes referred to herein as VEGF and VEGFr respectively. However, the various aspects and embodiments are also directed to other VEGF and/or VEGFr genes, such as mutant VEGF and/or VEGFr genes, splice variants of VEGF and/or VEGFr genes, other VEGF and/or VEGFr ligands and receptors. The various aspects and embodiments are also directed to other genes that are involved in VEGF and/or VEGFr mediated pathways of signal transduction or gene expression that are

involved in the progression, development, and/or maintenance of disease (e.g., cancer). Those additional genes can be analyzed for target sites using the methods described for VEGF and/or VEGFr genes herein. Thus, the inhibition and the effects of such inhibition of the other genes can be performed as described herein.

5        In one embodiment, the invention features a siNA molecule that down-regulates expression of a VEGF gene, for example, wherein the VEGF gene comprises VEGF encoding sequence.

10        In one embodiment, the invention features a siNA molecule that down-regulates expression of a VEGFr gene, for example, wherein the VEGFr gene comprises VEGFr encoding sequence.

15        In one embodiment, the invention features a siNA molecule having RNAi activity against VEGF and/or VEGFr RNA, wherein the siNA molecule comprises a sequence complementary to any RNA having VEGF and/or VEGFr or other VEGF and/or VEGFr encoding sequence, such as those sequences having GenBank Accession Nos. shown in Table I. Chemical modifications as shown in Tables III and IV or otherwise described herein can be applied to any siNA construct of the invention.

20        In one embodiment, the invention features a siNA molecule having RNAi activity against VEGF and/or VEGFr RNA, wherein the siNA molecule comprises a sequence complementary to any RNA having VEGF and/or VEGFr encoding sequence, such as those sequences having VEGF and/or VEGFr GenBank Accession Nos. shown in Table I. Chemical modifications as shown in Tables III and IV or otherwise described herein can be applied to any siNA construct of the invention.

25        In another embodiment, the invention features a siNA molecule having RNAi activity against a VEGF and/or VEGFr gene, wherein the siNA molecule comprises nucleotide sequence complementary to nucleotide sequence of a VEGF and/or VEGFr gene, such as those VEGF and/or VEGFr sequences having GenBank Accession Nos. shown in Table I. In another embodiment, a siNA molecule of the invention includes nucleotide sequence that can interact with nucleotide sequence of a VEGF and/or VEGFr gene and thereby mediate

silencing of VEGF and/or VEGFr gene expression, for example, wherein the siNA mediates regulation of VEGF and/or VEGFr gene expression by cellular processes that modulate the chromatin structure of the VEGF and/or VEGFr gene and prevent transcription of the VEGF and/or VEGFr gene.

5 In another embodiment, the invention features a siNA molecule comprising nucleotide sequence, for example, nucleotide sequence in the antisense region of the siNA molecule that is complementary to a nucleotide sequence or portion of sequence of a VEGF and/or VEGFr gene. In another embodiment, the invention features a siNA molecule comprising a region, for example, the antisense region of the siNA construct, complementary to a  
10 sequence or portion of sequence comprising a VEGF and/or VEGFr gene sequence.

In one embodiment, the antisense region of VEGFr1 siNA constructs can comprise a sequence complementary to sequence having any of SEQ ID NOs. 1-427 or 1997-2000. In one embodiment, the antisense region can also comprise sequence having any of SEQ ID NOs. 428-854, 2024-2027, 2032-2035, 2040-2043, 2104-2107, 2109, 2117, 2120-2122,  
15 2125-2132, 2137-2140, 2142, 2150, 2152, 2154, 2158-2160, 2164-2166, 2188-2190, 2197, 2199, 2203-2204, 2229, 2231, 2233, 2235, 2237, or 2238. In another embodiment, the sense region of VEGFr1 constructs can comprise sequence having any of SEQ ID NOs. 1-427, 1997-2000, 2009-2016, 2020-2023, 2028-2031, 2036-2039, 2092-2103, 2108, 2114, 2116, 2123-2124, 2133-2136, 2141, 2149, 2151, 2153, 2155-2157, 2161-2163, 2185-2187, 2198,  
20 2200-2202, 2228, 2230, 2232, 2234, or 2236. The sense region can comprise a sequence of SEQ ID NO. 2217 and the antisense region can comprise a sequence of SEQ ID NO. 2218. The sense region can comprise a sequence of SEQ ID NO. 2219 and the antisense region can comprise a sequence of SEQ ID NO. 2220. The sense region can comprise a sequence of SEQ ID NO. 2221 and the antisense region can comprise a sequence of SEQ ID NO. 2222.  
25 The sense region can comprise a sequence of SEQ ID NO. 2223 and the antisense region can comprise a sequence of SEQ ID NO. 2224. The sense region can comprise a sequence of SEQ ID NO. 2225 and the antisense region can comprise a sequence of SEQ ID NO. 2226. The sense region can comprise a sequence of SEQ ID NO. 2223 and the antisense region can comprise a sequence of SEQ ID NO. 2227.

In one embodiment, the antisense region of VEGFr2 siNA constructs can comprise a sequence complementary to sequence having any of SEQ ID NOs. 855-1178 or 2001-2004.

In one embodiment, the antisense region can also comprise sequence having any of SEQ ID NOs. 1179-1502, 2048-2051, 2056-2059, 2064-2067, 2208-2210, 2214-2216, or 2048-2051.

5 In another embodiment, the sense region of VEGFr2 constructs can comprise sequence having any of SEQ ID NOs. 855-1178, 2001-2004, 2044-2047, 2052-2055, 2060-2063, 2017-2019, 2205-2207, 2211-2213, or 2044-2047. The sense region can comprise a sequence of SEQ ID NO. 2217 and the antisense region can comprise a sequence of SEQ ID NO. 2218. The sense region can comprise a sequence of SEQ ID NO. 2219 and the  
10 antisense region can comprise a sequence of SEQ ID NO. 2220. The sense region can comprise a sequence of SEQ ID NO. 2221 and the antisense region can comprise a sequence of SEQ ID NO. 2222. The sense region can comprise a sequence of SEQ ID NO. 2223 and the antisense region can comprise a sequence of SEQ ID NO. 2224. The sense region can comprise a sequence of SEQ ID NO. 2225 and the antisense region can comprise a sequence  
15 of SEQ ID NO. 2226. The sense region can comprise a sequence of SEQ ID NO. 2223 and the antisense region can comprise a sequence of SEQ ID NO. 2227.

In one embodiment, the antisense region of VEGFr3 siNA constructs can comprise a sequence complementary to sequence having any of SEQ ID NOs. 1503-1749 or 2005-2008.

In one embodiment, the antisense region can also comprise sequence having any of SEQ ID  
20 NOs. 1750-1996, 2072-2075, 2080-2083, or 2088-2091. In another embodiment, the sense region of VEGFr3 constructs can comprise sequence having any of SEQ ID NOs. 1503-1749, 2005-2008, 2068-2071, 2076-2079, or 2034-2087. The sense region can comprise a sequence of SEQ ID NO. 2217 and the antisense region can comprise a sequence of SEQ ID NO. 2218. The sense region can comprise a sequence of SEQ ID NO. 2219 and the  
25 antisense region can comprise a sequence of SEQ ID NO. 2220. The sense region can comprise a sequence of SEQ ID NO. 2221 and the antisense region can comprise a sequence of SEQ ID NO. 2222. The sense region can comprise a sequence of SEQ ID NO. 2223 and the antisense region can comprise a sequence of SEQ ID NO. 2224. The sense region can comprise a sequence of SEQ ID NO. 2225 and the antisense region can comprise a sequence



of SEQ ID NO. 2226. The sense region can comprise a sequence of SEQ ID NO. 2223 and the antisense region can comprise a sequence of SEQ ID NO. 2227.

In one embodiment, a siNA molecule of the invention comprises any of SEQ ID NOs. 1-2238. The sequences shown in SEQ ID NOs: 1-2238 are not limiting. A siNA molecule  
5 of the invention can comprise any contiguous VEGF and/or VEGFr sequence (e.g., about 19 to about 25, or about 19, 20, 21, 22, 23, 24 or 25 contiguous VEGF and/or VEGFr nucleotides).

In yet another embodiment, the invention features a siNA molecule comprising a sequence, for example, the antisense sequence of the siNA construct, complementary to a  
10 sequence or portion of sequence comprising sequence represented by GenBank Accession Nos. shown in Table I. Chemical modifications in Tables III and IV and described herein can be applied to any siRNA construct of the invention.

In one embodiment of the invention a siNA molecule comprises an antisense strand having about 19 to about 29 nucleotides, wherein the antisense strand is complementary to a  
15 RNA sequence encoding a VEGF and/or VEGFr protein, and wherein said siNA further comprises a sense strand having about 19 to about 29 (e.g., about 19, 20, 21, 22, 23, 24, 25, 26, 27, 28 or 29) nucleotides, and wherein said sense strand and said antisense strand are distinct nucleotide sequences with at least about 19 complementary nucleotides.

In another embodiment of the invention a siNA molecule of the invention comprises  
20 an antisense region having about 19 to about 29 (e.g., about 19, 20, 21, 22, 23, 24, 25, 26, 27, 28 or 29) nucleotides, wherein the antisense region is complementary to a RNA sequence encoding a VEGF and/or VEGFr protein, and wherein said siNA further comprises a sense region having about 19 to about 29 nucleotides, wherein said sense region and said antisense region comprise a linear molecule with at least about 19 complementary  
25 nucleotides.

In one embodiment of the invention a siNA molecule comprises an antisense strand comprising a nucleotide sequence that is complementary to a nucleotide sequence or a portion thereof encoding a VEGF and/or VEGFr protein. The siNA further comprises a

sense strand, wherein said sense strand comprises a nucleotide sequence of a VEGF and/or VEGFr gene or a portion thereof.

In another embodiment, a siNA molecule comprises an antisense region comprising a nucleotide sequence that is complementary to a nucleotide sequence or a portion thereof encoding a VEGF and/or VEGFr protein. The siNA molecule further comprises a sense region, wherein said sense region comprises a nucleotide sequence of a VEGF and/or VEGFr gene or a portion thereof.

In one embodiment, a siNA molecule of the invention has RNAi activity that modulates expression of RNA encoded by a VEGF gene. Because VEGF genes can share some degree of sequence homology with each other, siNA molecules can be designed to target a class of VEGF genes (and associated receptor or ligand genes) or alternately specific VEGF genes by selecting sequences that are either shared amongst different VEGF targets or alternatively that are unique for a specific VEGF target. Therefore, in one embodiment, the siNA molecule can be designed to target conserved regions of VEGF RNA sequence having homology between several VEGF genes so as to target several VEGF genes (e.g., different VEGF isoforms, splice variants, mutant genes etc.) with one siNA molecule. In another embodiment, the siNA molecule can be designed to target a sequence that is unique to a specific VEGF RNA sequence due to the high degree of specificity that the siNA molecule requires to mediate RNAi activity.

In one embodiment, a siNA molecule of the invention has RNAi activity that modulates expression of RNA encoded by a VEGFr gene. Because VEGFr genes can share some degree of sequence homology with each other, siNA molecules can be designed to target a class of VEGFr genes (and associated receptor or ligand genes) or alternately specific VEGFr genes by selecting sequences that are either shared amongst different VEGFr targets or alternatively that are unique for a specific VEGFr target. Therefore, in one embodiment, the siNA molecule can be designed to target conserved regions of VEGFr RNA sequence having homology between several VEGFr genes so as to target several VEGFr genes (e.g., different VEGFr isoforms, splice variants, mutant genes etc.) with one siNA molecule. In another embodiment, the siNA molecule can be designed to target a

sequence that is unique to a specific VEGFr RNA sequence due to the high degree of specificity that the siNA molecule requires to mediate RNAi activity.

In one embodiment, a siNA molecule of the invention has RNAi activity that modulates expression of RNA encoded by a VEGFr gene. Because VEGFr genes can share some degree of sequence homology with each other, siNA molecules can be designed to target a class of VEGFr genes or alternately specific VEGFr genes by selecting sequences that are either shared amongst different VEGFr targets or alternatively that are unique for a specific VEGFr target. Therefore, in one embodiment, the siNA molecule can be designed to target conserved regions of VEGFr RNA sequence having homology between several VEGFr genes so as to target several VEGFr genes (e.g., VEGFr1, VEGFr2 and/or VEGFr3, different VEGFr isoforms, splice variants, mutant genes etc.) with one siNA molecule. In another embodiment, the siNA molecule can be designed to target a sequence that is unique to a specific VEGFr RNA sequence due to the high degree of specificity that the siNA molecule requires to mediate RNAi activity.

In one embodiment, a siNA molecule of the invention has RNAi activity that modulates expression of RNA encoded by a VEGF gene. Because VEGF genes can share some degree of sequence homology with each other, siNA molecules can be designed to target a class of VEGF genes or alternately specific VEGF genes by selecting sequences that are either shared amongst different VEGF targets or alternatively that are unique for a specific VEGF target. Therefore, in one embodiment, the siNA molecule can be designed to target conserved regions of VEGF RNA sequence having homology between several VEGF genes so as to target several VEGF genes (e.g., VEGF-A, VEGF-B, VEGF-C and/or VEGF-D, different VEGF isoforms, splice variants, mutant genes etc.) with one siNA molecule. In another embodiment, the siNA molecule can be designed to target a sequence that is unique to a specific VEGF RNA sequence due to the high degree of specificity that the siNA molecule requires to mediate RNAi activity.

In one embodiment, nucleic acid molecules of the invention that act as mediators of the RNA interference gene silencing response are double-stranded nucleic acid molecules. In another embodiment, the siNA molecules of the invention consist of duplexes containing

about 19 base pairs between oligonucleotides comprising about 19 to about 25 (e.g., about 19, 20, 21, 22, 23, 24 or 25) nucleotides. In yet another embodiment, siNA molecules of the invention comprise duplexes with overhanging ends of about about 1 to about 3 (e.g., about 1, 2, or 3) nucleotides, for example, about 21-nucleotide duplexes with about 19 base pairs and 3'-terminal mononucleotide, dinucleotide, or trinucleotide overhangs.

In one embodiment, the invention features one or more chemically-modified siNA constructs having specificity for VEGF and/or VEGFr expressing nucleic acid molecules, such as RNA encoding a VEGF and/or VEGFr protein. Non-limiting examples of such chemical modifications include without limitation phosphorothioate internucleotide linkages, 2'-deoxyribonucleotides, 2'-O-methyl ribonucleotides, 2'-deoxy-2'-fluoro ribonucleotides, "universal base" nucleotides, "acyclic" nucleotides, 5-C-methyl nucleotides, and terminal glyceryl and/or inverted deoxy abasic residue incorporation. These chemical modifications, when used in various siNA constructs, are shown to preserve RNAi activity in cells while at the same time, dramatically increasing the serum stability of these compounds. Furthermore, contrary to the data published by Parrish *et al.*, *supra*, applicant demonstrates that multiple (greater than one) phosphorothioate substitutions are well-tolerated and confer substantial increases in serum stability for modified siNA constructs.

In one embodiment, a siNA molecule of the invention comprises modified nucleotides while maintaining the ability to mediate RNAi. The modified nucleotides can be used to improve *in vitro* or *in vivo* characteristics such as stability, activity, and/or bioavailability. For example, a siNA molecule of the invention can comprise modified nucleotides as a percentage of the total number of nucleotides present in the siNA molecule. As such, a siNA molecule of the invention can generally comprise about 5% to about 100% modified nucleotides (e.g., 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 100% modified nucleotides). The actual percentage of modified nucleotides present in a given siNA molecule will depend on the total number of nucleotides present in the siNA. If the siNA molecule is single stranded, the percent modification can be based upon the total number of nucleotides present in the single stranded siNA molecules. Likewise, if the siNA molecule is double stranded, the percent

modification can be based upon the total number of nucleotides present in the sense strand, antisense strand, or both the sense and antisense strands.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a VEGF and/or VEGFr gene, wherein the siNA molecule comprises one or more chemical modifications and each strand of the double-stranded siNA is about 21 nucleotides long.

In one embodiment, a siNA molecule of the invention comprises no ribonucleotides. In another embodiment, a siNA molecule of the invention comprises ribonucleotides.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a VEGF and/or VEGFr gene, wherein one of the strands of the double-stranded siNA molecule comprises a nucleotide sequence that is complementary to a nucleotide sequence or a portion thereof of the VEGF and/or VEGFr gene, and wherein the second strand of the double-stranded siNA molecule comprises a nucleotide sequence substantially similar to the nucleotide sequence or a portion thereof of the VEGF and/or VEGFr gene.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a VEGF and/or VEGFr gene, wherein each strand of the siNA molecule comprises about 19 to about 23 nucleotides, and wherein each strand comprises at least about 19 nucleotides that are complementary to the nucleotides of the other strand.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a VEGF and/or VEGFr gene, wherein the siNA molecule comprises an antisense region comprising a nucleotide sequence that is complementary to a nucleotide sequence or a portion thereof of the VEGF and/or VEGFr gene, and wherein the siNA further comprises a sense region, wherein the sense region comprises a nucleotide sequence substantially similar to the nucleotide sequence or a portion thereof of the VEGF and/or VEGFr gene.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a VEGF and/or VEGFr gene, wherein the antisense region and the sense region each comprise about 19 to about 23 nucleotides, and wherein the antisense region comprises at least about 19 nucleotides that  
5 are complementary to nucleotides of the sense region.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a VEGF and/or VEGFr gene, wherein the siNA molecule comprises a sense region and an antisense region and wherein the antisense region comprises a nucleotide sequence that is complementary to a nucleotide  
10 sequence or a portion thereof of RNA encoded by the VEGF and/or VEGFr gene and the sense region comprises a nucleotide sequence that is complementary to the antisense region.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a VEGF and/or VEGFr gene, wherein the siNA molecule is assembled from two separate oligonucleotide fragments  
15 wherein one fragment comprises the sense region and the second fragment comprises the antisense region of the siNA molecule. The sense region can be connected to the antisense region via a linker molecule, such as a polynucleotide linker or a non-nucleotide linker.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a VEGF and/or VEGFr gene,  
20 wherein the siNA molecule comprises a sense region and an antisense region and wherein the antisense region comprises a nucleotide sequence that is complementary to a nucleotide sequence or a portion thereof of RNA encoded by the VEGF and/or VEGFr gene and the sense region comprises a nucleotide sequence that is complementary to the antisense region, and wherein pyrimidine nucleotides in the sense region are 2'-O-methyl pyrimidine  
25 nucleotides, 2'-deoxy purine nucleotides, or 2'-deoxy-2'-fluoro pyrimidine nucleotides.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a VEGF and/or VEGFr gene, wherein the siNA molecule is assembled from two separate oligonucleotide fragments

wherein one fragment comprises the sense region and the second fragment comprises the antisense region of the siNA molecule, and wherein the fragment comprising the sense region includes a terminal cap moiety at the 5'-end, the 3'-end, or both of the 5' and 3' ends of the fragment comprising the sense region. In another embodiment, the terminal cap moiety is an inverted deoxy abasic moiety or glyceryl moiety. In another embodiment, each of the two fragments of the siNA molecule comprise about 21 nucleotides.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a VEGF and/or VEGFr gene, wherein the siNA molecule comprises a sense region and an antisense region and wherein the antisense region comprises a nucleotide sequence that is complementary to a nucleotide sequence or a portion thereof of RNA encoded by the VEGF and/or VEGFr gene and the sense region comprises a nucleotide sequence that is complementary to the antisense region, and wherein the purine nucleotides present in the antisense region comprise 2'-deoxy- purine nucleotides. In another embodiment, the antisense region comprises a phosphorothioate internucleotide linkage at the 3' end of the antisense region. In another embodiment, the antisense region comprises a glyceryl modification at the 3' end of the antisense region.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a VEGF and/or VEGFr gene, wherein the siNA molecule is assembled from two separate oligonucleotide fragments wherein one fragment comprises the sense region and the second fragment comprises the antisense region of the siNA molecule, and wherein about 19 nucleotides of each fragment of the siNA molecule are base-paired to the complementary nucleotides of the other fragment of the siNA molecule and wherein at least two 3' terminal nucleotides of each fragment of the siNA molecule are not base-paired to the nucleotides of the other fragment of the siNA molecule. In another embodiment, each of the two 3' terminal nucleotides of each fragment of the siNA molecule are 2'-deoxy-pyrimidines, such as 2'-deoxy-thymidine. In another embodiment, all 21 nucleotides of each fragment of the siNA molecule are base-paired to the complementary nucleotides of the other fragment of the siNA molecule. In another embodiment, about 19 nucleotides of the antisense region are base-paired to the nucleotide sequence or a portion thereof of the RNA encoded by the VEGF and/or VEGFr

gene. In another embodiment, 21 nucleotides of the antisense region are base-paired to the nucleotide sequence or a portion thereof of the RNA encoded by the VEGF and/or VEGFr gene. In another embodiment, the 5'-end of the fragment comprising said antisense region optionally includes a phosphate group.

5 In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits the expression of a VEGF and/or VEGFr RNA sequence (e.g., wherein said target RNA sequence is encoded by a VEGF and/or VEGFr gene), wherein the siNA molecule comprises no ribonucleotides and wherein each strand of the double-stranded siNA molecule is about 21 nucleotides long.

10 In one embodiment, the invention features a medicament comprising a siNA molecule of the invention.

In one embodiment, the invention features an active ingredient comprising a siNA molecule of the invention.

15 In one embodiment, the invention features the use of a double-stranded short interfering nucleic acid (siNA) molecule to down-regulate expression of a VEGF and/or VEGFr gene, wherein the siNA molecule comprises one or more chemical modifications and each strand of the double-stranded siNA is about 21 nucleotides long.

In one embodiment, a VEGFr gene contemplated by the invention is a VEGFr1, VEGFr2, or VEGFr3 gene.

20 In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide  
25 sequence that is complementary to a nucleotide sequence of the antisense strand and wherein a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule



comprises a sugar modification. In one embodiment, the VEGFr gene is VEGFr2. In one embodiment, the VEGFr gene is VEGFr1.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises  
5 nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide sequence that is complementary to a nucleotide sequence of the antisense strand and wherein  
10 a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification, and wherein the nucleotide sequence of the antisense strand of the double-stranded siNA molecule is complementary to the nucleotide sequence of the VEGF and/or VEGFr RNA or a portion thereof which encodes an protein or a portion thereof.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one  
15 of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide sequence that is complementary to a nucleotide sequence of the antisense strand and wherein  
20 a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification, and wherein each strand of the siNA molecule comprises about 19 to about 29 nucleotides, and wherein each strand comprises at least about 19 nucleotides that are complementary to the nucleotides of the other strand.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one  
25 of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide sequence that is complementary to a nucleotide sequence of the antisense strand and wherein

a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification, and wherein the siNA molecule is assembled from two oligonucleotide fragments wherein one fragment comprises the nucleotide sequence of the antisense strand of the siNA molecule and a second fragment comprises nucleotide sequence of the sense region of the siNA molecule.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide sequence that is complementary to a nucleotide sequence of the antisense strand and wherein a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification, and wherein the sense strand is connected to the antisense strand via a linker molecule, such as a polynucleotide linker or a non-nucleotide linker.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide sequence that is complementary to a nucleotide sequence of the antisense strand and wherein a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification, and wherein pyrimidine nucleotides present in the sense strand are 2'-deoxy-2'-fluoro pyrimidine nucleotides and wherein purine nucleotides present in the sense region are 2'-deoxy purine nucleotides.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide

sequence that is complementary to a nucleotide sequence of the antisense strand and wherein a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification, and wherein the sense strand comprises a 3'-end and a 5'-end, and wherein a terminal cap moiety (e.g., an inverted deoxy abasic moiety) is present at the 5'-end, the 3'-end, or both of the 5' and 3' ends of the sense strand.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide sequence that is complementary to a nucleotide sequence of the antisense strand and wherein a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification, and wherein the antisense strand comprises one or more 2'-deoxy-2'-fluoro pyrimidine nucleotides and one or more 2'-O-methyl purine nucleotides.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide sequence that is complementary to a nucleotide sequence of the antisense strand and wherein a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification, and wherein the pyrimidine nucleotides present in the antisense strand are 2'-deoxy-2'-fluoro pyrimidine nucleotides and wherein any purine nucleotides present in the antisense strand are 2'-O-methyl purine nucleotides.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide

sequence that is complementary to a nucleotide sequence of the antisense strand and wherein a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification, and wherein the antisense strand comprises a phosphorothioate internucleotide linkage at the 3' end of the antisense strand.

5 In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide  
10 sequence that is complementary to a nucleotide sequence of the antisense strand and wherein a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification, and wherein the antisense strand comprises a glyceryl modification at the 3' end.

In one embodiment, the invention features a double-stranded short interfering nucleic  
15 acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide sequence that is complementary to a nucleotide sequence of the antisense strand and wherein  
20 a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification, and wherein each of the two strands of the siNA molecule comprises 21 nucleotides. In another embodiment, about 19 nucleotides of each strand of the siNA molecule are base-paired to the complementary nucleotides of the other strand of the siNA molecule and wherein at least two 3' terminal nucleotides of each strand of the  
25 siNA molecule are not base-paired to the nucleotides of the other strand of the siNA molecule. In another embodiment, each of the two 3' terminal nucleotides of each fragment of the siNA molecule are 2'-deoxy-pyrimidines, such as 2'-deoxy-thymidine. In another embodiment, each strand of the siNA molecule are base-paired to the complementary nucleotides of the other strand of the siNA molecule. In another embodiment, about 19  
30 nucleotides of the antisense strand are base-paired to the nucleotide sequence of the VEGF

and/or VEGFr RNA or a portion thereof. In another embodiment, 21 nucleotides of the antisense strand are base-paired to the nucleotide sequence of the VEGF and/or VEGFr RNA or a portion thereof.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide sequence that is complementary to a nucleotide sequence of the antisense strand and wherein a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification, and wherein the 5'-end of the antisense strand optionally includes a phosphate group.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide sequence that is complementary to a nucleotide sequence of the antisense strand and wherein a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification, and wherein the nucleotide sequence or a portion thereof of the antisense strand is complementary to a nucleotide sequence of the 5'-untranslated region or a portion thereof of the VEGF and/or VEGFr RNA.

In one embodiment, the invention features a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide sequence that is complementary to a nucleotide sequence of the antisense strand and wherein a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule

comprises a sugar modification, and wherein the nucleotide sequence or a portion thereof of the antisense strand is complementary to a nucleotide sequence of the VEGF and/or VEGFr RNA or a portion thereof that is present in the VEGF and/or VEGFr RNA.

In one embodiment, the invention features a pharmaceutical composition comprising a siNA molecule of the invention in an acceptable carrier or diluent.

In one embodiment, the invention features a medicament comprising an siNA molecule of the invention.

In one embodiment, the invention features an active ingredient comprising an siNA molecule of the invention.

In one embodiment, the invention features the use of a double-stranded short interfering nucleic acid (siNA) molecule that inhibits expression of a VEGF and/or VEGFr gene, wherein one of the strands of the double-stranded siNA molecule is an antisense strand which comprises nucleotide sequence that is complementary to nucleotide sequence of VEGF and/or VEGFr RNA or a portion thereof, the other strand is a sense strand which comprises nucleotide sequence that is complementary to a nucleotide sequence of the antisense strand and wherein a majority of the pyrimidine nucleotides present in the double-stranded siNA molecule comprises a sugar modification.


In a non-limiting example, the introduction of chemically-modified nucleotides into nucleic acid molecules provides a powerful tool in overcoming potential limitations of *in vivo* stability and bioavailability inherent to native RNA molecules that are delivered exogenously. For example, the use of chemically-modified nucleic acid molecules can enable a lower dose of a particular nucleic acid molecule for a given therapeutic effect since chemically-modified nucleic acid molecules tend to have a longer half-life in serum. Furthermore, certain chemical modifications can improve the bioavailability of nucleic acid molecules by targeting particular cells or tissues and/or improving cellular uptake of the nucleic acid molecule. Therefore, even if the activity of a chemically-modified nucleic acid molecule is reduced as compared to a native nucleic acid molecule, for example, when compared to an all-RNA nucleic acid molecule, the overall activity of the modified nucleic

acid molecule can be greater than that of the native molecule due to improved stability and/or delivery of the molecule. Unlike native unmodified siNA, chemically-modified siNA can also minimize the possibility of activating interferon activity in humans.

5 The antisense region of a siNA molecule of the invention can comprise a phosphorothioate internucleotide linkage at the 3'-end of said antisense region. The antisense region can comprise about one to about five phosphorothioate internucleotide linkages at the 5'-end of said antisense region. The 3'-terminal nucleotide overhangs of a siNA molecule of the invention can comprise ribonucleotides or deoxyribonucleotides that are chemically-modified at a nucleic acid sugar, base, or backbone. The 3'-terminal  
10 nucleotide overhangs can comprise one or more universal base ribonucleotides. The 3'-terminal nucleotide overhangs can comprise one or more acyclic nucleotides.

One embodiment of the invention provides an expression vector comprising a nucleic acid sequence encoding at least one siNA molecule of the invention in a manner that allows expression of the nucleic acid molecule. Another embodiment of the invention provides a  
15 mammalian cell comprising such an expression vector. The mammalian cell can be a human cell. The siNA molecule of the expression vector can comprise a sense region and an antisense region. The antisense region can comprise sequence complementary to a RNA or DNA sequence encoding VEGF and/or VEGFr and the sense region can comprise sequence complementary to the antisense region. The siNA molecule can comprise two distinct  
20 strands having complementary sense and antisense regions. The siNA molecule can comprise a single strand having complementary sense and antisense regions.

In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule capable of mediating RNA interference (RNAi) against a VEGF and/or VEGFr inside a cell or reconstituted *in vitro* system, wherein the chemical  
25 modification comprises one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) nucleotides comprising a backbone modified internucleotide linkage having Formula I:




wherein each R1 and R2 is independently any nucleotide, non-nucleotide, or polynucleotide which can be naturally-occurring or chemically-modified, each X and Y is independently O, S, N, alkyl, or substituted alkyl, each Z and W is independently O, S, N, alkyl, substituted alkyl, O-alkyl, S-alkyl, alkaryl, or aralkyl, and wherein W, X, Y, and Z are optionally not all O.

The chemically-modified internucleotide linkages having Formula I, for example, wherein any Z, W, X, and/or Y independently comprises a sulphur atom, can be present in one or both oligonucleotide strands of the siNA duplex, for example, in the sense strand, the antisense strand, or both strands. The siNA molecules of the invention can comprise one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) chemically-modified internucleotide linkages having Formula I at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the sense strand, the antisense strand, or both strands. For example, an exemplary siNA molecule of the invention can comprise about 1 to about 5 or more (*e.g.*, about 1, 2, 3, 4, 5, or more) chemically-modified internucleotide linkages having Formula I at the 5'-end of the sense strand, the antisense strand, or both strands. In another non-limiting example, an exemplary siNA molecule of the invention can comprise one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) pyrimidine nucleotides with chemically-modified internucleotide linkages having Formula I in the sense strand, the antisense strand, or both strands. In yet another non-limiting example, an exemplary siNA molecule of the invention can comprise one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) purine nucleotides with chemically-modified internucleotide linkages having Formula I in the sense strand, the antisense strand, or both strands. In another embodiment, a siNA molecule of the invention having internucleotide linkage(s) of Formula I also comprises a chemically-modified nucleotide or non-nucleotide having any of Formulae I-VII.

In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule capable of mediating RNA interference (RNAi) against a



VEGF and/or VEGFr inside a cell or reconstituted *in vitro* system, wherein the chemical modification comprises one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) nucleotides or non-nucleotides having Formula II:




- 5 wherein each R3, R4, R5, R6, R7, R8, R10, R11 and R12 is independently H, OH, alkyl, substituted alkyl, alkaryl or aralkyl, F, Cl, Br, CN, CF3, OCF3, OCN, O-alkyl, S-alkyl, N-alkyl, O-alkenyl, S-alkenyl, N-alkenyl, SO-alkyl, alkyl-OSH, alkyl-OH, O-alkyl-OH, O-alkyl-SH, S-alkyl-OH, S-alkyl-SH, alkyl-S-alkyl, alkyl-O-alkyl, ONO2, NO2, N3, NH2, aminoalkyl, aminoacid, aminoacyl, ONH2, O-aminoalkyl, O-aminoacid, O-aminoacyl,
- 10 heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, or group having Formula I; R9 is O, S, CH2, S=O, CHF, or CF2, and B is a nucleosidic base such as adenine, guanine, uracil, cytosine, thymine, 2-aminoadenosine, 5-methylcytosine, 2,6-diaminopurine, or any other non-naturally occurring base that can be complementary or non-complementary to target RNA or a non-nucleosidic base such as phenyl, naphthyl, 3-
- 15 nitropyrrole, 5-nitroindole, nebularine, pyridone, pyridinone, or any other non-naturally occurring universal base that can be complementary or non-complementary to target RNA.

- The chemically-modified nucleotide or non-nucleotide of Formula II can be present in one or both oligonucleotide strands of the siNA duplex, for example in the sense strand, the antisense strand, or both strands. The siNA molecules of the invention can comprise one or
- 20 more chemically-modified nucleotide or non-nucleotide of Formula II at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the sense strand, the antisense strand, or both strands. For example, an exemplary siNA molecule of the invention can comprise about 1 to about 5 or more (e.g., about 1, 2, 3, 4, 5, or more) chemically-modified nucleotides or non-nucleotides of Formula II at the 5'-end of the sense strand, the antisense strand, or both

strands. In another non-limiting example, an exemplary siRNA molecule of the invention can comprise about 1 to about 5 or more (e.g., about 1, 2, 3, 4, 5, or more) chemically-modified nucleotides or non-nucleotides of Formula II at the 3'-end of the sense strand, the antisense strand, or both strands.

- 5 In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siRNA) molecule capable of mediating RNA interference (RNAi) against a VEGF and/or VEGFR inside a cell or reconstituted *in vitro* system, wherein the chemical modification comprises one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) nucleotides or non-nucleotides having Formula III:




10

- wherein each R3, R4, R5, R6, R7, R8, R10, R11 and R12 is independently H, OH, alkyl, substituted alkyl, alkaryl or aralkyl, F, Cl, Br, CN, CF3, OCF3, OCN, O-alkyl, S-alkyl, N-alkyl, O-alkenyl, S-alkenyl, N-alkenyl, SO-alkyl, alkyl-OSH, alkyl-OH, O-alkyl-OH, O-alkyl-SH, S-alkyl-OH, S-alkyl-SH, alkyl-S-alkyl, alkyl-O-alkyl, ONO2, NO2, N3, NH2, aminoalkyl, aminoacid, aminoacyl, ONH2, O-aminoalkyl, O-aminoacid, O-aminoacyl, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, or group having Formula I; R9 is O, S, CH2, S=O, CHF, or CF2, and B is a nucleosidic base such as adenine, guanine, uracil, cytosine, thymine, 2-aminoadenosine, 5-methylcytosine, 2,6-diaminopurine, or any other non-naturally occurring base that can be employed to be complementary or non-complementary to target RNA or a non-nucleosidic base such as phenyl, naphthyl, 3-nitropyrrole, 5-nitroindole, nebularine, pyridone, pyridinone, or any other non-naturally occurring universal base that can be complementary or non-complementary to target RNA.
- 15
- 20

The chemically-modified nucleotide or non-nucleotide of Formula III can be present in one or both oligonucleotide strands of the siNA duplex, for example, in the sense strand, the antisense strand, or both strands. The siNA molecules of the invention can comprise one or more chemically-modified nucleotide or non-nucleotide of Formula III at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the sense strand, the antisense strand, or both strands. For example, an exemplary siNA molecule of the invention can comprise about 1 to about 5 or more (e.g., about 1, 2, 3, 4, 5, or more) chemically-modified nucleotide(s) or non-nucleotide(s) of Formula III at the 5'-end of the sense strand, the antisense strand, or both strands. In another non-limiting example, an exemplary siNA molecule of the invention can comprise about 1 to about 5 or more (e.g., about 1, 2, 3, 4, 5, or more) chemically-modified nucleotide or non-nucleotide of Formula III at the 3'-end of the sense strand, the antisense strand, or both strands.

In another embodiment, a siNA molecule of the invention comprises a nucleotide having Formula II or III, wherein the nucleotide having Formula II or III is in an inverted configuration. For example, the nucleotide having Formula II or III is connected to the siNA construct in a 3'-3', 3'-2', 2'-3', or 5'-5' configuration, such as at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of one or both siNA strands.

In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule capable of mediating RNA interference (RNAi) against a VEGF and/or VEGFr inside a cell or reconstituted *in vitro* system, wherein the chemical modification comprises a 5'-terminal phosphate group having Formula IV:



wherein each X and Y is independently O, S, N, alkyl, substituted alkyl, or alkylhalo; wherein each Z and W is independently O, S, N, alkyl, substituted alkyl, O-alkyl, S-alkyl, alkaryl, aralkyl, or alkylhalo; and wherein W, X, Y and Z are not all O.

In one embodiment, the invention features a siNA molecule having a 5'-terminal phosphate group having Formula IV on the target-complementary strand, for example, a strand complementary to a target RNA, wherein the siNA molecule comprises an all RNA siNA molecule. In another embodiment, the invention features a siNA molecule having a 5'-terminal phosphate group having Formula IV on the target-complementary strand wherein the siNA molecule also comprises about 1 to about 3 (e.g., about 1, 2, or 3) nucleotide 3'-terminal nucleotide overhangs having about 1 to about 4 (e.g., about 1, 2, 3, or 4) deoxyribonucleotides on the 3'-end of one or both strands. In another embodiment, a 5'-terminal phosphate group having Formula IV is present on the target-complementary strand of a siNA molecule of the invention, for example a siNA molecule having chemical modifications having any of Formulae I-VII.

In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule capable of mediating RNA interference (RNAi) against a VEGF and/or VEGFR inside a cell or reconstituted *in vitro* system, wherein the chemical modification comprises one or more phosphorothioate internucleotide linkages. For example, in a non-limiting example, the invention features a chemically-modified short interfering nucleic acid (siNA) having about 1, 2, 3, 4, 5, 6, 7, 8 or more phosphorothioate internucleotide linkages in one siNA strand. In yet another embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) individually having about 1, 2, 3, 4, 5, 6, 7, 8 or more phosphorothioate internucleotide linkages in both siNA strands. The phosphorothioate internucleotide linkages can be present in one or both oligonucleotide strands of the siNA duplex, for example in the sense strand, the antisense strand, or both strands. The siNA molecules of the invention can comprise one or more phosphorothioate internucleotide linkages at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the sense strand, the antisense strand, or both strands. For example, an exemplary siNA molecule of the invention can comprise about 1 to about 5 or more (e.g., about 1, 2, 3, 4, 5, or more) consecutive phosphorothioate internucleotide linkages at the 5'-end of the sense strand, the antisense strand, or both strands. In another non-limiting example, an exemplary siNA molecule of the invention can comprise one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) pyrimidine phosphorothioate internucleotide linkages in the sense

strand, the antisense strand, or both strands. In yet another non-limiting example, an exemplary siNA molecule of the invention can comprise one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) purine phosphorothioate internucleotide linkages in the sense strand, the antisense strand, or both strands.

5 In one embodiment, the invention features a siNA molecule, wherein the sense strand comprises one or more, for example, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more phosphorothioate internucleotide linkages, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or about one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally  
10 a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the sense strand; and wherein the antisense strand comprises about 1 to about 10 or more, specifically about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more phosphorothioate internucleotide linkages, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base  
15 modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the antisense strand. In another embodiment, one or more, for example about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more, pyrimidine nucleotides of the sense and/or antisense siNA strand are chemically-modified with 2'-deoxy, 2'-O-methyl and/or 2'-deoxy-2'-fluoro nucleotides, with or without one or more, for example about 1, 2, 3, 4, 5, 6,  
20 7, 8, 9, 10, or more, phosphorothioate internucleotide linkages and/or a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends, being present in the same or different strand.

In another embodiment, the invention features a siNA molecule, wherein the sense strand comprises about 1 to about 5, specifically about 1, 2, 3, 4, or 5 phosphorothioate  
25 internucleotide linkages, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the sense strand; and wherein the antisense strand comprises about 1 to about 5 or more, specifically about 1, 2, 3, 4, 5, or more  
30 phosphorothioate internucleotide linkages, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7,

8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the antisense strand. In another embodiment, one or more, for example about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 5 or more, pyrimidine nucleotides of the sense and/or antisense siNA strand are chemically-modified with 2'-deoxy, 2'-O-methyl and/or 2'-deoxy-2'-fluoro nucleotides, with or without about 1 to about 5 or more, for example about 1, 2, 3, 4, 5, or more phosphorothioate internucleotide linkages and/or a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends, being present in the same or different strand.

10 In one embodiment, the invention features a siNA molecule, wherein the antisense strand comprises one or more, for example, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more phosphorothioate internucleotide linkages, and/or about one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally 15 a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the sense strand; and wherein the antisense strand comprises about 1 to about 10 or more, specifically about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more phosphorothioate internucleotide linkages, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base 20 modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the antisense strand. In another embodiment, one or more, for example about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more pyrimidine nucleotides of the sense and/or antisense siNA strand are chemically-modified with 2'-deoxy, 2'-O-methyl and/or 2'-deoxy-2'-fluoro nucleotides, with or without one or more, for example, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 25 10 or more phosphorothioate internucleotide linkages and/or a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3' and 5'-ends, being present in the same or different strand.

In another embodiment, the invention features a siNA molecule, wherein the antisense strand comprises about 1 to about 5 or more, specifically about 1, 2, 3, 4, 5 or more phosphorothioate internucleotide linkages, and/or one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 30 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (*e.g.*, about

1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the sense strand; and wherein the antisense strand comprises about 1 to about 5 or more, specifically about 1, 2, 3, 4, 5 or more phosphorothioate internucleotide linkages, and/or one or more  
5 (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the antisense strand. In another embodiment, one or more, for example about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more pyrimidine nucleotides of the sense and/or antisense  
10 siNA strand are chemically-modified with 2'-deoxy, 2'-O-methyl and/or 2'-deoxy-2'-fluoro nucleotides, with or without about 1 to about 5, for example about 1, 2, 3, 4, 5 or more phosphorothioate internucleotide linkages and/or a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends, being present in the same or different strand.

In one embodiment, the invention features a chemically-modified short interfering  
15 nucleic acid (siNA) molecule having about 1 to about 5, specifically about 1, 2, 3, 4, 5 or more phosphorothioate internucleotide linkages in each strand of the siNA molecule.

In another embodiment, the invention features a siNA molecule comprising 2'-5' internucleotide linkages. The 2'-5' internucleotide linkage(s) can be at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of one or both siNA sequence strands. In addition, the 2'-5'  
20 internucleotide linkage(s) can be present at various other positions within one or both siNA sequence strands, for example, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more including every internucleotide linkage of a pyrimidine nucleotide in one or both strands of the siNA molecule can comprise a 2'-5' internucleotide linkage, or about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more including every internucleotide linkage of a purine nucleotide in one or both strands of  
25 the siNA molecule can comprise a 2'-5' internucleotide linkage.

In another embodiment, a chemically-modified siNA molecule of the invention comprises a duplex having two strands, one or both of which can be chemically-modified, wherein each strand is about 18 to about 27 (e.g., about 18, 19, 20, 21, 22, 23, 24, 25, 26, or 27) nucleotides in length, wherein the duplex has about 18 to about 23 (e.g., about 18, 19,

20, 21, 22, or 23) base pairs, and wherein the chemical modification comprises a structure having any of Formulae I-VII. For example, an exemplary chemically-modified siNA molecule of the invention comprises a duplex having two strands, one or both of which can be chemically-modified with a chemical modification having any of Formulae I-VII or any combination thereof, wherein each strand consists of about 21 nucleotides, each having a 2-nucleotide 3'-terminal nucleotide overhang, and wherein the duplex has about 19 base pairs. In another embodiment, a siNA molecule of the invention comprises a single stranded hairpin structure, wherein the siNA is about 36 to about 70 (e.g., about 36, 40, 45, 50, 55, 60, 65, or 70) nucleotides in length having about 18 to about 23 (e.g., about 18, 19, 20, 21, 22, or 23) base pairs, and wherein the siNA can include a chemical modification comprising a structure having any of Formulae I-VII or any combination thereof. For example, an exemplary chemically-modified siNA molecule of the invention comprises a linear oligonucleotide having about 42 to about 50 (e.g., about 42, 43, 44, 45, 46, 47, 48, 49, or 50) nucleotides that is chemically-modified with a chemical modification having any of Formulae I-VII or any combination thereof, wherein the linear oligonucleotide forms a hairpin structure having about 19 base pairs and a 2-nucleotide 3'-terminal nucleotide overhang. In another embodiment, a linear hairpin siNA molecule of the invention contains a stem loop motif, wherein the loop portion of the siNA molecule is biodegradable. For example, a linear hairpin siNA molecule of the invention is designed such that degradation of the loop portion of the siNA molecule *in vivo* can generate a double-stranded siNA molecule with 3'-terminal overhangs, such as 3'-terminal nucleotide overhangs comprising about 2 nucleotides.


In another embodiment, a siNA molecule of the invention comprises a circular nucleic acid molecule, wherein the siNA is about 38 to about 70 (e.g., about 38, 40, 45, 50, 55, 60, 65, or 70) nucleotides in length having about 18 to about 23 (e.g., about 18, 19, 20, 21, 22, or 23) base pairs, and wherein the siNA can include a chemical modification, which comprises a structure having any of Formulae I-VII or any combination thereof. For example, an exemplary chemically-modified siNA molecule of the invention comprises a circular oligonucleotide having about 42 to about 50 (e.g., about 42, 43, 44, 45, 46, 47, 48, 49, or 50) nucleotides that is chemically-modified with a chemical modification having any



of Formulae I-VII or any combination thereof, wherein the circular oligonucleotide forms a dumbbell shaped structure having about 19 base pairs and 2 loops.


In another embodiment, a circular siNA molecule of the invention contains two loop motifs, wherein one or both loop portions of the siNA molecule is biodegradable. For example, a circular siNA molecule of the invention is designed such that degradation of the loop portions of the siNA molecule *in vivo* can generate a double-stranded siNA molecule with 3'-terminal overhangs, such as 3'-terminal nucleotide overhangs comprising about 2 nucleotides.

In one embodiment, a siNA molecule of the invention comprises at least one (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) abasic moiety, for example a compound having Formula V:




wherein each R3, R4, R5, R6, R7, R8, R10, R11, R12, and R13 is independently H, OH, alkyl, substituted alkyl, alkaryl or aralkyl, F, Cl, Br, CN, CF3, OCF3, OCN, O-alkyl, S-alkyl, N-alkyl, O-alkenyl, S-alkenyl, N-alkenyl, SO-alkyl, alkyl-OSH, alkyl-OH, O-alkyl-OH, O-alkyl-SH, S-alkyl-OH, S-alkyl-SH, alkyl-S-alkyl, alkyl-O-alkyl, ONO2, NO2, N3, NH2, aminoalkyl, aminoacid, aminoacyl, ONH2, O-aminoalkyl, O-aminoacid, O-aminoacyl, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, or group having Formula I; R9 is O, S, CH2, S=O, CHF, or CF2.

In one embodiment, a siNA molecule of the invention comprises at least one (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) inverted abasic moiety, for example a compound having Formula VI:



wherein each R3, R4, R5, R6, R7, R8, R10, R11, R12, and R13 is independently H, OH, alkyl, substituted alkyl, alkaryl or aralkyl, F, Cl, Br, CN, CF3, OCF3, OCN, O-alkyl, S-alkyl, N-alkyl, O-alkenyl, S-alkenyl, N-alkenyl, SO-alkyl, alkyl-OSH, alkyl-OH, O-alkyl-OH, O-alkyl-SH, S-alkyl-OH, S-alkyl-SH, alkyl-S-alkyl, alkyl-O-alkyl, ONO2, NO2, N3, NH2, aminoalkyl, aminoacid, aminoacyl, ONH2, O-aminoalkyl, O-aminoacid, O-aminoacyl, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, or group having Formula I; R9 is O, S, CH2, S=O, CHF, or CF2, and either R2, R3, R8 or R13 serve as points of attachment to the siNA molecule of the invention.

- 10 In another embodiment, a siNA molecule of the invention comprises at least one (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) substituted polyalkyl moieties, for example a compound having Formula VII:



wherein each n is independently an integer from 1 to 12, each R1, R2 and R3 is independently H, OH, alkyl, substituted alkyl, alkaryl or aralkyl, F, Cl, Br, CN, CF3, OCF3, OCN, O-alkyl, S-alkyl, N-alkyl, O-alkenyl, S-alkenyl, N-alkenyl, SO-alkyl, alkyl-OSH, alkyl-OH, O-alkyl-OH, O-alkyl-SH, S-alkyl-OH, S-alkyl-SH, alkyl-S-alkyl, alkyl-O-alkyl, ONO2, NO2, N3, NH2, aminoalkyl, aminoacid, aminoacyl, ONH2, O-aminoalkyl, O-aminoacid, O-aminoacyl, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, or a group having Formula I, and R1, R2 or R3 serves as points of attachment to the siNA molecule of the invention.

In another embodiment, the invention features a compound having Formula VII, wherein R1 and R2 are hydroxyl (OH) groups,  $n = 1$ , and R3 comprises O and is the point of attachment to the 3'-end, the 5'-end, or both of the 3' and 5'-ends of one or both strands of a double-stranded siNA molecule of the invention or to a single-stranded siNA molecule of the invention. This modification is referred to herein as "glyceryl" (for example modification 6 in Figure 10).

In another embodiment, a moiety having any of Formula V, VI or VII of the invention is at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of a siNA molecule of the invention. For example, a moiety having Formula V, VI or VII can be present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense strand, the sense strand, or both antisense and sense strands of the siNA molecule. In addition, a moiety having Formula VII can be present at the 3'-end or the 5'-end of a hairpin siNA molecule as described herein.

In another embodiment, a siNA molecule of the invention comprises an abasic residue having Formula V or VI, wherein the abasic residue having Formula VI or VI is connected to the siNA construct in a 3'-3', 3'-2', 2'-3', or 5'-5' configuration, such as at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of one or both siNA strands.

In one embodiment, a siNA molecule of the invention comprises one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) locked nucleic acid (LNA) nucleotides, for example at the 5'-end, the 3'-end, both of the 5' and 3'-ends, or any combination thereof, of the siNA molecule.

In another embodiment, a siNA molecule of the invention comprises one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) acyclic nucleotides, for example at the 5'-end, the 3'-end, both of the 5' and 3'-ends, or any combination thereof, of the siNA molecule.

In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule of the invention, wherein the chemically-modified siNA comprises a sense region, where any (*e.g.*, one or more or all) pyrimidine nucleotides present in the sense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (*e.g.*, wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a

plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and where any (*e.g.*, one or more or all) purine nucleotides present in the sense region are 2'-deoxy purine nucleotides (*e.g.*, wherein all purine nucleotides are 2'-deoxy purine nucleotides or alternately a plurality of purine nucleotides are 2'-deoxy purine nucleotides).

5        In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule of the invention, wherein the chemically-modified siNA comprises a sense region, where any (*e.g.*, one or more or all) pyrimidine nucleotides present in the sense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (*e.g.*, wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a  
10    plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and where any (*e.g.*, one or more or all) purine nucleotides present in the sense region are 2'-deoxy purine nucleotides (*e.g.*, wherein all purine nucleotides are 2'-deoxy purine nucleotides or alternately a plurality of purine nucleotides are 2'-deoxy purine nucleotides), wherein any nucleotides comprising a 3'-terminal nucleotide overhang that are present in  
15    said sense region are 2'-deoxy nucleotides.

      In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule of the invention, wherein the chemically-modified siNA comprises an antisense region, where any (*e.g.*, one or more or all) pyrimidine nucleotides present in the antisense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (*e.g.*, wherein  
20    all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein any (*e.g.*, one or more or all) purine nucleotides present in the antisense region are 2'-O-methyl purine nucleotides (*e.g.*, wherein all purine nucleotides are 2'-O-methyl purine nucleotides or alternately a plurality of purine nucleotides are 2'-O-methyl purine  
25    nucleotides).

      In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule of the invention, wherein the chemically-modified siNA comprises an antisense region, where any (*e.g.*, one or more or all) pyrimidine nucleotides present in the antisense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (*e.g.*, wherein

all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein any (e.g., one or more or all) purine nucleotides present in the antisense region are 2'-O-methyl purine nucleotides (e.g., wherein all purine nucleotides are 2'-O-methyl purine nucleotides or alternately a plurality of purine nucleotides are 2'-O-methyl purine nucleotides), wherein any nucleotides comprising a 3'-terminal nucleotide overhang that are present in said antisense region are 2'-deoxy nucleotides.

In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule of the invention, wherein the chemically-modified siNA comprises an antisense region, where any (e.g., one or more or all) pyrimidine nucleotides present in the antisense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and where any (e.g., one or more or all) purine nucleotides present in the antisense region are 2'-deoxy purine nucleotides (e.g., wherein all purine nucleotides are 2'-deoxy purine nucleotides or alternately a plurality of purine nucleotides are 2'-deoxy purine nucleotides).

In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule of the invention capable of mediating RNA interference (RNAi) against a VEGF and/or VEGFr inside a cell or reconstituted *in vitro* system, wherein the chemically-modified siNA comprises a sense region, where one or more pyrimidine nucleotides present in the sense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and where one or more purine nucleotides present in the sense region are 2'-deoxy purine nucleotides (e.g., wherein all purine nucleotides are 2'-deoxy purine nucleotides or alternately a plurality of purine nucleotides are 2'-deoxy purine nucleotides), and inverted deoxy abasic modifications that are optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the sense region, the sense region optionally further comprising a 3'-terminal overhang having about 1 to about 4 (e.g., about 1, 2, 3, or 4) 2'-deoxyribonucleotides; and wherein the chemically-modified short interfering nucleic acid

molecule comprises an antisense region, where one or more pyrimidine nucleotides present in the antisense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and  
5 wherein one or more purine nucleotides present in the antisense region are 2'-O-methyl purine nucleotides (e.g., wherein all purine nucleotides are 2'-O-methyl purine nucleotides or alternately a plurality of purine nucleotides are 2'-O-methyl purine nucleotides), and a terminal cap modification, such as any modification described herein or shown in Figure 10, that is optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the  
10 antisense sequence, the antisense region optionally further comprising a 3'-terminal nucleotide overhang having about 1 to about 4 (e.g., about 1, 2, 3, or 4) 2'-deoxynucleotides, wherein the overhang nucleotides can further comprise one or more (e.g., 1, 2, 3, or 4) phosphorothioate internucleotide linkages. Non-limiting examples of these chemically-modified siNAs are shown in Figures 4 and 5 and Tables III and IV herein.

15 In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule of the invention capable of mediating RNA interference (RNAi) against a VEGF and/or VEGFr inside a cell or reconstituted *in vitro* system, wherein the siNA comprises a sense region, where one or more pyrimidine nucleotides present in the sense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine  
20 nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and where one or more purine nucleotides present in the sense region are purine ribonucleotides (e.g., wherein all purine nucleotides are purine ribonucleotides or alternately a plurality of purine nucleotides are purine ribonucleotides), and inverted deoxy abasic modifications that are  
25 optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the sense region, the sense region optionally further comprising a 3'-terminal overhang having about 1 to about 4 (e.g., about 1, 2, 3, or 4) 2'-deoxyribonucleotides; and wherein the siNA comprises an antisense region, where one or more pyrimidine nucleotides present in the antisense  
30 nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of

pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein any purine nucleotides present in the antisense region are 2'-O-methyl purine nucleotides (e.g., wherein all purine nucleotides are 2'-O-methyl purine nucleotides or alternately a plurality of purine nucleotides are 2'-O-methyl purine nucleotides), and a terminal cap modification, such as any modification described herein or shown in Figure 10, that is optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense sequence, the antisense region optionally further comprising a 3'-terminal nucleotide overhang having about 1 to about 4 (e.g., about 1, 2, 3, or 4) 2'-deoxynucleotides, wherein the overhang nucleotides can further comprise one or more (e.g., 1, 2, 3, or 4) phosphorothioate internucleotide linkages. Non-limiting examples of these chemically-modified siNAs are shown in Figures 4 and 5 and Tables III and IV herein.

In one embodiment, the invention features a chemically-modified short interfering nucleic acid (siNA) molecule of the invention capable of mediating RNA interference (RNAi) against a VEGF and/or VEGFr inside a cell or reconstituted *in vitro* system, wherein the chemically-modified siNA comprises a sense region, where one or more pyrimidine nucleotides present in the sense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and for example where one or more purine nucleotides present in the sense region are selected from the group consisting of 2'-deoxy nucleotides, locked nucleic acid (LNA) nucleotides, 2'-methoxyethyl nucleotides, 4'-thionucleotides, and 2'-O-methyl nucleotides (e.g., wherein all purine nucleotides are selected from the group consisting of 2'-deoxy nucleotides, locked nucleic acid (LNA) nucleotides, 2'-methoxyethyl nucleotides, 4'-thionucleotides, and 2'-O-methyl nucleotides or alternately a plurality of purine nucleotides are selected from the group consisting of 2'-deoxy nucleotides, locked nucleic acid (LNA) nucleotides, 2'-methoxyethyl nucleotides, 4'-thionucleotides, and 2'-O-methyl nucleotides), and wherein inverted deoxy abasic modifications are optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the sense region, the sense region optionally further comprising a 3'-terminal overhang having about 1 to about 4 (e.g., about 1, 2, 3, or 4) 2'-deoxyribonucleotides; and wherein the chemically-modified short interfering nucleic acid

molecule comprises an antisense region, where one or more pyrimidine nucleotides present in the antisense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein one or more purine nucleotides present in the antisense region are selected from the group consisting of 2'-deoxy nucleotides, locked nucleic acid (LNA) nucleotides, 2'-methoxyethyl nucleotides, 4'-thionucleotides, and 2'-O-methyl nucleotides (e.g., wherein all purine nucleotides are selected from the group consisting of 2'-deoxy nucleotides, locked nucleic acid (LNA) nucleotides, 2'-methoxyethyl nucleotides, 4'-thionucleotides, and 2'-O-methyl nucleotides or alternately a plurality of purine nucleotides are selected from the group consisting of 2'-deoxy nucleotides, locked nucleic acid (LNA) nucleotides, 2'-methoxyethyl nucleotides, 4'-thionucleotides, and 2'-O-methyl nucleotides), and a terminal cap modification, such as any modification described herein or shown in Figure 10, that is optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense sequence, the antisense region optionally further comprising a 3'-terminal nucleotide overhang having about 1 to about 4 (e.g., about 1, 2, 3, or 4) 2'-deoxynucleotides, wherein the overhang nucleotides can further comprise one or more (e.g., 1, 2, 3, or 4) phosphorothioate internucleotide linkages.

In another embodiment, any modified nucleotides present in the siNA molecules of the invention, preferably in the antisense strand of the siNA molecules of the invention, but also optionally in the sense and/or both antisense and sense strands, comprise modified nucleotides having properties or characteristics similar to naturally occurring ribonucleotides. For example, the invention features siNA molecules including modified nucleotides having a Northern conformation (e.g., Northern pseudorotation cycle, see for example Saenger, *Principles of Nucleic Acid Structure*, Springer-Verlag ed., 1984). As such, chemically modified nucleotides present in the siNA molecules of the invention, preferably in the antisense strand of the siNA molecules of the invention, but also optionally in the sense and/or both antisense and sense strands, are resistant to nuclease degradation while at the same time maintaining the capacity to mediate RNAi. Non-limiting examples of nucleotides having a northern configuration include locked nucleic acid (LNA)



nucleotides (e.g., 2'-O, 4'-C-methylene-(D-ribofuranosyl) nucleotides); 2'-methoxyethoxy (MOE) nucleotides; 2'-methyl-thio-ethyl, 2'-deoxy-2'-fluoro nucleotides, 2'-deoxy-2'-chloro nucleotides, 2'-azido nucleotides, and 2'-O-methyl nucleotides.

In one embodiment, the invention features a chemically-modified short interfering  
5 nucleic acid molecule (siNA) capable of mediating RNA interference (RNAi) against a VEGF and/or VEGFR inside a cell or reconstituted *in vitro* system, wherein the chemical modification comprises a conjugate covalently attached to the chemically-modified siNA molecule. In another embodiment, the conjugate is covalently attached to the chemically-modified siNA molecule via a biodegradable linker. In one embodiment, the conjugate  
10 molecule is attached at the 3'-end of either the sense strand, the antisense strand, or both strands of the chemically-modified siNA molecule. In another embodiment, the conjugate molecule is attached at the 5'-end of either the sense strand, the antisense strand, or both strands of the chemically-modified siNA molecule. In yet another embodiment, the conjugate molecule is attached both the 3'-end and 5'-end of either the sense strand, the  
15 antisense strand, or both strands of the chemically-modified siNA molecule, or any combination thereof. In one embodiment, a conjugate molecule of the invention comprises a molecule that facilitates delivery of a chemically-modified siNA molecule into a biological system, such as a cell. In another embodiment, the conjugate molecule attached to the chemically-modified siNA molecule is a poly ethylene glycol, human serum albumin, or a  
20 ligand for a cellular receptor that can mediate cellular uptake. Examples of specific conjugate molecules contemplated by the instant invention that can be attached to chemically-modified siNA molecules are described in Vargeese *et al.*, U.S. Serial No. 10/201,394, incorporated by reference herein. The type of conjugates used and the extent of conjugation of siNA molecules of the invention can be evaluated for improved  
25 pharmacokinetic profiles, bioavailability, and/or stability of siNA constructs while at the same time maintaining the ability of the siNA to mediate RNAi activity. As such, one skilled in the art can screen siNA constructs that are modified with various conjugates to determine whether the siNA conjugate complex possesses improved properties while maintaining the ability to mediate RNAi, for example in animal models as are generally  
30 known in the art.

In one embodiment, the invention features a short interfering nucleic acid (siNA) molecule of the invention, wherein the siNA further comprises a nucleotide, non-nucleotide, or mixed nucleotide/non-nucleotide linker that joins the sense region of the siNA to the antisense region of the siNA. In one embodiment, a nucleotide linker of the invention can be a linker of  $\geq 2$  nucleotides in length, for example 3, 4, 5, 6, 7, 8, 9, or 10 nucleotides in length. In another embodiment, the nucleotide linker can be a nucleic acid aptamer. By "aptamer" or "nucleic acid aptamer" as used herein is meant a nucleic acid molecule that binds specifically to a target molecule wherein the nucleic acid molecule has sequence that comprises a sequence recognized by the target molecule in its natural setting. Alternately, an aptamer can be a nucleic acid molecule that binds to a target molecule where the target molecule does not naturally bind to a nucleic acid. The target molecule can be any molecule of interest. For example, the aptamer can be used to bind to a ligand-binding domain of a protein, thereby preventing interaction of the naturally occurring ligand with the protein. This is a non-limiting example and those in the art will recognize that other embodiments can be readily generated using techniques generally known in the art. (See, for example, Gold *et al.*, 1995, *Annu. Rev. Biochem.*, 64, 763; Brody and Gold, 2000, *J. Biotechnol.*, 74, 5; Sun, 2000, *Curr. Opin. Mol. Ther.*, 2, 100; Kusser, 2000, *J. Biotechnol.*, 74, 27; Hermann and Patel, 2000, *Science*, 287, 820; and Jayasena, 1999, *Clinical Chemistry*, 45, 1628.)

In yet another embodiment, a non-nucleotide linker of the invention comprises abasic nucleotide, polyether, polyamine, polyamide, peptide, carbohydrate, lipid, polyhydrocarbon, or other polymeric compounds (e.g. polyethylene glycols such as those having between 2 and 100 ethylene glycol units). Specific examples include those described by Seela and Kaiser, *Nucleic Acids Res.* 1990, 18:6353 and *Nucleic Acids Res.* 1987, 15:3113; Cload and Schepartz, *J. Am. Chem. Soc.* 1991, 113:6324; Richardson and Schepartz, *J. Am. Chem. Soc.* 1991, 113:5109; Ma *et al.*, *Nucleic Acids Res.* 1993, 21:2585 and *Biochemistry* 1993, 32:1751; Durand *et al.*, *Nucleic Acids Res.* 1990, 18:6353; McCurdy *et al.*, *Nucleosides & Nucleotides* 1991, 10:287; Jsckke *et al.*, *Tetrahedron Lett.* 1993, 34:301; Ono *et al.*, *Biochemistry* 1991, 30:9914; Arnold *et al.*, International Publication No. WO 89/02439; Usman *et al.*, International Publication No. WO 95/06731; Dudycz *et al.*, International Publication No. WO 95/11910 and Ferentz and Verdine, *J. Am. Chem. Soc.* 1991, 113:4000,

all hereby incorporated by reference herein. A "non-nucleotide" further means any group or compound that can be incorporated into a nucleic acid chain in the place of one or more nucleotide units, including either sugar and/or phosphate substitutions, and allows the remaining bases to exhibit their enzymatic activity. The group or compound can be abasic  
5 in that it does not contain a commonly recognized nucleotide base, such as adenosine, guanine, cytosine, uracil or thymine, for example at the C1 position of the sugar.

In one embodiment, the invention features a short interfering nucleic acid (siNA) molecule capable of mediating RNA interference (RNAi) inside a cell or reconstituted in vitro system, wherein one or both strands of the siNA molecule that are assembled from two  
10 separate oligonucleotides do not comprise any ribonucleotides. For example, a siNA molecule can be assembled from a single oligonucleotide where the sense and antisense regions of the siNA comprise separate oligonucleotides not having any ribonucleotides (e.g., nucleotides having a 2'-OH group) present in the oligonucleotides. In another example, a siNA molecule can be assembled from a single oligonucleotide where the sense and  
15 antisense regions of the siNA are linked or circularized by a nucleotide or non-nucleotide linker as described herein, wherein the oligonucleotide does not have any ribonucleotides (e.g., nucleotides having a 2'-OH group) present in the oligonucleotide. Applicant has surprisingly found that the presense of ribonucleotides (e.g., nucleotides having a 2'-hydroxyl group) within the siNA molecule is not required or essential to support RNAi  
20 activity. As such, in one embodiment, all positions within the siNA can include chemically modified nucleotides and/or non-nucleotides such as nucleotides and or non-nucleotides having Formula I, II, III, IV, V, VI, or VII or any combination thereof to the extent that the ability of the siNA molecule to support RNAi activity in a cell is maintained.

In one embodiment, a siNA molecule of the invention is a single stranded siNA  
25 molecule that mediates RNAi activity in a cell or reconstituted in vitro system, wherein the siNA molecule comprises a single stranded polynucleotide having complementarity to a target nucleic acid sequence. In another embodiment, the single stranded siNA molecule of the invention comprises a 5'-terminal phosphate group. In another embodiment, the single stranded siNA molecule of the invention comprises a 5'-terminal phosphate group and a 3'-  
30 terminal phosphate group (e.g., a 2',3'-cyclic phosphate). In another embodiment, the single

stranded siNA molecule of the invention comprises about 19 to about 29 nucleotides. In yet another embodiment, the single stranded siNA molecule of the invention comprises one or more chemically modified nucleotides or non-nucleotides described herein. For example, all the positions within the siNA molecule can include chemically-modified nucleotides such as nucleotides having any of Formulae I-VII, or any combination thereof to the extent that the ability of the siNA molecule to support RNAi activity in a cell is maintained.

In one embodiment, a siNA molecule of the invention is a single stranded siNA molecule that mediates RNAi activity in a cell or reconstituted in vitro system, wherein the siNA molecule comprises a single stranded polynucleotide having complementarity to a target nucleic acid sequence, and wherein one or more pyrimidine nucleotides present in the siNA are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein any purine nucleotides present in the antisense region are 2'-O-methyl purine nucleotides (e.g., wherein all purine nucleotides are 2'-O-methyl purine nucleotides or alternately a plurality of purine nucleotides are 2'-O-methyl purine nucleotides), and a terminal cap modification, such as any modification described herein or shown in Figure 10, that is optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense sequence, the siNA optionally further comprising about 1 to about 4 (e.g., about 1, 2, 3, or 4) terminal 2'-deoxynucleotides at the 3'-end of the siNA molecule, wherein the terminal nucleotides can further comprise one or more (e.g., 1, 2, 3, or 4) phosphorothioate internucleotide linkages, and wherein the siNA optionally further comprises a terminal phosphate group, such as a 5'-terminal phosphate group.

In one embodiment, a siNA molecule of the invention is a single stranded siNA molecule that mediates RNAi activity in a cell or reconstituted in vitro system, wherein the siNA molecule comprises a single stranded polynucleotide having complementarity to a target nucleic acid sequence, and wherein one or more pyrimidine nucleotides present in the siNA are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine

nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein any purine nucleotides present in the antisense region are 2'-deoxy purine nucleotides (e.g., wherein all purine nucleotides are 2'-deoxy purine nucleotides or alternately a plurality of purine nucleotides are 2'-deoxy purine nucleotides), and a terminal cap modification, such as any  
5 modification described herein or shown in Figure 10, that is optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense sequence, the siNA optionally further comprising about 1 to about 4 (e.g., about 1, 2, 3, or 4) terminal 2'-deoxynucleotides at the 3'-end of the siNA molecule, wherein the terminal nucleotides can further comprise one or more (e.g., 1, 2, 3, or 4 ) phosphorothioate internucleotide linkages, and wherein the  
10 siNA optionally further comprises a terminal phosphate group, such as a 5'-terminal phosphate group.

In one embodiment, a siNA molecule of the invention is a single stranded siNA molecule that mediates RNAi activity in a cell or reconstituted in vitro system, wherein the siNA molecule comprises a single stranded polynucleotide having complementarity to a  
15 target nucleic acid sequence, and wherein one or more pyrimidine nucleotides present in the siNA are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein any purine nucleotides present in the antisense region are locked nucleic acid (LNA) nucleotides (e.g.,  
20 wherein all purine nucleotides are LNA nucleotides or alternately a plurality of purine nucleotides are LNA nucleotides), and a terminal cap modification, such as any modification described herein or shown in Figure 10, that is optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense sequence, the siNA optionally further comprising about 1 to about 4 (e.g., about 1, 2, 3, or 4) terminal 2'-deoxynucleotides at the 3'-end of the  
25 siNA molecule, wherein the terminal nucleotides can further comprise one or more (e.g., 1, 2, 3, or 4 ) phosphorothioate internucleotide linkages, and wherein the siNA optionally further comprises a terminal phosphate group, such as a 5'-terminal phosphate group.

In one embodiment, a siNA molecule of the invention is a single stranded siNA molecule that mediates RNAi activity in a cell or reconstituted in vitro system, wherein the  
30 siNA molecule comprises a single stranded polynucleotide having complementarity to a

target nucleic acid sequence, and wherein one or more pyrimidine nucleotides present in the siNA are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein any purine nucleotides present in the antisense region are 2'-methoxyethyl purine nucleotides (e.g., wherein all purine nucleotides are 2'-methoxyethyl purine nucleotides or alternately a plurality of purine nucleotides are 2'-methoxyethyl purine nucleotides), and a terminal cap modification, such as any modification described herein or shown in Figure 10, that is optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense sequence, the siNA optionally further comprising about 1 to about 4 (e.g., about 1, 2, 3, or 4) terminal 2'-deoxynucleotides at the 3'-end of the siNA molecule, wherein the terminal nucleotides can further comprise one or more (e.g., 1, 2, 3, or 4 ) phosphorothioate internucleotide linkages, and wherein the siNA optionally further comprises a terminal phosphate group, such as a 5'-terminal phosphate group.

In another embodiment, any modified nucleotides present in the single stranded siNA molecules of the invention comprise modified nucleotides having properties or characteristics similar to naturally occurring ribonucleotides. For example, the invention features siNA molecules including modified nucleotides having a Northern conformation (e.g., Northern pseudorotation cycle, see for example Saenger, *Principles of Nucleic Acid Structure*, Springer-Verlag ed., 1984). As such, chemically modified nucleotides present in the single stranded siNA molecules of the invention are preferably resistant to nuclease degradation while at the same time maintaining the capacity to mediate RNAi.

In one embodiment, the invention features a method for modulating the expression of a VEGF and/or VEGFr gene within a cell comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein one of the siNA strands comprises a sequence complementary to RNA of the VEGF and/or VEGFr gene; and (b) introducing the siNA molecule into a cell under conditions suitable to modulate the expression of the VEGF and/or VEGFr gene in the cell.

In one embodiment, the invention features a method for modulating the expression of a VEGF and/or VEGFr gene within a cell comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein one of the siNA strands comprises a sequence complementary to RNA of the VEGF and/or VEGFr gene and  
5 wherein the sense strand sequence of the siNA comprises a sequence identical to the sequence of the target RNA; and (b) introducing the siNA molecule into a cell under conditions suitable to modulate the expression of the VEGF and/or VEGFr gene in the cell.

In another embodiment, the invention features a method for modulating the expression of more than one VEGF and/or VEGFr gene within a cell comprising: (a) synthesizing  
10 siNA molecules of the invention, which can be chemically-modified, wherein one of the siNA strands comprises a sequence complementary to RNA of the VEGF and/or VEGFr genes; and (b) introducing the siNA molecules into a cell under conditions suitable to modulate the expression of the VEGF and/or VEGFr genes in the cell.

In another embodiment, the invention features a method for modulating the expression of more than one VEGF and/or VEGFr gene within a cell comprising: (a) synthesizing a  
15 siNA molecule of the invention, which can be chemically-modified, wherein one of the siNA strands comprises a sequence complementary to RNA of the VEGF and/or VEGFr gene and wherein the sense strand sequence of the siNA comprises a sequence identical to the sequence of the target RNA; and (b) introducing the siNA molecules into a cell under  
20 conditions suitable to modulate the expression of the VEGF and/or VEGFr genes in the cell.

In one embodiment, siNA molecules of the invention are used as reagents in *ex vivo* applications. For example, siNA reagents are introduced into tissue or cells that are transplanted into a subject for therapeutic effect. The cells and/or tissue can be derived from an organism or subject that later receives the explant, or can be derived from another  
25 organism or subject prior to transplantation. The siNA molecules can be used to modulate the expression of one or more genes in the cells or tissue, such that the cells or tissue obtain a desired phenotype or are able to perform a function when transplanted in vivo. In one embodiment, certain target cells from a patient are extracted. These extracted cells are contacted with siNAs targeting a specific nucleotide sequence within the cells under

conditions suitable for uptake of the siNAs by these cells (e.g. using delivery reagents such as cationic lipids, liposomes and the like or using techniques such as electroporation to facilitate the delivery of siNAs into cells). The cells are then reintroduced back into the same patient or other patients. In one embodiment, the invention features a method of modulating the expression of a VEGF and/or VEGFr gene in a tissue explant comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein one of the siNA strands comprises a sequence complementary to RNA of the VEGF and/or VEGFr gene; and (b) introducing the siNA molecule into a cell of the tissue explant derived from a particular organism under conditions suitable to modulate the expression of the VEGF and/or VEGFr gene in the tissue explant. In another embodiment, the method further comprises introducing the tissue explant back into the organism the tissue was derived from or into another organism under conditions suitable to modulate the expression of the VEGF and/or VEGFr gene in that organism.

In one embodiment, the invention features a method of modulating the expression of a VEGF and/or VEGFr gene in a tissue explant comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein one of the siNA strands comprises a sequence complementary to RNA of the VEGF and/or VEGFr gene and wherein the sense strand sequence of the siNA comprises a sequence identical to the sequence of the target RNA; and (b) introducing the siNA molecule into a cell of the tissue explant derived from a particular organism under conditions suitable to modulate the expression of the VEGF and/or VEGFr gene in the tissue explant. In another embodiment, the method further comprises introducing the tissue explant back into the organism the tissue was derived from or into another organism under conditions suitable to modulate the expression of the VEGF and/or VEGFr gene in that organism.

In another embodiment, the invention features a method of modulating the expression of more than one VEGF and/or VEGFr gene in a tissue explant comprising: (a) synthesizing siNA molecules of the invention, which can be chemically-modified, wherein one of the siNA strands comprises a sequence complementary to RNA of the VEGF and/or VEGFr genes; and (b) introducing the siNA molecules into a cell of the tissue explant derived from a particular organism under conditions suitable to modulate the expression of the VEGF



and/or VEGFr genes in the tissue explant. In another embodiment, the method further comprises introducing the tissue explant back into the organism the tissue was derived from or into another organism under conditions suitable to modulate the expression of the VEGF and/or VEGFr genes in that organism.

5        In one embodiment, the invention features a method of modulating the expression of a VEGF and/or VEGFr gene in an organism comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein one of the siNA strands comprises a sequence complementary to RNA of the VEGF and/or VEGFr gene; and (b) introducing the siNA molecule into the organism under conditions suitable to modulate the  
10        expression of the VEGF and/or VEGFr gene in the organism.

      In another embodiment, the invention features a method of modulating the expression of more than one VEGF and/or VEGFr gene in an organism comprising: (a) synthesizing siNA molecules of the invention, which can be chemically-modified, wherein one of the siNA strands comprises a sequence complementary to RNA of the VEGF and/or VEGFr  
15        genes; and (b) introducing the siNA molecules into the organism under conditions suitable to modulate the expression of the VEGF and/or VEGFr genes in the organism.

      In one embodiment, the invention features a method for modulating the expression of a VEGF and/or VEGFr gene within a cell comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein the siNA comprises a single  
20        stranded sequence having complementarity to RNA of the VEGF and/or VEGFr gene; and (b) introducing the siNA molecule into a cell under conditions suitable to modulate the expression of the VEGF and/or VEGFr gene in the cell.

      In another embodiment, the invention features a method for modulating the expression of more than one VEGF and/or VEGFr gene within a cell comprising: (a) synthesizing  
25        siNA molecules of the invention, which can be chemically-modified, wherein the siNA comprises a single stranded sequence having complementarity to RNA of the VEGF and/or VEGFr gene; and (b) contacting the siNA molecule with a cell in vitro or in vivo under conditions suitable to modulate the expression of the VEGF and/or VEGFr genes in the cell.

In one embodiment, the invention features a method of modulating the expression of a VEGF and/or VEGFr gene in a tissue explant comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein the siNA comprises a single stranded sequence having complementarity to RNA of the VEGF and/or VEGFr gene; and  
5 (b) contacting the siNA molecule with a cell of the tissue explant derived from a particular organism under conditions suitable to modulate the expression of the VEGF and/or VEGFr gene in the tissue explant. In another embodiment, the method further comprises introducing the tissue explant back into the organism the tissue was derived from or into another organism under conditions suitable to modulate the expression of the VEGF and/or VEGFr  
10 gene in that organism.

In another embodiment, the invention features a method of modulating the expression of more than one VEGF and/or VEGFr gene in a tissue explant comprising: (a) synthesizing siNA molecules of the invention, which can be chemically-modified, wherein the siNA comprises a single stranded sequence having complementarity to RNA of the VEGF and/or  
15 VEGFr gene; and (b) introducing the siNA molecules into a cell of the tissue explant derived from a particular organism under conditions suitable to modulate the expression of the VEGF and/or VEGFr genes in the tissue explant. In another embodiment, the method further comprises introducing the tissue explant back into the organism the tissue was derived from or into another organism under conditions suitable to modulate the expression  
20 of the VEGF and/or VEGFr genes in that organism.

In one embodiment, the invention features a method of modulating the expression of a VEGF and/or VEGFr gene in an organism comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein the siNA comprises a single stranded sequence having complementarity to RNA of the VEGF and/or VEGFr gene; and  
25 (b) introducing the siNA molecule into the organism under conditions suitable to modulate the expression of the VEGF and/or VEGFr gene in the organism.

In another embodiment, the invention features a method of modulating the expression of more than one VEGF and/or VEGFr gene in an organism comprising: (a) synthesizing siNA molecules of the invention, which can be chemically-modified, wherein the siNA

comprises a single stranded sequence having complementarity to RNA of the VEGF and/or VEGFr gene; and (b) introducing the siNA molecules into the organism under conditions suitable to modulate the expression of the VEGF and/or VEGFr genes in the organism.

In one embodiment, the invention features a method of modulating the expression of a VEGF and/or VEGFr gene in an organism comprising contacting the organism with a siNA molecule of the invention under conditions suitable to modulate the expression of the VEGF and/or VEGFr gene in the organism.

In another embodiment, the invention features a method of modulating the expression of more than one VEGF and/or VEGFr gene in an organism comprising contacting the organism with one or more siNA molecules of the invention under conditions suitable to modulate the expression of the VEGF and/or VEGFr genes in the organism.

The siNA molecules of the invention can be designed to inhibit target (VEGF and/or VEGFr) gene expression through RNAi targeting of a variety of RNA molecules. In one embodiment, the siNA molecules of the invention are used to target various RNAs corresponding to a target gene. Non-limiting examples of such RNAs include messenger RNA (mRNA), alternate RNA splice variants of target gene(s), post-transcriptionally modified RNA of target gene(s), pre-mRNA of target gene(s), and/or RNA templates. If alternate splicing produces a family of transcripts that are distinguished by usage of appropriate exons, the instant invention can be used to inhibit gene expression through appropriate exons to specifically inhibit or to distinguish among the functions of gene family members. For example, a protein that contains an alternatively spliced transmembrane domain can be expressed in both membrane bound and secreted forms. Use of the invention to target the exon containing the transmembrane domain can be used to determine the functional consequences of pharmaceutical targeting of membrane bound as opposed to the secreted form of the protein. Non-limiting examples of applications of the invention relating to targeting these RNA molecules include therapeutic pharmaceutical applications, pharmaceutical discovery applications, molecular diagnostic and gene function applications, and gene mapping, for example using single nucleotide polymorphism mapping with siNA

molecules of the invention. Such applications can be implemented using known gene sequences or from partial sequences available from an expressed sequence tag (EST).

In another embodiment, the siNA molecules of the invention are used to target conserved sequences corresponding to a gene family or gene families such as VEGF and/or VEGFr family genes. As such, siNA molecules targeting multiple VEGF and/or VEGFr targets can provide increased therapeutic effect. In addition, siNA can be used to characterize pathways of gene function in a variety of applications. For example, the present invention can be used to inhibit the activity of target gene(s) in a pathway to determine the function of uncharacterized gene(s) in gene function analysis, mRNA function analysis, or translational analysis. The invention can be used to determine potential target gene pathways involved in various diseases and conditions toward pharmaceutical development. The invention can be used to understand pathways of gene expression involved in, for example, the progression and/or maintenance of cancer.

In one embodiment, siNA molecule(s) and/or methods of the invention are used to inhibit the expression of gene(s) that encode RNA referred to by Genbank Accession, for example VEGF and/or VEGFr genes encoding RNA sequence(s) referred to herein by Genbank Accession number, for example, Genbank Accession Nos. shown in Table I.

In one embodiment, the invention features a method comprising: (a) generating a library of siNA constructs having a predetermined complexity; and (b) assaying the siNA constructs of (a) above, under conditions suitable to determine RNAi target sites within the target RNA sequence. In another embodiment, the siNA molecules of (a) have strands of a fixed length, for example, about 23 nucleotides in length. In yet another embodiment, the siNA molecules of (a) are of differing length, for example having strands of about 19 to about 25 (e.g., about 19, 20, 21, 22, 23, 24, or 25) nucleotides in length. In one embodiment, the assay can comprise a reconstituted *in vitro* siNA assay as described herein. In another embodiment, the assay can comprise a cell culture system in which target RNA is expressed. In another embodiment, fragments of target RNA are analyzed for detectable levels of cleavage, for example by gel electrophoresis, northern blot analysis, or RNase protection assays, to determine the most suitable target site(s) within the target RNA

sequence. The target RNA sequence can be obtained as is known in the art, for example, by cloning and/or transcription for *in vitro* systems, and by cellular expression in *in vivo* systems.

In one embodiment, the invention features a method comprising: (a) generating a  
5 randomized library of siNA constructs having a predetermined complexity, such as of  $4^N$ ,  
where N represents the number of base paired nucleotides in each of the siNA construct  
strands (eg. for a siNA construct having 21 nucleotide sense and antisense strands with 19  
base pairs, the complexity would be  $4^{19}$ ); and (b) assaying the siNA constructs of (a) above,  
under conditions suitable to determine RNAi target sites within the target VEGF and/or  
10 VEGFr RNA sequence. In another embodiment, the siNA molecules of (a) have strands of a  
fixed length, for example about 23 nucleotides in length. In yet another embodiment, the  
siNA molecules of (a) are of differing length, for example having strands of about 19 to  
about 25 (e.g., about 19, 20, 21, 22, 23, 24, or 25) nucleotides in length. In one  
embodiment, the assay can comprise a reconstituted *in vitro* siNA assay as described in  
15 Example 7 herein. In another embodiment, the assay can comprise a cell culture system in  
which target RNA is expressed. In another embodiment, fragments of VEGF and/or VEGFr  
RNA are analyzed for detectable levels of cleavage, for example by gel electrophoresis,  
northern blot analysis, or RNase protection assays, to determine the most suitable target  
site(s) within the target VEGF and/or VEGFr RNA sequence. The target VEGF and/or  
20 VEGFr RNA sequence can be obtained as is known in the art, for example, by cloning  
and/or transcription for *in vitro* systems, and by cellular expression in *in vivo* systems.

In another embodiment, the invention features a method comprising: (a) analyzing the  
sequence of a RNA target encoded by a target gene; (b) synthesizing one or more sets of  
siNA molecules having sequence complementary to one or more regions of the RNA of (a);  
25 and (c) assaying the siNA molecules of (b) under conditions suitable to determine RNAi  
targets within the target RNA sequence. In one embodiment, the siNA molecules of (b)  
have strands of a fixed length, for example about 23 nucleotides in length. In another  
embodiment, the siNA molecules of (b) are of differing length, for example having strands  
of about 19 to about 25 (e.g., about 19, 20, 21, 22, 23, 24, or 25) nucleotides in length. In  
30 one embodiment, the assay can comprise a reconstituted *in vitro* siNA assay as described

herein. In another embodiment, the assay can comprise a cell culture system in which target RNA is expressed. Fragments of target RNA are analyzed for detectable levels of cleavage, for example by gel electrophoresis, northern blot analysis, or RNase protection assays, to determine the most suitable target site(s) within the target RNA sequence. The target RNA  
5 sequence can be obtained as is known in the art, for example, by cloning and/or transcription for *in vitro* systems, and by expression in *in vivo* systems.

By "target site" is meant a sequence within a target RNA that is "targeted" for cleavage mediated by a siNA construct which contains sequences within its antisense region that are complementary to the target sequence.

10 By "detectable level of cleavage" is meant cleavage of target RNA (and formation of cleaved product RNAs) to an extent sufficient to discern cleavage products above the background of RNAs produced by random degradation of the target RNA. Production of cleavage products from 1-5% of the target RNA is sufficient to detect above the background for most methods of detection.

15 In one embodiment, the invention features a composition comprising a siNA molecule of the invention, which can be chemically-modified, in a pharmaceutically acceptable carrier or diluent. In another embodiment, the invention features a pharmaceutical composition comprising siNA molecules of the invention, which can be chemically-modified, targeting one or more genes in a pharmaceutically acceptable carrier or diluent. In another  
20 embodiment, the invention features a method for treating or preventing a disease or condition in a subject, comprising administering to the subject a composition of the invention under conditions suitable for the treatment or prevention of the disease or condition in the subject, alone or in conjunction with one or more other therapeutic compounds. In yet another embodiment, the invention features a method for reducing or  
25 preventing tissue rejection in a subject comprising administering to the subject a composition of the invention under conditions suitable for the reduction or prevention of tissue rejection in the subject.

In another embodiment, the invention features a method for validating a VEGF and/or VEGFr gene target, comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein one of the siNA strands includes a sequence complementary to RNA of a VEGF and/or VEGFr target gene; (b) introducing the siNA molecule into a cell, tissue, or organism under conditions suitable for modulating expression of the VEGF and/or VEGFr target gene in the cell, tissue, or organism; and (c) determining the function of the gene by assaying for any phenotypic change in the cell, tissue, or organism.

In another embodiment, the invention features a method for validating a VEGF and/or VEGFr target comprising: (a) synthesizing a siNA molecule of the invention, which can be chemically-modified, wherein one of the siNA strands includes a sequence complementary to RNA of a VEGF and/or VEGFr target gene; (b) introducing the siNA molecule into a biological system under conditions suitable for modulating expression of the VEGF and/or VEGFr target gene in the biological system; and (c) determining the function of the gene by assaying for any phenotypic change in the biological system.

By "biological system" is meant, material, in a purified or unpurified form, from biological sources, including but not limited to human, animal, plant, insect, bacterial, viral or other sources, wherein the system comprises the components required for RNAi activity. The term "biological system" includes, for example, a cell, tissue, or organism, or extract thereof. The term biological system also includes reconstituted RNAi systems that can be used in an *in vitro* setting.

By "phenotypic change" is meant any detectable change to a cell that occurs in response to contact or treatment with a nucleic acid molecule of the invention (e.g., siNA). Such detectable changes include, but are not limited to, changes in shape, size, proliferation, motility, protein expression or RNA expression or other physical or chemical changes as can be assayed by methods known in the art. The detectable change can also include expression of reporter genes/molecules such as Green Florescent Protein (GFP) or various tags that are used to identify an expressed protein or any other cellular component that can be assayed.

In one embodiment, the invention features a kit containing a siNA molecule of the invention, which can be chemically-modified, that can be used to modulate the expression of a VEGF and/or VEGFr target gene in a cell, tissue, or organism. In another embodiment, the invention features a kit containing more than one siNA molecule of the invention, which  
5 can be chemically-modified, that can be used to modulate the expression of more than one VEGF and/or VEGFr target gene in a cell, tissue, or organism.

In one embodiment, the invention features a cell containing one or more siNA molecules of the invention, which can be chemically-modified. In another embodiment, the cell containing a siNA molecule of the invention is a mammalian cell. In yet another  
10 embodiment, the cell containing a siNA molecule of the invention is a human cell.

In one embodiment, the synthesis of a siNA molecule of the invention, which can be chemically-modified, comprises: (a) synthesis of two complementary strands of the siNA molecule; (b) annealing the two complementary strands together under conditions suitable to obtain a double-stranded siNA molecule. In another embodiment, synthesis of the two  
15 complementary strands of the siNA molecule is by solid phase oligonucleotide synthesis. In yet another embodiment, synthesis of the two complementary strands of the siNA molecule is by solid phase tandem oligonucleotide synthesis.

In one embodiment, the invention features a method for synthesizing a siNA duplex molecule comprising: (a) synthesizing a first oligonucleotide sequence strand of the siNA  
20 molecule, wherein the first oligonucleotide sequence strand comprises a cleavable linker molecule that can be used as a scaffold for the synthesis of the second oligonucleotide sequence strand of the siNA; (b) synthesizing the second oligonucleotide sequence strand of siNA on the scaffold of the first oligonucleotide sequence strand, wherein the second oligonucleotide sequence strand further comprises a chemical moiety than can be used to  
25 purify the siNA duplex; (c) cleaving the linker molecule of (a) under conditions suitable for the two siNA oligonucleotide strands to hybridize and form a stable duplex; and (d) purifying the siNA duplex utilizing the chemical moiety of the second oligonucleotide sequence strand. In one embodiment, cleavage of the linker molecule in (c) above takes place during deprotection of the oligonucleotide, for example under hydrolysis conditions



using an alkylamine base such as methylamine. In one embodiment, the method of synthesis comprises solid phase synthesis on a solid support such as controlled pore glass (CPG) or polystyrene, wherein the first sequence of (a) is synthesized on a cleavable linker, such as a succinyl linker, using the solid support as a scaffold. The cleavable linker in (a) used as a scaffold for synthesizing the second strand can comprise similar reactivity as the solid support derivatized linker, such that cleavage of the solid support derivatized linker and the cleavable linker of (a) takes place concomitantly. In another embodiment, the chemical moiety of (b) that can be used to isolate the attached oligonucleotide sequence comprises a trityl group, for example a dimethoxytrityl group, which can be employed in a trityl-on synthesis strategy as described herein. In yet another embodiment, the chemical moiety, such as a dimethoxytrityl group, is removed during purification, for example, using acidic conditions.

In a further embodiment, the method for siNA synthesis is a solution phase synthesis or hybrid phase synthesis wherein both strands of the siNA duplex are synthesized in tandem using a cleavable linker attached to the first sequence which acts a scaffold for synthesis of the second sequence. Cleavage of the linker under conditions suitable for hybridization of the separate siNA sequence strands results in formation of the double-stranded siNA molecule.

In another embodiment, the invention features a method for synthesizing a siNA duplex molecule comprising: (a) synthesizing one oligonucleotide sequence strand of the siNA molecule, wherein the sequence comprises a cleavable linker molecule that can be used as a scaffold for the synthesis of another oligonucleotide sequence; (b) synthesizing a second oligonucleotide sequence having complementarity to the first sequence strand on the scaffold of (a), wherein the second sequence comprises the other strand of the double-stranded siNA molecule and wherein the second sequence further comprises a chemical moiety than can be used to isolate the attached oligonucleotide sequence; (c) purifying the product of (b) utilizing the chemical moiety of the second oligonucleotide sequence strand under conditions suitable for isolating the full-length sequence comprising both siNA oligonucleotide strands connected by the cleavable linker and under conditions suitable for the two siNA oligonucleotide strands to hybridize and form a stable duplex. In one

embodiment, cleavage of the linker molecule in (c) above takes place during deprotection of the oligonucleotide, for example under hydrolysis conditions. In another embodiment, cleavage of the linker molecule in (c) above takes place after deprotection of the oligonucleotide. In another embodiment, the method of synthesis comprises solid phase  
5 synthesis on a solid support such as controlled pore glass (CPG) or polystyrene, wherein the first sequence of (a) is synthesized on a cleavable linker, such as a succinyl linker, using the solid support as a scaffold. The cleavable linker in (a) used as a scaffold for synthesizing the second strand can comprise similar reactivity or differing reactivity as the solid support derivatized linker, such that cleavage of the solid support derivatized linker and the  
10 cleavable linker of (a) takes place either concomitantly or sequentially. In one embodiment, the chemical moiety of (b) that can be used to isolate the attached oligonucleotide sequence comprises a trityl group, for example a dimethoxytrityl group.

In another embodiment, the invention features a method for making a double-stranded siNA molecule in a single synthetic process comprising: (a) synthesizing an oligonucleotide  
15 having a first and a second sequence, wherein the first sequence is complementary to the second sequence, and the first oligonucleotide sequence is linked to the second sequence via a cleavable linker, and wherein a terminal 5'-protecting group, for example, a 5'-O-dimethoxytrityl group (5'-O-DMT) remains on the oligonucleotide having the second sequence; (b) deprotecting the oligonucleotide whereby the deprotection results in the  
20 cleavage of the linker joining the two oligonucleotide sequences; and (c) purifying the product of (b) under conditions suitable for isolating the double-stranded siNA molecule, for example using a trityl-on synthesis strategy as described herein.

In another embodiment, the method of synthesis of siNA molecules of the invention comprises the teachings of Scaringe *et al.*, US Patent Nos. 5,889,136; 6,008,400; and  
25 6,111,086, incorporated by reference herein in their entirety.

In one embodiment, the invention features siNA constructs that mediate RNAi against a VEGF and/or VEGFr, wherein the siNA construct comprises one or more chemical modifications, for example, one or more chemical modifications having any of Formulae I-VII or any combination thereof that increases the nuclease resistance of the siNA construct.

In another embodiment, the invention features a method for generating siNA molecules with increased nuclease resistance comprising (a) introducing nucleotides having any of Formula I-VII or any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having increased nuclease resistance.

In one embodiment, the invention features siNA constructs that mediate RNAi against a VEGF and/or VEGFr, wherein the siNA construct comprises one or more chemical modifications described herein that modulates the binding affinity between the sense and antisense strands of the siNA construct.

In another embodiment, the invention features a method for generating siNA molecules with increased binding affinity between the sense and antisense strands of the siNA molecule comprising (a) introducing nucleotides having any of Formula I-VII or any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having increased binding affinity between the sense and antisense strands of the siNA molecule.

In one embodiment, the invention features siNA constructs that mediate RNAi against a VEGF and/or VEGFr, wherein the siNA construct comprises one or more chemical modifications described herein that modulates the binding affinity between the antisense strand of the siNA construct and a complementary target RNA sequence within a cell.

In one embodiment, the invention features siNA constructs that mediate RNAi against a VEGF and/or VEGFr, wherein the siNA construct comprises one or more chemical modifications described herein that modulates the binding affinity between the antisense strand of the siNA construct and a complementary target DNA sequence within a cell.

In another embodiment, the invention features a method for generating siNA molecules with increased binding affinity between the antisense strand of the siNA molecule and a complementary target RNA sequence comprising (a) introducing nucleotides having any of Formula I-VII or any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having

increased binding affinity between the antisense strand of the siNA molecule and a complementary target RNA sequence.

In another embodiment, the invention features a method for generating siNA molecules with increased binding affinity between the antisense strand of the siNA molecule and a complementary target DNA sequence comprising (a) introducing nucleotides having any of Formula I-VII or any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having increased binding affinity between the antisense strand of the siNA molecule and a complementary target DNA sequence.

10 In one embodiment, the invention features siNA constructs that mediate RNAi against a VEGF and/or VEGFr, wherein the siNA construct comprises one or more chemical modifications described herein that modulate the polymerase activity of a cellular polymerase capable of generating additional endogenous siNA molecules having sequence homology to the chemically-modified siNA construct.

15 In another embodiment, the invention features a method for generating siNA molecules capable of mediating increased polymerase activity of a cellular polymerase capable of generating additional endogenous siNA molecules having sequence homology to a chemically-modified siNA molecule comprising (a) introducing nucleotides having any of Formula I-VII or any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules capable of mediating increased polymerase activity of a cellular polymerase capable of generating additional endogenous siNA molecules having sequence homology to the chemically-modified siNA molecule.

25 In one embodiment, the invention features chemically-modified siNA constructs that mediate RNAi against a VEGF and/or VEGFr in a cell, wherein the chemical modifications do not significantly effect the interaction of siNA with a target RNA molecule, DNA molecule and/or proteins or other factors that are essential for RNAi in a manner that would decrease the efficacy of RNAi mediated by such siNA constructs.

In another embodiment, the invention features a method for generating siNA molecules with improved RNAi activity against VEGF and/or VEGFr comprising (a) introducing nucleotides having any of Formula I-VII or any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having improved RNAi activity.

In yet another embodiment, the invention features a method for generating siNA molecules with improved RNAi activity against a VEGF and/or VEGFr target RNA comprising (a) introducing nucleotides having any of Formula I-VII or any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having improved RNAi activity against the target RNA.

In yet another embodiment, the invention features a method for generating siNA molecules with improved RNAi activity against a VEGF and/or VEGFr target DNA comprising (a) introducing nucleotides having any of Formula I-VII or any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having improved RNAi activity against the target DNA.

In one embodiment, the invention features siNA constructs that mediate RNAi against a VEGF and/or VEGFr, wherein the siNA construct comprises one or more chemical modifications described herein that modulates the cellular uptake of the siNA construct.

In another embodiment, the invention features a method for generating siNA molecules against VEGF and/or VEGFr with improved cellular uptake comprising (a) introducing nucleotides having any of Formula I-VII or any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having improved cellular uptake.

In one embodiment, the invention features siNA constructs that mediate RNAi against a VEGF and/or VEGFr, wherein the siNA construct comprises one or more chemical modifications described herein that increases the bioavailability of the siNA construct, for

example, by attaching polymeric conjugates such as polyethyleneglycol or equivalent conjugates that improve the pharmacokinetics of the siNA construct, or by attaching conjugates that target specific tissue types or cell types *in vivo*. Non-limiting examples of such conjugates are described in Vargeese *et al.*, U.S. Serial No. 10/201,394 incorporated by  
5 reference herein.

In one embodiment, the invention features a method for generating siNA molecules of the invention with improved bioavailability, comprising (a) introducing a conjugate into the structure of a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having improved bioavailability. Such  
10 conjugates can include ligands for cellular receptors, such as peptides derived from naturally occurring protein ligands; protein localization sequences, including cellular ZIP code sequences; antibodies; nucleic acid aptamers; vitamins and other co-factors, such as folate and N-acetylgalactosamine; polymers, such as polyethyleneglycol (PEG); phospholipids; polyamines, such as spermine or spermidine; and others.

15 In another embodiment, the invention features a method for generating siNA molecules of the invention with improved bioavailability comprising (a) introducing an excipient formulation to a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA molecules having improved bioavailability. Such excipients include polymers such as cyclodextrins, lipids, cationic lipids, polyamines,  
20 phospholipids, and others.

In another embodiment, the invention features a method for generating siNA molecules of the invention with improved bioavailability comprising (a) introducing nucleotides having any of Formulae I-VII or any combination thereof into a siNA molecule, and (b) assaying the siNA molecule of step (a) under conditions suitable for isolating siNA  
25 molecules having improved bioavailability.

In another embodiment, polyethylene glycol (PEG) can be covalently attached to siNA compounds of the present invention. The attached PEG can be any molecular weight, preferably from about 2,000 to about 50,000 daltons (Da).

The present invention can be used alone or as a component of a kit having at least one of the reagents necessary to carry out the *in vitro* or *in vivo* introduction of RNA to test samples and/or subjects. For example, preferred components of the kit include a siNA molecule of the invention and a vehicle that promotes introduction of the siNA into cells of interest as described herein (e.g., using lipids and other methods of transfection known in the art, see for example Beigelman *et al.*, US 6,395,713). The kit can be used for target validation, such as in determining gene function and/or activity, or in drug optimization, and in drug discovery (see for example Usman *et al.*, USSN 60/402,996). Such a kit can also include instructions to allow a user of the kit to practice the invention.

The term "short interfering nucleic acid", "siNA", "short interfering RNA", "siRNA", "short interfering nucleic acid molecule", "short interfering oligonucleotide molecule", or "chemically-modified short interfering nucleic acid molecule" as used herein refers to any nucleic acid molecule capable of inhibiting or down regulating gene expression, for example by mediating RNA interference "RNAi" or gene silencing in a sequence-specific manner; see for example Bass, 2001, *Nature*, 411, 428-429; Elbashir *et al.*, 2001, *Nature*, 411, 494-498; and Kreutzer *et al.*, International PCT Publication No. WO 00/44895; Zernicka-Goetz *et al.*, International PCT Publication No. WO 01/36646; Fire, International PCT Publication No. WO 99/32619; Plaetinck *et al.*, International PCT Publication No. WO 00/01846; Mello and Fire, International PCT Publication No. WO 01/29058; Deschamps-Depaillette, International PCT Publication No. WO 99/07409; and Li *et al.*, International PCT Publication No. WO 00/44914; Allshire, 2002, *Science*, 297, 1818-1819; Volpe *et al.*, 2002, *Science*, 297, 1833-1837; Jenuwein, 2002, *Science*, 297, 2215-2218; and Hall *et al.*, 2002, *Science*, 297, 2232-2237; Hutvagner and Zamore, 2002, *Science*, 297, 2056-60; McManus *et al.*, 2002, *RNA*, 8, 842-850; Reinhart *et al.*, 2002, *Gene & Dev.*, 16, 1616-1626; and Reinhart & Bartel, 2002, *Science*, 297, 1831). Non limiting examples of siNA molecules of the invention are shown in Figures 4-6, and Tables II, III, and IV herein. For example the siNA can be a double-stranded polynucleotide molecule comprising self-complementary sense and antisense regions, wherein the antisense region comprises nucleotide sequence that is complementary to nucleotide sequence in a target nucleic acid molecule or a portion thereof and the sense region having nucleotide sequence corresponding to the target nucleic

acid sequence or a portion thereof. The siNA can be assembled from two separate oligonucleotides, where one strand is the sense strand and the other is the antisense strand, wherein the antisense and sense strands are self-complementary (i.e. each strand comprises nucleotide sequence that is complementary to nucleotide sequence in the other strand; such as where the antisense strand and sense strand form a duplex or double stranded structure, for example wherein the double stranded region is about 19 base pairs); the antisense strand comprises nucleotide sequence that is complementary to nucleotide sequence in a target nucleic acid molecule or a portion thereof and the sense strand comprises nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof.

Alternatively, the siNA is assembled from a single oligonucleotide, where the self-complementary sense and antisense regions of the siNA are linked by means of a nucleic acid based or non-nucleic acid-based linker(s). The siNA can be a polynucleotide with a hairpin secondary structure, having self-complementary sense and antisense regions, wherein the antisense region comprises nucleotide sequence that is complementary to nucleotide sequence in a separate target nucleic acid molecule or a portion thereof and the sense region having nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof. The siNA can be a circular single-stranded polynucleotide having two or more loop structures and a stem comprising self-complementary sense and antisense regions, wherein the antisense region comprises nucleotide sequence that is complementary to nucleotide sequence in a target nucleic acid molecule or a portion thereof and the sense region having nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof, and wherein the circular polynucleotide can be processed either *in vivo* or *in vitro* to generate an active siNA molecule capable of mediating RNAi. The siNA can also comprise a single stranded polynucleotide having nucleotide sequence complementary to nucleotide sequence in a target nucleic acid molecule or a portion thereof (for example, where such siNA molecule does not require the presence within the siNA molecule of nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof), wherein the single stranded polynucleotide can further comprise a terminal phosphate group, such as a 5'-phosphate (see for example Martinez *et al.*, 2002, *Cell.*, 110, 563-574 and Schwarz *et al.*, 2002, *Molecular Cell*, 10, 537-568), or 5',3'-diphosphate. In certain embodiment, the siNA molecule of the invention comprises separate sense and antisense



sequences or regions, wherein the sense and antisense regions are covalently linked by nucleotide or non-nucleotide linkers molecules as is known in the art, or are alternately non-covalently linked by ionic interactions, hydrogen bonding, van der Waals interactions, hydrophobic interactions, and/or stacking interactions. In certain embodiments, the siNA molecules of the invention comprise nucleotide sequence that is complementary to nucleotide sequence of a target gene. In another embodiment, the siNA molecule of the invention interacts with nucleotide sequence of a target gene in a manner that causes inhibition of expression of the target gene. As used herein, siNA molecules need not be limited to those molecules containing only RNA, but further encompasses chemically-modified nucleotides and non-nucleotides. In certain embodiments, the short interfering nucleic acid molecules of the invention lack 2'-hydroxy (2'-OH) containing nucleotides. Applicant describes in certain embodiments short interfering nucleic acids that do not require the presence of nucleotides having a 2'-hydroxy group for mediating RNAi and as such, short interfering nucleic acid molecules of the invention optionally do not include any ribonucleotides (e.g., nucleotides having a 2'-OH group). Such siNA molecules that do not require the presence of ribonucleotides within the siNA molecule to support RNAi can however have an attached linker or linkers or other attached or associated groups, moieties, or chains containing one or more nucleotides with 2'-OH groups. Optionally, siNA molecules can comprise ribonucleotides at about 5, 10, 20, 30, 40, or 50% of the nucleotide positions. The modified short interfering nucleic acid molecules of the invention can also be referred to as short interfering modified oligonucleotides "siMON." As used herein, the term siNA is meant to be equivalent to other terms used to describe nucleic acid molecules that are capable of mediating sequence specific RNAi, for example short interfering RNA (siRNA), double-stranded RNA (dsRNA), micro-RNA (miRNA), short hairpin RNA (shRNA), short interfering oligonucleotide, short interfering nucleic acid, short interfering modified oligonucleotide, chemically-modified siRNA, post-transcriptional gene silencing RNA (ptgsRNA), and others. In addition, as used herein, the term RNAi is meant to be equivalent to other terms used to describe sequence specific RNA interference, such as post transcriptional gene silencing, or epigenetics. For example, siNA molecules of the invention can be used to epigenetically silence genes at both the post-transcriptional level or the pre-transcriptional level. In a non-limiting example, epigenetic regulation of gene expression by

siNA molecules of the invention can result from siNA mediated modification of chromatin structure to alter gene expression (see, for example, Allshire, 2002, *Science*, 297, 1818-1819; Volpe *et al.*, 2002, *Science*, 297, 1833-1837; Jenuwein, 2002, *Science*, 297, 2215-2218; and Hall *et al.*, 2002, *Science*, 297, 2232-2237).

5 By "modulate" is meant that the expression of the gene, or level of RNA molecule or equivalent RNA molecules encoding one or more proteins or protein subunits, or activity of one or more proteins or protein subunits is up regulated or down regulated, such that expression, level, or activity is greater than or less than that observed in the absence of the modulator. For example, the term "modulate" can mean "inhibit," but the use of the word  
10 "modulate" is not limited to this definition.

By "inhibit", "down-regulate", or "reduce", it is meant that the expression of the gene, or level of RNA molecules or equivalent RNA molecules encoding one or more proteins or protein subunits, or activity of one or more proteins or protein subunits, is reduced below that observed in the absence of the nucleic acid molecules (e.g., siNA) of the invention. In  
15 one embodiment, inhibition, down-regulation or reduction with an siNA molecule is below that level observed in the presence of an inactive or attenuated molecule. In another embodiment, inhibition, down-regulation, or reduction with siNA molecules is below that level observed in the presence of, for example, an siNA molecule with scrambled sequence or with mismatches. In another embodiment, inhibition, down-regulation, or reduction of  
20 gene expression with a nucleic acid molecule of the instant invention is greater in the presence of the nucleic acid molecule than in its absence.

By "gene" or "target gene" is meant, a nucleic acid that encodes an RNA, for example, nucleic acid sequences including, but not limited to, structural genes encoding a polypeptide. The target gene can be a gene derived from a cell, an endogenous gene, a transgene, or  
25 exogenous genes such as genes of a pathogen, for example a virus, which is present in the cell after infection thereof. The cell containing the target gene can be derived from or contained in any organism, for example a plant, animal, protozoan, virus, bacterium, or fungus. Non-limiting examples of plants include monocots, dicots, or gymnosperms. Non-

limiting examples of animals include vertebrates or invertebrates. Non-limiting examples of fungi include molds or yeasts.

By "VEGF" as used herein is meant, any vascular endothelial growth factor (e.g., VEGF, VEGF-A, VEGF-B, VEGF-C, VEGF-D) protein, peptide, or polypeptide having  
5 vascular endothelial growth factor activity, such as encoded by VEGF Genbank Accession Nos. shown in Table I. The term VEGF also refers to nucleic acid sequences encoding any vascular endothelial growth factor protein, peptide, or polypeptide having vascular endothelial growth factor activity.

By "VEGF-B" is meant, protein, peptide, or polypeptide receptor or a derivative  
10 thereof, such as encoded by Genbank Accession No. NM\_003377, having vascular endothelial growth factor type B activity. The term VEGF-B also refers to nucleic acid sequences encoding any VEGF-B protein, peptide, or polypeptide having VEGF-B activity.

By "VEGF-C" is meant, protein, peptide, or polypeptide receptor or a derivative thereof, such as encoded by Genbank Accession No. NM\_005429, having vascular  
15 endothelial growth factor type C activity. The term VEGF-C also refers to nucleic acid sequences encoding any VEGF-C protein, peptide, or polypeptide having VEGF-C activity.

By "VEGF-D" is meant, protein, peptide, or polypeptide receptor or a derivative thereof, such as encoded by Genbank Accession No. NM\_004469, having vascular  
20 endothelial growth factor type D activity. The term VEGF-D also refers to nucleic acid sequences encoding any VEGF-D protein, peptide, or polypeptide having VEGF-D activity.

By "VEGFr" as used herein is meant, any vascular endothelial growth factor receptor protein, peptide, or polypeptide (e.g., VEGFr1, VEGFr2, or VEGFr3, including both membrane bound and/or soluble forms thereof) having vascular endothelial growth factor receptor activity, such as encoded by VEGFr Genbank Accession Nos. shown in Table I.  
25 The term VEGFr also refers to nucleic acid sequences encoding any vascular endothelial growth factor receptor protein, peptide, or polypeptide having vascular endothelial growth factor receptor activity.

By "VEGFr1" is meant, protein, peptide, or polypeptide receptor or a derivative thereof, such as encoded by Genbank Accession No. NM\_002019, having vascular endothelial growth factor receptor type 1 (*flt*) activity, for example, having the ability to bind a vascular endothelial growth factor. The term VEGF1 also refers to nucleic acid sequences encoding any VEGFr1 protein, peptide, or polypeptide having VEGFr1 activity.

By "VEGFr2" is meant, protein, peptide, or polypeptide receptor or a derivative thereof, such as encoded by Genbank Accession No. NM\_002253, having vascular endothelial growth factor receptor type 2 (*kdr*) activity, for example, having the ability to bind a vascular endothelial growth factor. The term VEGF2 also refers to nucleic acid sequences encoding any VEGFr2 protein, peptide, or polypeptide having VEGFr2 activity.

By "VEGFr3" is meant, protein, peptide, or polypeptide receptor or a derivative thereof, such as encoded by Genbank Accession No. NM\_002020 having vascular endothelial growth factor receptor type 3 (*kdr*) activity, for example, having the ability to bind a vascular endothelial growth factor. The term VEGF3 also refers to nucleic acid sequences encoding any VEGFr3 protein, peptide, or polypeptide having VEGFr3 activity.

By "highly conserved sequence region" is meant, a nucleotide sequence of one or more regions in a target gene does not vary significantly from one generation to the other or from one biological system to the other.

By "sense region" is meant a nucleotide sequence of a siNA molecule having complementarity to an antisense region of the siNA molecule. In addition, the sense region of a siNA molecule can comprise a nucleic acid sequence having homology with a target nucleic acid sequence.

By "antisense region" is meant a nucleotide sequence of a siNA molecule having complementarity to a target nucleic acid sequence. In addition, the antisense region of a siNA molecule can optionally comprise a nucleic acid sequence having complementarity to a sense region of the siNA molecule.

By "target nucleic acid" is meant any nucleic acid sequence whose expression or activity is to be modulated. The target nucleic acid can be DNA or RNA.

By "complementarity" is meant that a nucleic acid can form hydrogen bond(s) with another nucleic acid sequence by either traditional Watson-Crick or other non-traditional types. In reference to the nucleic molecules of the present invention, the binding free energy for a nucleic acid molecule with its complementary sequence is sufficient to allow the relevant function of the nucleic acid to proceed, e.g., RNAi activity. Determination of binding free energies for nucleic acid molecules is well known in the art (see, e.g., Turner *et al.*, 1987, *CSH Symp. Quant. Biol.* LII pp.123-133; Frier *et al.*, 1986, *Proc. Nat. Acad. Sci.* USA 83:9373-9377; Turner *et al.*, 1987, *J. Am. Chem. Soc.* 109:3783-3785). A percent complementarity indicates the percentage of contiguous residues in a nucleic acid molecule that can form hydrogen bonds (e.g., Watson-Crick base pairing) with a second nucleic acid sequence (e.g., 5, 6, 7, 8, 9, 10 out of 10 being 50%, 60%, 70%, 80%, 90%, and 100% complementary). "Perfectly complementary" means that all the contiguous residues of a nucleic acid sequence will hydrogen bond with the same number of contiguous residues in a second nucleic acid sequence.

The siRNA molecules of the invention represent a novel therapeutic approach to treat a variety of pathologic indications or other conditions, such as tumor angiogenesis and cancer, including but not limited to breast cancer, lung cancer (including non-small cell lung carcinoma), prostate cancer, colorectal cancer, brain cancer, esophageal cancer, bladder cancer, pancreatic cancer, cervical cancer, head and neck cancer, skin cancers, nasopharyngeal carcinoma, liposarcoma, epithelial carcinoma, renal cell carcinoma, gallbladder adeno carcinoma, parotid adenocarcinoma, ovarian cancer, melanoma, lymphoma, glioma, endometrial sarcoma, multidrug resistant cancers, diabetic retinopathy, macular degeneration, neovascular glaucoma, myopic degeneration, arthritis, psoriasis, endometriosis, female reproduction, verruca vulgaris, angiofibroma of tuberous sclerosis, pot-wine stains, Sturge Weber syndrome, Kippel-Trenaunay-Weber syndrome, Osler-Weber-Rendu syndrome, renal disease such as Autosomal dominant polycystic kidney disease (ADPKD), and any other diseases or conditions that are related to or will respond to the levels of VEGF, VEGFr1, VEGFr2 and/or VEGFr3 in a cell or tissue, alone or in

combination with other therapies. The reduction of VEGF, VEGFr1, VEGFr2 and/or VEGFr3 expression (specifically VEGF, VEGFr1, VEGFr2 and/or VEGFr3 gene RNA levels) and thus reduction in the level of the respective protein relieves, to some extent, the symptoms of the disease or condition.ue

5 In one embodiment of the present invention, each sequence of a siNA molecule of the invention is independently about 18 to about 24 nucleotides in length, in specific embodiments about 18, 19, 20, 21, 22, 23, or 24 nucleotides in length. In another embodiment, the siNA duplexes of the invention independently comprise about 17 to about 23 base pairs (*e.g.*, about 17, 18, 19, 20, 21, 22 or 23). In yet another embodiment, siNA  
10 molecules of the invention comprising hairpin or circular structures are about 35 to about 55 (*e.g.*, about 35, 40, 45, 50 or 55) nucleotides in length, or about 38 to about 44 (*e.g.*, 38, 39, 40, 41, 42, 43 or 44) nucleotides in length and comprising about 16 to about 22 (*e.g.*, about 16, 17, 18, 19, 20, 21 or 22) base pairs. Exemplary siNA molecules of the invention are shown in Table II. Exemplary synthetic siNA molecules of the invention are shown in  
15 Tables III and IV and/or Figures 4-5.

As used herein "cell" is used in its usual biological sense, and does not refer to an entire multicellular organism, *e.g.*, specifically does not refer to a human. The cell can be present in an organism, *e.g.*, birds, plants and mammals such as humans, cows, sheep, apes, monkeys, swine, dogs, and cats. The cell can be prokaryotic (*e.g.*, bacterial cell) or  
20 eukaryotic (*e.g.*, mammalian or plant cell). The cell can be of somatic or germ line origin, totipotent or pluripotent, dividing or non-dividing. The cell can also be derived from or can comprise a gamete or embryo, a stem cell, or a fully differentiated cell.

The siNA molecules of the invention are added directly, or can be complexed with cationic lipids, packaged within liposomes, or otherwise delivered to target cells or tissues.  
25 The nucleic acid or nucleic acid complexes can be locally administered to relevant tissues *ex vivo*, or *in vivo* through injection, infusion pump or stent, with or without their incorporation in biopolymers. In particular embodiments, the nucleic acid molecules of the invention comprise sequences shown in Tables II-III and/or Figures 4-5. Examples of such nucleic acid molecules consist essentially of sequences defined in these tables and figures.

Furthermore, the chemically modified constructs described in Table IV can be applied to any siNA sequence of the invention.

In another aspect, the invention provides mammalian cells containing one or more siNA molecules of this invention. The one or more siNA molecules can independently be  
5 targeted to the same or different sites.

By "RNA" is meant a molecule comprising at least one ribonucleotide residue. By "ribonucleotide" is meant a nucleotide with a hydroxyl group at the 2' position of a  $\beta$ -D-ribo-furanose moiety. The terms include double-stranded RNA, single-stranded RNA, isolated RNA such as partially purified RNA, essentially pure RNA, synthetic RNA,  
10 recombinantly produced RNA, as well as altered RNA that differs from naturally occurring RNA by the addition, deletion, substitution and/or alteration of one or more nucleotides. Such alterations can include addition of non-nucleotide material, such as to the end(s) of the siNA or internally, for example at one or more nucleotides of the RNA. Nucleotides in the RNA molecules of the instant invention can also comprise non-standard nucleotides, such as  
15 non-naturally occurring nucleotides or chemically synthesized nucleotides or deoxynucleotides. These altered RNAs can be referred to as analogs or analogs of naturally-occurring RNA.

By "subject" is meant an organism, which is a donor or recipient of explanted cells or the cells themselves. "Subject" also refers to an organism to which the nucleic acid  
20 molecules of the invention can be administered. In one embodiment, a subject is a mammal or mammalian cells. In another embodiment, a subject is a human or human cells.

The term "phosphorothioate" as used herein refers to an internucleotide linkage having Formula I, wherein Z and/or W comprise a sulfur atom. Hence, the term phosphorothioate refers to both phosphorothioate and phosphorodithioate internucleotide linkages.

25 The term "universal base" as used herein refers to nucleotide base analogs that form base pairs with each of the natural DNA/RNA bases with little discrimination between them. Non-limiting examples of universal bases include C-phenyl, C-naphthyl and other aromatic derivatives, inosine, azole carboxamides, and nitroazole derivatives such as 3-nitropyrrole,

4-nitroindole, 5-nitroindole, and 6-nitroindole as known in the art (see for example Loakes, 2001, *Nucleic Acids Research*, 29, 2437-2447).

The term "acyclic nucleotide" as used herein refers to any nucleotide having an acyclic ribose sugar, for example where any of the ribose carbons (C1, C2, C3, C4, or C5), are  
5 independently or in combination absent from the nucleotide.

The nucleic acid molecules of the instant invention, individually, or in combination or in conjunction with other drugs, can be used to treat diseases or conditions discussed herein (e.g., cancers and othe proliferative conditions). For example, to treat a particular disease or condition, the siNA molecules can be administered to a subject or can be administered to  
10 other appropriate cells evident to those skilled in the art, individually or in combination with one or more drugs under conditions suitable for the treatment.

In a further embodiment, the siNA molecules can be used in combination with other known treatments to treat conditions or diseases discussed above. For example, the described molecules could be used in combination with one or more known therapeutic  
15 agents to treat a disease or condition. Non-limiting examples of other therapeutic agents that can be readily combined with a siNA molecule of the invention are enzymatic nucleic acid molecules, allosteric nucleic acid molecules, antisense, decoy, or aptamer nucleic acid molecules, antibodies such as monoclonal antibodies, small molecules, and other organic and/or inorganic compounds including metals, salts and ions.

20 In one embodiment, the invention features an expression vector comprising a nucleic acid sequence encoding at least one siNA molecule of the invention, in a manner which allows expression of the siNA molecule. For example, the vector can contain sequence(s) encoding both strands of a siNA molecule comprising a duplex. The vector can also contain sequence(s) encoding a single nucleic acid molecule that is self-complementary and thus  
25 forms a siNA molecule. Non-limiting examples of such expression vectors are described in Paul *et al.*, 2002, *Nature Biotechnology*, 19, 505; Miyagishi and Taira, 2002, *Nature Biotechnology*, 19, 497; Lee *et al.*, 2002, *Nature Biotechnology*, 19, 500; and Novina *et al.*, 2002, *Nature Medicine*, advance online publication doi:10.1038/nm725.



In another embodiment, the invention features a mammalian cell, for example, a human cell, including an expression vector of the invention.

In yet another embodiment, the expression vector of the invention comprises a sequence for a siNA molecule having complementarity to a RNA molecule referred to by a  
5 Genbank Accession numbers, for example Genbank Accession Nos. shown in Table I.

In one embodiment, an expression vector of the invention comprises a nucleic acid sequence encoding two or more siNA molecules, which can be the same or different.

In another aspect of the invention, siNA molecules that interact with target RNA molecules and down-regulate gene encoding target RNA molecules (for example target  
10 RNA molecules referred to by Genbank Accession numbers herein) are expressed from transcription units inserted into DNA or RNA vectors. The recombinant vectors can be DNA plasmids or viral vectors. siNA expressing viral vectors can be constructed based on, but not limited to, adeno-associated virus, retrovirus, adenovirus, or alphavirus. The recombinant vectors capable of expressing the siNA molecules can be delivered as described herein, and  
15 persist in target cells. Alternatively, viral vectors can be used that provide for transient expression of siNA molecules. Such vectors can be repeatedly administered as necessary. Once expressed, the siNA molecules bind and down-regulate gene function or expression via RNA interference (RNAi). Delivery of siNA expressing vectors can be systemic, such as by intravenous or intramuscular administration, by administration to target cells ex-planted  
20 from a subject followed by reintroduction into the subject, or by any other means that would allow for introduction into the desired target cell.

By "vectors" is meant any nucleic acid- and/or viral-based technique used to deliver a desired nucleic acid.

Other features and advantages of the invention will be apparent from the following  
25 description of the preferred embodiments thereof, and from the claims.

### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows a non-limiting example of a scheme for the synthesis of siNA molecules. The complementary siNA sequence strands, strand 1 and strand 2, are synthesized in tandem and are connected by a cleavable linkage, such as a nucleotide succinate or abasic succinate, which can be the same or different from the cleavable linker used for solid phase synthesis on a solid support. The synthesis can be either solid phase or solution phase, in the example shown, the synthesis is a solid phase synthesis. The synthesis is performed such that a protecting group, such as a dimethoxytrityl group, remains intact on the terminal nucleotide of the tandem oligonucleotide. Upon cleavage and deprotection of the oligonucleotide, the two siNA strands spontaneously hybridize to form a siNA duplex, which allows the purification of the duplex by utilizing the properties of the terminal protecting group, for example by applying a trityl on purification method wherein only duplexes/oligonucleotides with the terminal protecting group are isolated.

Figure 2 shows a MALDI-TOV mass spectrum of a purified siNA duplex synthesized by a method of the invention. The two peaks shown correspond to the predicted mass of the separate siNA sequence strands. This result demonstrates that the siNA duplex generated from tandem synthesis can be purified as a single entity using a simple trityl-on purification methodology.

Figure 3 shows a non-limiting proposed mechanistic representation of target RNA degradation involved in RNAi. Double-stranded RNA (dsRNA), which is generated by RNA-dependent RNA polymerase (RdRP) from foreign single-stranded RNA, for example viral, transposon, or other exogenous RNA, activates the DICER enzyme that in turn generates siNA duplexes. Alternately, synthetic or expressed siNA can be introduced directly into a cell by appropriate means. An active siNA complex forms which recognizes a target RNA, resulting in degradation of the target RNA by the RISC endonuclease complex or in the synthesis of additional RNA by RNA-dependent RNA polymerase (RdRP), which can activate DICER and result in additional siNA molecules, thereby amplifying the RNAi response.

Figure 4A-F shows non-limiting examples of chemically-modified siNA constructs of the present invention. In the figure, N stands for any nucleotide (adenosine, guanosine, cytosine, uridine, or optionally thymidine, for example thymidine can be substituted in the overhanging regions designated by parenthesis (N N)). Various modifications are shown for the sense and antisense strands of the siNA constructs.

Figure 4A: The sense strand comprises 21 nucleotides having four phosphorothioate 5'- and 3'-terminal internucleotide linkages, wherein the two terminal 3'-nucleotides are optionally base paired and wherein all pyrimidine nucleotides that may be present are 2'-O-methyl or 2'-deoxy-2'-fluoro modified nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein. The antisense strand comprises 21 nucleotides, optionally having a 3'-terminal glyceryl moiety and wherein the two terminal 3'-nucleotides are optionally complementary to the target RNA sequence, and having one 3'-terminal phosphorothioate internucleotide linkage and four 5'-terminal phosphorothioate internucleotide linkages and wherein all pyrimidine nucleotides that may be present are 2'-deoxy-2'-fluoro modified nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein.

Figure 4B: The sense strand comprises 21 nucleotides wherein the two terminal 3'-nucleotides are optionally base paired and wherein all pyrimidine nucleotides that may be present are 2'-O-methyl or 2'-deoxy-2'-fluoro modified nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein. The antisense strand comprises 21 nucleotides, optionally having a 3'-terminal glyceryl moiety and wherein the two terminal 3'-nucleotides are optionally complementary to the target RNA sequence, and wherein all pyrimidine nucleotides that may be present are 2'-deoxy-2'-fluoro modified nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein.

**Figure 4C:** The sense strand comprises 21 nucleotides having 5'- and 3'- terminal cap moieties wherein the two terminal 3'-nucleotides are optionally base paired and wherein all pyrimidine nucleotides that may be present are 2'-O-methyl or 2'-deoxy-2'-fluoro modified nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein. The antisense strand comprises 21 nucleotides, optionally having a 3'-terminal glyceryl moiety and wherein the two terminal 3'-nucleotides are optionally complementary to the target RNA sequence, and having one 3'-terminal phosphorothioate internucleotide linkage and wherein all pyrimidine nucleotides that may be present are 2'-deoxy-2'-fluoro modified nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein.

**Figure 4D:** The sense strand comprises 21 nucleotides having 5'- and 3'- terminal cap moieties wherein the two terminal 3'-nucleotides are optionally base paired and wherein all pyrimidine nucleotides that may be present are 2'-deoxy-2'-fluoro modified nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein and wherein all purine nucleotides that may be present are 2'-deoxy nucleotides. The antisense strand comprises 21 nucleotides, optionally having a 3'-terminal glyceryl moiety and wherein the two terminal 3'-nucleotides are optionally complementary to the target RNA sequence, and having one 3'-terminal phosphorothioate internucleotide linkage and wherein all pyrimidine nucleotides that may be present are 2'-deoxy-2'-fluoro modified nucleotides and all purine nucleotides that may be present are 2'-O-methyl modified nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein.

**Figure 4E:** The sense strand comprises 21 nucleotides having 5'- and 3'- terminal cap moieties wherein the two terminal 3'-nucleotides are optionally base paired and wherein all pyrimidine nucleotides that may be present are 2'-deoxy-2'-fluoro modified nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein. The antisense strand comprises 21 nucleotides, optionally having a 3'-terminal glyceryl moiety and wherein the

two terminal 3'-nucleotides are optionally complementary to the target RNA sequence, and wherein all pyrimidine nucleotides that may be present are 2'-deoxy-2'-fluoro modified nucleotides and all purine nucleotides that may be present are 2'-O-methyl modified nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein.

**Figure 4F:** The sense strand comprises 21 nucleotides having 5'- and 3'- terminal cap moieties wherein the two terminal 3'-nucleotides are optionally base paired and wherein all pyrimidine nucleotides that may be present are 2'-deoxy-2'-fluoro modified nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein. The antisense strand comprises 21 nucleotides, optionally having a 3'-terminal glyceryl moiety and wherein the two terminal 3'-nucleotides are optionally complementary to the target RNA sequence, and having one 3'-terminal phosphorothioate internucleotide linkage and wherein all pyrimidine nucleotides that may be present are 2'-deoxy-2'-fluoro modified nucleotides and all purine nucleotides that may be present are 2'-deoxy nucleotides except for (N N) nucleotides, which can comprise ribonucleotides, deoxynucleotides, universal bases, or other chemical modifications described herein. The antisense strand of constructs A-F comprise sequence complementary to any target nucleic acid sequence of the invention.

**Figure 5A-F** shows non-limiting examples of specific chemically-modified siNA sequences of the invention. A-F applies the chemical modifications described in **Figure 4A-F** to a VEGFr1 siNA sequence. Such chemical modifications can be applied to any sequence herein, such as any VEGF, VEGFr1, VEGFr2, or VEGFr3 sequence.

**Figure 6** shows non-limiting examples of different siNA constructs of the invention. The examples shown (constructs 1, 2, and 3) have 19 representative base pairs; however, different embodiments of the invention include any number of base pairs described herein. Bracketed regions represent nucleotide overhangs, for example comprising about 1, 2, 3, or 4 nucleotides in length, preferably about 2 nucleotides. Constructs 1 and 2 can be used independently for RNAi activity. Construct 2 can comprise a polynucleotide or non-nucleotide linker, which can optionally be designed as a biodegradable linker. In one

embodiment, the loop structure shown in construct 2 can comprise a biodegradable linker that results in the formation of construct 1 *in vivo* and/or *in vitro*. In another example, construct 3 can be used to generate construct 2 under the same principle wherein a linker is used to generate the active siNA construct 2 *in vivo* and/or *in vitro*, which can optionally  
5 utilize another biodegradable linker to generate the active siNA construct 1 *in vivo* and/or *in vitro*. As such, the stability and/or activity of the siNA constructs can be modulated based on the design of the siNA construct for use *in vivo* or *in vitro* and/or *in vitro*.

Figure 7A-C is a diagrammatic representation of a scheme utilized in generating an expression cassette to generate siNA hairpin constructs.

10 **Figure 7A:** A DNA oligomer is synthesized with a 5'-restriction site (R1) sequence followed by a region having sequence identical (sense region of siNA) to a predetermined VEGF and/or VEGFr target sequence, wherein the sense region comprises, for example, about 19, 20, 21, or 22 nucleotides (N) in length, which is followed by a loop sequence of defined sequence (X), comprising, for example, about 3 to about 10 nucleotides.

15 **Figure 7B:** The synthetic construct is then extended by DNA polymerase to generate a hairpin structure having self-complementary sequence that will result in a siNA transcript having specificity for a VEGF and/or VEGFr target sequence and having self-complementary sense and antisense regions.

20 **Figure 7C:** The construct is heated (for example to about 95°C) to linearize the sequence, thus allowing extension of a complementary second DNA strand using a primer to the 3'-restriction sequence of the first strand. The double-stranded DNA is then inserted into an appropriate vector for expression in cells. The construct can be designed such that a 3'-terminal nucleotide overhang results from the transcription, for example by engineering restriction sites and/or utilizing a poly-U termination region as described in Paul *et al.*, 2002,  
25 *Nature Biotechnology*, 29, 505-508.

**Figure 8A-C** is a diagrammatic representation of a scheme utilized in generating an expression cassette to generate double-stranded siNA constructs.

**Figure 8A:** A DNA oligomer is synthesized with a 5'-restriction (R1) site sequence followed by a region having sequence identical (sense region of siNA) to a predetermined VEGF and/or VEGFr target sequence, wherein the sense region comprises, for example, about 19, 20, 21, or 22 nucleotides (N) in length, and which is followed by a 3'-restriction site (R2) which is adjacent to a loop sequence of defined sequence (X).

**Figure 8B:** The synthetic construct is then extended by DNA polymerase to generate a hairpin structure having self-complementary sequence.

**Figure 8C:** The construct is processed by restriction enzymes specific to R1 and R2 to generate a double-stranded DNA which is then inserted into an appropriate vector for expression in cells. The transcription cassette is designed such that a U6 promoter region flanks each side of the dsDNA which generates the separate sense and antisense strands of the siNA. Poly T termination sequences can be added to the constructs to generate U overhangs in the resulting transcript.

**Figure 9A-E** is a diagrammatic representation of a method used to determine target sites for siNA mediated RNAi within a particular target nucleic acid sequence, such as messenger RNA.

**Figure 9A:** A pool of siNA oligonucleotides are synthesized wherein the antisense region of the siNA constructs has complementarity to target sites across the target nucleic acid sequence, and wherein the sense region comprises sequence complementary to the antisense region of the siNA.

**Figure 9B&C:** (Figure 9B) The sequences are pooled and are inserted into vectors such that (Figure 9C) transfection of a vector into cells results in the expression of the siNA.

**Figure 9D:** Cells are sorted based on phenotypic change that is associated with modulation of the target nucleic acid sequence.

**Figure 9E:** The siNA is isolated from the sorted cells and is sequenced to identify efficacious target sites within the target nucleic acid sequence.

Figure 10 shows non-limiting examples of different stabilization chemistries (1-10) that can be used, for example, to stabilize the 3'-end of siNA sequences of the invention, including (1) [3-3']-inverted deoxyribose; (2) deoxyribonucleotide; (3) [5'-3']-3'-deoxyribonucleotide; (4) [5'-3']-ribonucleotide; (5) [5'-3']-3'-O-methyl ribonucleotide; (6) 3'-glyceryl; (7) [3'-5']-3'-deoxyribonucleotide; (8) [3'-3']-deoxyribonucleotide; (9) [5'-2']-deoxyribonucleotide; and (10) [5'-3']-dideoxyribonucleotide. In addition to modified and unmodified backbone chemistries indicated in the figure, these chemistries can be combined with different backbone modifications as described herein, for example, backbone modifications having Formula I. In addition, the 2'-deoxy nucleotide shown 5' to the terminal modifications shown can be another modified or unmodified nucleotide or non-nucleotide described herein, for example modifications having any of Formulae I-VII or any combination thereof.

Figure 11 shows a non-limiting example of a strategy used to identify chemically modified siNA constructs of the invention that are nuclease resistance while preserving the ability to mediate RNAi activity. Chemical modifications are introduced into the siNA construct based on educated design parameters (e.g. introducing 2'-modifications, base modifications, backbone modifications, terminal cap modifications etc). The modified construct is tested in an appropriate system (e.g. human serum for nuclease resistance, shown, or an animal model for PK/delivery parameters). In parallel, the siNA construct is tested for RNAi activity, for example in a cell culture system such as a luciferase reporter assay). Lead siNA constructs are then identified which possess a particular characteristic while maintaining RNAi activity, and can be further modified and assayed once again. This same approach can be used to identify siNA-conjugate molecules with improved pharmacokinetic profiles, delivery, and RNAi activity.

Figure 12 shows a non-limiting example of siNA mediated inhibition of VEGF-induced angiogenesis using the rat corneal model of angiogenesis. siNA targeting site 2340 of VEGFr1 RNA 29695/29699 (shown as RPI No. sense strand/antisense strand) was compared to an inverted control siNA 29983/29984 (shown as RPI No. sense strand/antisense strand) at three different concentrations (1ug, 3ug, and 10ug) and compared to a VEGF control in which no siNA was administered. As shown in the Figure, siNA



constructs targeting VEGFr1 RNA can provide significant inhibition of angiogenesis in the rat corneal model.

Figure 13 shows a non-limiting example of reduction of VEGFr1 mRNA in A375 cells mediated by chemically-modified siNAs that target VEGFr1 mRNA. A549 cells were transfected with 0.25 ug/well of lipid complexed with 25 nM siNA. A screen of siNA constructs (Stabilization "Stab" chemistries are shown in Table IV, constructs are referred to by RPI number, see Table III) comprising Stab 4/5 chemistry (RPI 31190/31193), Stab 1/2 chemistry (RPI 31183/31186 and RPI 31184/31187), and unmodified RNA (RPI 30075/30076) were compared to untreated cells, matched chemistry inverted control siNA constructs, (RPI 31208/31211, RPI 31201/31204, RPI 31202/31205, and RPI 30077/30078) scrambled siNA control constructs (Scram1 and Scram2), and cells transfected with lipid alone (transfection control). All of the siNA constructs show significant reduction of VEGFr1 RNA expression.

## DETAILED DESCRIPTION OF THE INVENTION

### 15 Mechanism of action of Nucleic Acid Molecules of the Invention

The discussion that follows discusses the proposed mechanism of RNA interference mediated by short interfering RNA as is presently known, and is not meant to be limiting and is not an admission of prior art. Applicant demonstrates herein that chemically-modified short interfering nucleic acids possess similar or improved capacity to mediate RNAi as do siRNA molecules and are expected to possess improved stability and activity *in vivo*; therefore, this discussion is not meant to be limiting only to siRNA and can be applied to siNA as a whole. By "improved capacity to mediate RNAi" or "improved RNAi activity" is meant to include RNAi activity measured *in vitro* and/or *in vivo* where the RNAi activity is a reflection of both the ability of the siNA to mediate RNAi and the stability of the siNAs of the invention. In this invention, the product of these activities can be increased *in vitro* and/or *in vivo* compared to an all RNA siRNA or a siNA containing a plurality of ribonucleotides. In some cases, the activity or stability of the siNA molecule can be

decreased (i.e., less than ten-fold), but the overall activity of the siNA molecule is enhanced *in vitro* and/or *in vivo*.

RNA interference refers to the process of sequence specific post-transcriptional gene silencing in animals mediated by short interfering RNAs (siRNAs) (Fire *et al.*, 1998, *Nature*, 391, 806). The corresponding process in plants is commonly referred to as post-transcriptional gene silencing or RNA silencing and is also referred to as quelling in fungi. The process of post-transcriptional gene silencing is thought to be an evolutionarily-conserved cellular defense mechanism used to prevent the expression of foreign genes which is commonly shared by diverse flora and phyla (Fire *et al.*, 1999, *Trends Genet.*, 15, 358). Such protection from foreign gene expression may have evolved in response to the production of double-stranded RNAs (dsRNAs) derived from viral infection or the random integration of transposon elements into a host genome via a cellular response that specifically destroys homologous single-stranded RNA or viral genomic RNA. The presence of dsRNA in cells triggers the RNAi response through a mechanism that has yet to be fully characterized. This mechanism appears to be different from the interferon response that results from dsRNA-mediated activation of protein kinase PKR and 2', 5'-oligoadenylate synthetase resulting in non-specific cleavage of mRNA by ribonuclease L.

The presence of long dsRNAs in cells stimulates the activity of a ribonuclease III enzyme referred to as Dicer. Dicer is involved in the processing of the dsRNA into short pieces of dsRNA known as short interfering RNAs (siRNAs) (Bernstein *et al.*, 2001, *Nature*, 409, 363). Short interfering RNAs derived from Dicer activity are typically about 21 to about 23 nucleotides in length and comprise about 19 base pair duplexes. Dicer has also been implicated in the excision of 21- and 22-nucleotide small temporal RNAs (stRNAs) from precursor RNA of conserved structure that are implicated in translational control (Hutvagner *et al.*, 2001, *Science*, 293, 834). The RNAi response also features an endonuclease complex containing a siRNA, commonly referred to as an RNA-induced silencing complex (RISC), which mediates cleavage of single-stranded RNA having sequence homologous to the siRNA. Cleavage of the target RNA takes place in the middle of the region complementary to the guide sequence of the siRNA duplex (Elbashir *et al.*, 2001, *Genes Dev.*, 15, 188). In addition, RNA interference can also involve small RNA

(e.g., micro-RNA or miRNA) mediated gene silencing, presumably through cellular mechanisms that regulate chromatin structure and thereby prevent transcription of target gene sequences (see for example Allshire, 2002, *Science*, 297, 1818-1819; Volpe *et al.*, 2002, *Science*, 297, 1833-1837; Jenuwein, 2002, *Science*, 297, 2215-2218; and Hall *et al.*, 5 2002, *Science*, 297, 2232-2237). As such, siRNA molecules of the invention can be used to mediate gene silencing via interaction with RNA transcripts or alternately by interaction with particular gene sequences, wherein such interaction results in gene silencing either at the transcriptional level or post-transcriptional level.

RNAi has been studied in a variety of systems. Fire *et al.*, 1998, *Nature*, 391, 806, 10 were the first to observe RNAi in *C. elegans*. Wianny and Goetz, 1999, *Nature Cell Biol.*, 2, 70, describe RNAi mediated by dsRNA in mouse embryos. Hammond *et al.*, 2000, *Nature*, 404, 293, describe RNAi in *Drosophila* cells transfected with dsRNA. Elbashir *et al.*, 2001, *Nature*, 411, 494, describe RNAi induced by introduction of duplexes of synthetic 21-nucleotide RNAs in cultured mammalian cells including human embryonic kidney and HeLa 15 cells. Recent work in *Drosophila* embryonic lysates has revealed certain requirements for siRNA length, structure, chemical composition, and sequence that are essential to mediate efficient RNAi activity. These studies have shown that 21 nucleotide siRNA duplexes are most active when containing two 2-nucleotide 3'-terminal nucleotide overhangs. Furthermore, substitution of one or both siRNA strands with 2'-deoxy or 2'-O-methyl 20 nucleotides abolishes RNAi activity, whereas substitution of 3'-terminal siRNA nucleotides with deoxy nucleotides was shown to be tolerated. Mismatch sequences in the center of the siRNA duplex were also shown to abolish RNAi activity. In addition, these studies also indicate that the position of the cleavage site in the target RNA is defined by the 5'-end of the siRNA guide sequence rather than the 3'-end (Elbashir *et al.*, 2001, *EMBO J.*, 20, 6877). 25 Other studies have indicated that a 5'-phosphate on the target-complementary strand of a siRNA duplex is required for siRNA activity and that ATP is utilized to maintain the 5'-phosphate moiety on the siRNA (Nykanen *et al.*, 2001, *Cell*, 107, 309); however, siRNA molecules lacking a 5'-phosphate are active when introduced exogenously, suggesting that 5'-phosphorylation of siRNA constructs may occur *in vivo*.

### Synthesis of Nucleic acid Molecules

Synthesis of nucleic acids greater than 100 nucleotides in length is difficult using automated methods, and the therapeutic cost of such molecules is prohibitive. In this invention, small nucleic acid motifs ("small" refers to nucleic acid motifs no more than 100  
5 nucleotides in length, preferably no more than 80 nucleotides in length, and most preferably no more than 50 nucleotides in length; e.g., individual siNA oligonucleotide sequences or siNA sequences synthesized in tandem) are preferably used for exogenous delivery. The simple structure of these molecules increases the ability of the nucleic acid to invade targeted regions of protein and/or RNA structure. Exemplary molecules of the instant  
10 invention are chemically synthesized, and others can similarly be synthesized.

Oligonucleotides (e.g., certain modified oligonucleotides or portions of oligonucleotides lacking ribonucleotides) are synthesized using protocols known in the art, for example as described in Caruthers *et al.*, 1992, *Methods in Enzymology* 211, 3-19, Thompson *et al.*, International PCT Publication No. WO 99/54459, Wincott *et al.*, 1995,  
15 *Nucleic Acids Res.* 23, 2677-2684, Wincott *et al.*, 1997, *Methods Mol. Bio.*, 74, 59, Brennan *et al.*, 1998, *Biotechnol Bioeng.*, 61, 33-45, and Brennan, U.S. Pat. No. 6,001,311. All of these references are incorporated herein by reference. The synthesis of oligonucleotides makes use of common nucleic acid protecting and coupling groups, such as dimethoxytrityl at the 5'-end, and phosphoramidites at the 3'-end. In a non-limiting example, small scale  
20 syntheses are conducted on a 394 Applied Biosystems, Inc. synthesizer using a 0.2  $\mu$ mol scale protocol with a 2.5 min coupling step for 2'-O-methylated nucleotides and a 45 sec coupling step for 2'-deoxy nucleotides or 2'-deoxy-2'-fluoro nucleotides. Table V outlines the amounts and the contact times of the reagents used in the synthesis cycle. Alternatively, syntheses at the 0.2  $\mu$ mol scale can be performed on a 96-well plate synthesizer, such as the  
25 instrument produced by Protogene (Palo Alto, CA) with minimal modification to the cycle. A 33-fold excess (60  $\mu$ L of 0.11 M = 6.6  $\mu$ mol) of 2'-O-methyl phosphoramidite and a 105-fold excess of S-ethyl tetrazole (60  $\mu$ L of 0.25 M = 15  $\mu$ mol) can be used in each coupling cycle of 2'-O-methyl residues relative to polymer-bound 5'-hydroxyl. A 22-fold excess (40  
30  $\mu$ L of 0.11 M = 4.4  $\mu$ mol) of deoxy phosphoramidite and a 70-fold excess of S-ethyl tetrazole (40  $\mu$ L of 0.25 M = 10  $\mu$ mol) can be used in each coupling cycle of deoxy residues

relative to polymer-bound 5'-hydroxyl. Average coupling yields on the 394 Applied Biosystems, Inc. synthesizer, determined by colorimetric quantitation of the trityl fractions, are typically 97.5-99%. Other oligonucleotide synthesis reagents for the 394 Applied Biosystems, Inc. synthesizer include the following: detritylation solution is 3% TCA in  
5 methylene chloride (ABI); capping is performed with 16% *N*-methyl imidazole in THF (ABI) and 10% acetic anhydride/10% 2,6-lutidine in THF (ABI); and oxidation solution is 16.9 mM I<sub>2</sub>, 49 mM pyridine, 9% water in THF (PERSEPTIVE™). Burdick & Jackson Synthesis Grade acetonitrile is used directly from the reagent bottle. S-Ethyltetrazole solution (0.25 M in acetonitrile) is made up from the solid obtained from American  
10 International Chemical, Inc. Alternately, for the introduction of phosphorothioate linkages, Beaucage reagent (3H-1,2-Benzodithiol-3-one 1,1-dioxide, 0.05 M in acetonitrile) is used.

Deprotection of the DNA-based oligonucleotides is performed as follows: the polymer-bound trityl-on oligoribonucleotide is transferred to a 4 mL glass screw top vial and suspended in a solution of 40% aq. methylamine (1 mL) at 65 °C for 10 min. After cooling  
15 to -20 °C, the supernatant is removed from the polymer support. The support is washed three times with 1.0 mL of EtOH:MeCN:H<sub>2</sub>O/3:1:1, vortexed and the supernatant is then added to the first supernatant. The combined supernatants, containing the oligoribonucleotide, are dried to a white powder.

The method of synthesis used for RNA including certain siNA molecules of the  
20 invention follows the procedure as described in Usman *et al.*, 1987, *J. Am. Chem. Soc.*, 109, 7845; Scaringe *et al.*, 1990, *Nucleic Acids Res.*, 18, 5433; and Wincott *et al.*, 1995, *Nucleic Acids Res.* 23, 2677-2684 Wincott *et al.*, 1997, *Methods Mol. Bio.*, 74, 59, and makes use of common nucleic acid protecting and coupling groups, such as dimethoxytrityl at the 5'-end, and phosphoramidites at the 3'-end. In a non-limiting example, small scale syntheses are  
25 conducted on a 394 Applied Biosystems, Inc. synthesizer using a 0.2 μmol scale protocol with a 7.5 min coupling step for alkylsilyl protected nucleotides and a 2.5 min coupling step for 2'-O-methylated nucleotides. Table V outlines the amounts and the contact times of the reagents used in the synthesis cycle. Alternatively, syntheses at the 0.2 μmol scale can be done on a 96-well plate synthesizer, such as the instrument produced by Protogene (Palo  
30 Alto, CA) with minimal modification to the cycle. A 33-fold excess (60 μL of 0.11 M = 6.6

μmol) of 2'-O-methyl phosphoramidite and a 75-fold excess of S-ethyl tetrazole (60 μL of 0.25 M = 15 μmol) can be used in each coupling cycle of 2'-O-methyl residues relative to polymer-bound 5'-hydroxyl. A 66-fold excess (120 μL of 0.11 M = 13.2 μmol) of alkylsilyl (ribo) protected phosphoramidite and a 150-fold excess of S-ethyl tetrazole (120 μL of 0.25 M = 30 μmol) can be used in each coupling cycle of ribo residues relative to polymer-bound 5'-hydroxyl. Average coupling yields on the 394 Applied Biosystems, Inc. synthesizer, determined by colorimetric quantitation of the trityl fractions, are typically 97.5-99%. Other oligonucleotide synthesis reagents for the 394 Applied Biosystems, Inc. synthesizer include the following: detritylation solution is 3% TCA in methylene chloride (ABI); capping is performed with 16% *N*-methyl imidazole in THF (ABI) and 10% acetic anhydride/10% 2,6-lutidine in THF (ABI); oxidation solution is 16.9 mM I<sub>2</sub>, 49 mM pyridine, 9% water in THF (PERSEPTIVE™). Burdick & Jackson Synthesis Grade acetonitrile is used directly from the reagent bottle. S-Ethyltetrazole solution (0.25 M in acetonitrile) is made up from the solid obtained from American International Chemical, Inc. Alternately, for the introduction of phosphorothioate linkages, Beaucage reagent (3H-1,2-Benzodithiol-3-one 1,1-dioxide 0.05 M in acetonitrile) is used.

Deprotection of the RNA is performed using either a two-pot or one-pot protocol. For the two-pot protocol, the polymer-bound trityl-on oligoribonucleotide is transferred to a 4 mL glass screw top vial and suspended in a solution of 40% aq. methylamine (1 mL) at 65 °C for 10 min. After cooling to -20 °C, the supernatant is removed from the polymer support. The support is washed three times with 1.0 mL of EtOH:MeCN:H<sub>2</sub>O/3:1:1, vortexed and the supernatant is then added to the first supernatant. The combined supernatants, containing the oligoribonucleotide, are dried to a white powder. The base deprotected oligoribonucleotide is resuspended in anhydrous TEA/HF/NMP solution (300 μL of a solution of 1.5 mL *N*-methylpyrrolidinone, 750 μL TEA and 1 mL TEA•3HF to provide a 1.4 M HF concentration) and heated to 65 °C. After 1.5 h, the oligomer is quenched with 1.5 M NH<sub>4</sub>HCO<sub>3</sub>.

Alternatively, for the one-pot protocol, the polymer-bound trityl-on oligoribonucleotide is transferred to a 4 mL glass screw top vial and suspended in a solution

of 33% ethanolic methylamine/DMSO: 1/1 (0.8 mL) at 65 °C for 15 min. The vial is brought to rt. TEA•3HF (0.1 mL) is added and the vial is heated at 65 °C for 15 min. The sample is cooled at -20 °C and then quenched with 1.5 M NH<sub>4</sub>HCO<sub>3</sub>.

For purification of the trityl-on oligomers, the quenched NH<sub>4</sub>HCO<sub>3</sub> solution is loaded onto a C-18 containing cartridge that had been prewashed with acetonitrile followed by 50 mM TEAA. After washing the loaded cartridge with water, the RNA is detritylated with 0.5% TFA for 13 min. The cartridge is then washed again with water, salt exchanged with 1 M NaCl and washed with water again. The oligonucleotide is then eluted with 30% acetonitrile.

The average stepwise coupling yields are typically >98% (Wincott *et al.*, 1995 *Nucleic Acids Res.* 23, 2677-2684). Those of ordinary skill in the art will recognize that the scale of synthesis can be adapted to be larger or smaller than the example described above including but not limited to 96-well format.

Alternatively, the nucleic acid molecules of the present invention can be synthesized separately and joined together post-synthetically, for example, by ligation (Moore *et al.*, 1992, *Science* 256, 9923; Draper *et al.*, International PCT publication No. WO 93/23569; Shabarova *et al.*, 1991, *Nucleic Acids Research* 19, 4247; Bellon *et al.*, 1997, *Nucleosides & Nucleotides*, 16, 951; Bellon *et al.*, 1997, *Bioconjugate Chem.* 8, 204), or by hybridization following synthesis and/or deprotection.

The siNA molecules of the invention can also be synthesized via a tandem synthesis methodology as described in Example 1 herein, wherein both siNA strands are synthesized as a single contiguous oligonucleotide fragment or strand separated by a cleavable linker which is subsequently cleaved to provide separate siNA fragments or strands that hybridize and permit purification of the siNA duplex. The linker can be a polynucleotide linker or a non-nucleotide linker. The tandem synthesis of siNA as described herein can be readily adapted to both multiwell/multiplate synthesis platforms such as 96 well or similarly larger multi-well platforms. The tandem synthesis of siNA as described herein can also be readily

adapted to large scale synthesis platforms employing batch reactors, synthesis columns and the like.

A siNA molecule can also be assembled from two distinct nucleic acid strands or fragments wherein one fragment includes the sense region and the second fragment includes the antisense region of the RNA molecule.

The nucleic acid molecules of the present invention can be modified extensively to enhance stability by modification with nuclease resistant groups, for example, 2'-amino, 2'-C-allyl, 2'-fluoro, 2'-O-methyl, 2'-H (for a review see Usman and Cedergren, 1992, *TIBS* 17, 34; Usman *et al.*, 1994, *Nucleic Acids Symp. Ser.* 31, 163). siNA constructs can be purified by gel electrophoresis using general methods or can be purified by high pressure liquid chromatography (HPLC; see Wincott *et al.*, *supra*, the totality of which is hereby incorporated herein by reference) and re-suspended in water.

In another aspect of the invention, siNA molecules of the invention are expressed from transcription units inserted into DNA or RNA vectors. The recombinant vectors can be DNA plasmids or viral vectors. siNA expressing viral vectors can be constructed based on, but not limited to, adeno-associated virus, retrovirus, adenovirus, or alphavirus. The recombinant vectors capable of expressing the siNA molecules can be delivered as described herein, and persist in target cells. Alternatively, viral vectors can be used that provide for transient expression of siNA molecules.

#### Optimizing Activity of the nucleic acid molecule of the invention.

Chemically synthesizing nucleic acid molecules with modifications (base, sugar and/or phosphate) can prevent their degradation by serum ribonucleases, which can increase their potency (see *e.g.*, Eckstein *et al.*, International Publication No. WO 92/07065; Perrault *et al.*, 1990 *Nature* 344, 565; Pieken *et al.*, 1991, *Science* 253, 314; Usman and Cedergren, 1992, *Trends in Biochem. Sci.* 17, 334; Usman *et al.*, International Publication No. WO 93/15187; and Rossi *et al.*, International Publication No. WO 91/03162; Sproat, U.S. Pat. No. 5,334,711; Gold *et al.*, U.S. Pat. No. 6,300,074; and Burgin *et al.*, *supra*; all of which are incorporated by reference herein). All of the above references describe various chemical



modifications that can be made to the base, phosphate and/or sugar moieties of the nucleic acid molecules described herein. Modifications that enhance their efficacy in cells, and removal of bases from nucleic acid molecules to shorten oligonucleotide synthesis times and reduce chemical requirements are desired.

5        There are several examples in the art describing sugar, base and phosphate modifications that can be introduced into nucleic acid molecules with significant enhancement in their nuclease stability and efficacy. For example, oligonucleotides are modified to enhance stability and/or enhance biological activity by modification with  
10        nuclease resistant groups, for example, 2'-amino, 2'-C-allyl, 2'-fluoro, 2'-O-methyl, 2'-O-allyl, 2'-H, nucleotide base modifications (for a review see Usman and Cedergren, 1992, *TIBS*, 17, 34; Usman *et al.*, 1994, *Nucleic Acids Symp. Ser.* 31, 163; Burgin *et al.*, 1996, *Biochemistry*, 35, 14090). Sugar modification of nucleic acid molecules have been extensively described in the art (see Eckstein *et al.*, *International Publication* PCT No. WO 92/07065; Perrault *et al.* *Nature*, 1990, 344, 565-568; Pieken *et al.* *Science*, 1991, 253,  
15        314-317; Usman and Cedergren, *Trends in Biochem. Sci.*, 1992, 17, 334-339; Usman *et al.* *International Publication* PCT No. WO 93/15187; Sproat, *U.S. Pat.* No. 5,334,711 and Beigelman *et al.*, 1995, *J. Biol. Chem.*, 270, 25702; Beigelman *et al.*, *International PCT* publication No. WO 97/26270; Beigelman *et al.*, *U.S. Pat.* No. 5,716,824; Usman *et al.*, *U.S. Pat.* No. 5,627,053; Woolf *et al.*, *International PCT Publication* No. WO 98/13526;  
20        Thompson *et al.*, *USSN* 60/082,404 which was filed on April 20, 1998; Karpeisky *et al.*, 1998, *Tetrahedron Lett.*, 39, 1131; Earnshaw and Gait, 1998, *Biopolymers (Nucleic Acid Sciences)*, 48, 39-55; Verma and Eckstein, 1998, *Annu. Rev. Biochem.*, 67, 99-134; and Burlina *et al.*, 1997, *Bioorg. Med. Chem.*, 5, 1999-2010; all of the references are hereby incorporated in their totality by reference herein). Such publications describe general  
25        methods and strategies to determine the location of incorporation of sugar, base and/or phosphate modifications and the like into nucleic acid molecules without modulating catalysis, and are incorporated by reference herein. In view of such teachings, similar modifications can be used as described herein to modify the siNA nucleic acid molecules of the instant invention so long as the ability of siNA to promote RNAi in cells is not  
30        significantly inhibited.

While chemical modification of oligonucleotide internucleotide linkages with phosphorothioate, phosphorodithioate, and/or 5'-methylphosphonate linkages improves stability, excessive modifications can cause some toxicity or decreased activity. Therefore, when designing nucleic acid molecules, the amount of these internucleotide linkages should be minimized. The reduction in the concentration of these linkages should lower toxicity, resulting in increased efficacy and higher specificity of these molecules.

Short interfering nucleic acid (siNA) molecules having chemical modifications that maintain or enhance activity are provided. Such a nucleic acid is also generally more resistant to nucleases than an unmodified nucleic acid. Accordingly, the *in vitro* and/or *in vivo* activity should not be significantly lowered. In cases in which modulation is the goal, therapeutic nucleic acid molecules delivered exogenously should optimally be stable within cells until translation of the target RNA has been modulated long enough to reduce the levels of the undesirable protein. This period of time varies between hours to days depending upon the disease state. Improvements in the chemical synthesis of RNA and DNA (Wincott *et al.*, 1995, *Nucleic Acids Res.* 23, 2677; Caruthers *et al.*, 1992, *Methods in Enzymology* 211,3-19 (incorporated by reference herein)) have expanded the ability to modify nucleic acid molecules by introducing nucleotide modifications to enhance their nuclease stability, as described above.

In one embodiment, nucleic acid molecules of the invention include one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) G-clamp nucleotides. A G-clamp nucleotide is a modified cytosine analog wherein the modifications confer the ability to hydrogen bond both Watson-Crick and Hoogsteen faces of a complementary guanine within a duplex, see for example Lin and Matteucci, 1998, *J. Am. Chem. Soc.*, 120, 8531-8532. A single G-clamp analog substitution within an oligonucleotide can result in substantially enhanced helical thermal stability and mismatch discrimination when hybridized to complementary oligonucleotides. The inclusion of such nucleotides in nucleic acid molecules of the invention results in both enhanced affinity and specificity to nucleic acid targets, complementary sequences, or template strands. In another embodiment, nucleic acid molecules of the invention include one or more (*e.g.*, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) LNA "locked nucleic acid" nucleotides such as a 2', 4'-C methylene bicyclo

nucleotide (see for example Wengel *et al.*, International PCT Publication No. WO 00/66604 and WO 99/14226).

In another embodiment, the invention features conjugates and/or complexes of siNA molecules of the invention. Such conjugates and/or complexes can be used to facilitate delivery of siNA molecules into a biological system, such as a cell. The conjugates and complexes provided by the instant invention can impart therapeutic activity by transferring therapeutic compounds across cellular membranes, altering the pharmacokinetics, and/or modulating the localization of nucleic acid molecules of the invention. The present invention encompasses the design and synthesis of novel conjugates and complexes for the delivery of molecules, including, but not limited to, small molecules, lipids, phospholipids, nucleosides, nucleotides, nucleic acids, antibodies, toxins, negatively charged polymers and other polymers, for example proteins, peptides, hormones, carbohydrates, polyethylene glycols, or polyamines, across cellular membranes. In general, the transporters described are designed to be used either individually or as part of a multi-component system, with or without degradable linkers. These compounds are expected to improve delivery and/or localization of nucleic acid molecules of the invention into a number of cell types originating from different tissues, in the presence or absence of serum (see Sullenger and Cech, U.S. Pat. No. 5,854,038). Conjugates of the molecules described herein can be attached to biologically active molecules via linkers that are biodegradable, such as biodegradable nucleic acid linker molecules.

The term "biodegradable linker" as used herein, refers to a nucleic acid or non-nucleic acid linker molecule that is designed as a biodegradable linker to connect one molecule to another molecule, for example, a biologically active molecule to a siNA molecule of the invention or the sense and antisense strands of a siNA molecule of the invention. The biodegradable linker is designed such that its stability can be modulated for a particular purpose, such as delivery to a particular tissue or cell type. The stability of a nucleic acid-based biodegradable linker molecule can be modulated by using various chemistries, for example combinations of ribonucleotides, deoxyribonucleotides, and chemically-modified nucleotides, such as 2'-O-methyl, 2'-fluoro, 2'-amino, 2'-O-amino, 2'-C-allyl, 2'-O-allyl, and other 2'-modified or base modified nucleotides. The biodegradable nucleic acid linker

molecule can be a dimer, trimer, tetramer or longer nucleic acid molecule, for example, an oligonucleotide of about 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 nucleotides in length, or can comprise a single nucleotide with a phosphorus-based linkage, for example, a phosphoramidate or phosphodiester linkage. The biodegradable nucleic acid linker molecule can also comprise nucleic acid backbone, nucleic acid sugar, or nucleic acid base modifications.

The term "biodegradable" as used herein, refers to degradation in a biological system, for example enzymatic degradation or chemical degradation.

The term "biologically active molecule" as used herein, refers to compounds or molecules that are capable of eliciting or modifying a biological response in a system. Non-limiting examples of biologically active siNA molecules either alone or in combination with other molecules contemplated by the instant invention include therapeutically active molecules such as antibodies, hormones, antivirals, peptides, proteins, chemotherapeutics, small molecules, vitamins, co-factors, nucleosides, nucleotides, oligonucleotides, enzymatic nucleic acids, antisense nucleic acids, triplex forming oligonucleotides, 2,5-A chimeras, siNA, dsRNA, allozymes, aptamers, decoys and analogs thereof. Biologically active molecules of the invention also include molecules capable of modulating the pharmacokinetics and/or pharmacodynamics of other biologically active molecules, for example, lipids and polymers such as polyamines, polyamides, polyethylene glycol and other polyethers.

The term "phospholipid" as used herein, refers to a hydrophobic molecule comprising at least one phosphorus group. For example, a phospholipid can comprise a phosphorus-containing group and saturated or unsaturated alkyl group, optionally substituted with OH, COOH, oxo, amine, or substituted or unsubstituted aryl groups.

Therapeutic nucleic acid molecules (e.g., siNA molecules) delivered exogenously optimally are stable within cells until reverse transcription of the RNA has been modulated long enough to reduce the levels of the RNA transcript. The nucleic acid molecules are resistant to nucleases in order to function as effective intracellular therapeutic agents.

Improvements in the chemical synthesis of nucleic acid molecules described in the instant invention and in the art have expanded the ability to modify nucleic acid molecules by introducing nucleotide modifications to enhance their nuclease stability as described above.

In yet another embodiment, siNA molecules having chemical modifications that  
5 maintain or enhance enzymatic activity of proteins involved in RNAi are provided. Such nucleic acids are also generally more resistant to nucleases than unmodified nucleic acids. Thus, *in vitro* and/or *in vivo* the activity should not be significantly lowered.

Use of the nucleic acid-based molecules of the invention will lead to better treatment of the disease progression by affording the possibility of combination therapies (*e.g.*,  
10 multiple siNA molecules targeted to different genes; nucleic acid molecules coupled with known small molecule modulators; or intermittent treatment with combinations of molecules, including different motifs and/or other chemical or biological molecules). The treatment of subjects with siNA molecules can also include combinations of different types of nucleic acid molecules, such as enzymatic nucleic acid molecules (ribozymes),  
15 allozymes, antisense, 2,5-A oligoadenylate, decoys, and aptamers.

In another aspect a siNA molecule of the invention comprises one or more 5' and/or a 3'-cap structure, for example on only the sense siNA strand, the antisense siNA strand, or both siNA strands.

By "cap structure" is meant chemical modifications, which have been incorporated at  
20 either terminus of the oligonucleotide (see, for example, Adamic *et al.*, U.S. Pat. No. 5,998,203, incorporated by reference herein). These terminal modifications protect the nucleic acid molecule from exonuclease degradation, and may help in delivery and/or localization within a cell. The cap may be present at the 5'-terminus (5'-cap) or at the 3'-terminal (3'-cap) or may be present on both termini. In non-limiting examples, the 5'-cap is  
25 selected from the group consisting of glyceryl, inverted deoxy abasic residue (moiety); 4',5'-methylene nucleotide; 1-(beta-D-erythrofuransyl) nucleotide, 4'-thio nucleotide; carbocyclic nucleotide; 1,5-anhydrohexitol nucleotide; L-nucleotides; alpha-nucleotides; modified base nucleotide; phosphorodithioate linkage; *threo*-pentofuransyl nucleotide;

acyclic 3',4'-seco nucleotide; acyclic 3,4-dihydroxybutyl nucleotide; acyclic 3,5-dihydroxypentyl nucleotide, 3'-3'-inverted nucleotide moiety; 3'-3'-inverted abasic moiety; 3'-2'-inverted nucleotide moiety; 3'-2'-inverted abasic moiety; 1,4-butanediol phosphate; 3'-phosphoramidate; hexylphosphate; aminohexyl phosphate; 3'-phosphate; 3'-phosphorothioate; phosphorodithioate; or bridging or non-bridging methylphosphonate moiety.

In non-limiting examples, the 3'-cap is selected from the group consisting of glyceryl, inverted deoxy abasic residue (moiety), 4', 5'-methylene nucleotide; 1-(beta-D-erythrofuranosyl) nucleotide; 4'-thio nucleotide, carbocyclic nucleotide; 5'-amino-alkyl phosphate; 1,3-diamino-2-propyl phosphate; 3-aminopropyl phosphate; 6-aminohexyl phosphate; 1,2-aminododecyl phosphate; hydroxypropyl phosphate; 1,5-anhydrohexitol nucleotide; L-nucleotide; alpha-nucleotide; modified base nucleotide; phosphorodithioate; *threo*-pentofuranosyl nucleotide; acyclic 3',4'-seco nucleotide; 3,4-dihydroxybutyl nucleotide; 3,5-dihydroxypentyl nucleotide, 5'-5'-inverted nucleotide moiety; 5'-5'-inverted abasic moiety; 5'-phosphoramidate; 5'-phosphorothioate; 1,4-butanediol phosphate; 5'-amino; bridging and/or non-bridging 5'-phosphoramidate, phosphorothioate and/or phosphorodithioate, bridging or non bridging methylphosphonate and 5'-mercapto moieties (for more details see Beaucage and Iyer, 1993, *Tetrahedron* 49, 1925; incorporated by reference herein).

By the term "non-nucleotide" is meant any group or compound which can be incorporated into a nucleic acid chain in the place of one or more nucleotide units, including either sugar and/or phosphate substitutions, and allows the remaining bases to exhibit their enzymatic activity. The group or compound is abasic in that it does not contain a commonly recognized nucleotide base, such as adenosine, guanine, cytosine, uracil or thymine and therefore lacks a base at the 1'-position.

An "alkyl" group refers to a saturated aliphatic hydrocarbon, including straight-chain, branched-chain, and cyclic alkyl groups. Preferably, the alkyl group has 1 to 12 carbons. More preferably, it is a lower alkyl of from 1 to 7 carbons, more preferably 1 to 4 carbons. The alkyl group can be substituted or unsubstituted. When substituted the substituted

group(s) is preferably, hydroxyl, cyano, alkoxy, =O, =S, NO<sub>2</sub> or N(CH<sub>3</sub>)<sub>2</sub>, amino, or SH. The term also includes alkenyl groups that are unsaturated hydrocarbon groups containing at least one carbon-carbon double bond, including straight-chain, branched-chain, and cyclic groups. Preferably, the alkenyl group has 1 to 12 carbons. More preferably, it is a lower alkenyl of from 1 to 7 carbons, more preferably 1 to 4 carbons. The alkenyl group may be substituted or unsubstituted. When substituted the substituted group(s) is preferably, hydroxyl, cyano, alkoxy, =O, =S, NO<sub>2</sub>, halogen, N(CH<sub>3</sub>)<sub>2</sub>, amino, or SH. The term "alkyl" also includes alkynyl groups that have an unsaturated hydrocarbon group containing at least one carbon-carbon triple bond, including straight-chain, branched-chain, and cyclic groups. Preferably, the alkynyl group has 1 to 12 carbons. More preferably, it is a lower alkynyl of from 1 to 7 carbons, more preferably 1 to 4 carbons. The alkynyl group may be substituted or unsubstituted. When substituted the substituted group(s) is preferably, hydroxyl, cyano, alkoxy, =O, =S, NO<sub>2</sub> or N(CH<sub>3</sub>)<sub>2</sub>, amino or SH.

Such alkyl groups can also include aryl, alkylaryl, carbocyclic aryl, heterocyclic aryl, amide and ester groups. An "aryl" group refers to an aromatic group that has at least one ring having a conjugated pi electron system and includes carbocyclic aryl, heterocyclic aryl and biaryl groups, all of which may be optionally substituted. The preferred substituent(s) of aryl groups are halogen, trihalomethyl, hydroxyl, SH, OH, cyano, alkoxy, alkyl, alkenyl, alkynyl, and amino groups. An "alkylaryl" group refers to an alkyl group (as described above) covalently joined to an aryl group (as described above). Carbocyclic aryl groups are groups wherein the ring atoms on the aromatic ring are all carbon atoms. The carbon atoms are optionally substituted. Heterocyclic aryl groups are groups having from 1 to 3 heteroatoms as ring atoms in the aromatic ring and the remainder of the ring atoms are carbon atoms. Suitable heteroatoms include oxygen, sulfur, and nitrogen, and include furanyl, thienyl, pyridyl, pyrrolyl, N-lower alkyl pyrrolo, pyrimidyl, pyrazinyl, imidazolyl and the like, all optionally substituted. An "amide" refers to an -C(O)-NH-R, where R is either alkyl, aryl, alkylaryl or hydrogen. An "ester" refers to an -C(O)-OR', where R is either alkyl, aryl, alkylaryl or hydrogen.

By "nucleotide" as used herein is as recognized in the art to include natural bases (standard), and modified bases well known in the art. Such bases are generally located at the 1' position of a nucleotide sugar moiety. Nucleotides generally comprise a base, sugar and a phosphate group. The nucleotides can be unmodified or modified at the sugar, phosphate and/or base moiety, (also referred to interchangeably as nucleotide analogs, modified nucleotides, non-natural nucleotides, non-standard nucleotides and other; see, for example, Usman and McSwiggen, *supra*; Eckstein *et al.*, International PCT Publication No. WO 92/07065; Usman *et al.*, International PCT Publication No. WO 93/15187; Uhlman & Peyman, *supra*, all are hereby incorporated by reference herein). There are several examples of modified nucleic acid bases known in the art as summarized by Limbach *et al.*, 1994, *Nucleic Acids Res.* 22, 2183. Some of the non-limiting examples of base modifications that can be introduced into nucleic acid molecules include, inosine, purine, pyridin-4-one, pyridin-2-one, phenyl, pseudouracil, 2, 4, 6-trimethoxy benzene, 3-methyl uracil, dihydrouridine, naphthyl, aminophenyl, 5-alkylcytidines (*e.g.*, 5-methylcytidine), 5-alkyluridines (*e.g.*, ribothymidine), 5-halouridine (*e.g.*, 5-bromouridine) or 6-azapyrimidines or 6-alkylpyrimidines (*e.g.* 6-methyluridine), propyne, and others (Burgin *et al.*, 1996, *Biochemistry*, 35, 14090; Uhlman & Peyman, *supra*). By "modified bases" in this aspect is meant nucleotide bases other than adenine, guanine, cytosine and uracil at 1' position or their equivalents.

In one embodiment, the invention features modified siNA molecules, with phosphate backbone modifications comprising one or more phosphorothioate, phosphorodithioate, methylphosphonate, phosphotriester, morpholino, amidate carbamate, carboxymethyl, acetamidate, polyamide, sulfonate, sulfonamide, sulfamate, formacetal, thioformacetal, and/or alkylsilyl, substitutions. For a review of oligonucleotide backbone modifications, see Hunziker and Leumann, 1995, *Nucleic Acid Analogues: Synthesis and Properties*, in *Modern Synthetic Methods*, VCH, 331-417, and Mesmaeker *et al.*, 1994, *Novel Backbone Replacements for Oligonucleotides*, in *Carbohydrate Modifications in Antisense Research*, ACS, 24-39.

By "abasic" is meant sugar moieties lacking a base or having other chemical groups in place of a base at the 1' position, see for example Adamic *et al.*, U.S. Pat. No. 5,998,203.



By "unmodified nucleoside" is meant one of the bases adenine, cytosine, guanine, thymine, or uracil joined to the 1' carbon of  $\beta$ -D-ribo-furanose.

By "modified nucleoside" is meant any nucleotide base which contains a modification in the chemical structure of an unmodified nucleotide base, sugar and/or phosphate. Non-limiting examples of modified nucleotides are shown by Formulae I-VII and/or other  
5 modifications described herein.

In connection with 2'-modified nucleotides as described for the present invention, by "amino" is meant 2'-NH<sub>2</sub> or 2'-O- NH<sub>2</sub>, which can be modified or unmodified. Such modified groups are described, for example, in Eckstein *et al.*, U.S. Pat. No. 5,672,695 and  
10 Matulic-Adamic *et al.*, U.S. Pat. No. 6,248,878, which are both incorporated by reference in their entireties.

Various modifications to nucleic acid siNA structure can be made to enhance the utility of these molecules. Such modifications will enhance shelf-life, half-life *in vitro*, stability, and ease of introduction of such oligonucleotides to the target site, *e.g.*, to enhance  
15 penetration of cellular membranes, and confer the ability to recognize and bind to targeted cells.

#### Administration of Nucleic Acid Molecules

A siNA molecule of the invention can be adapted for use to treat, for example, tumor angiogenesis and cancer, including but not limited to breast cancer, lung cancer (including  
20 non-small cell lung carcinoma), prostate cancer, colorectal cancer, brain cancer, esophageal cancer, bladder cancer, pancreatic cancer, cervical cancer, head and neck cancer, skin cancers, nasopharyngeal carcinoma, liposarcoma, epithelial carcinoma, renal cell carcinoma, gallbladder adeno carcinoma, parotid adenocarcinoma, ovarian cancer, melanoma, lymphoma, glioma, endometrial sarcoma, multidrug resistant cancers, diabetic retinopathy,  
25 macular degeneration, neovascular glaucoma, myopic degeneration, arthritis, psoriasis, endometriosis, female reproduction, verruca vulgaris, angiofibroma of tuberous sclerosis, pot-wine stains, Sturge Weber syndrome, Kippel-Trenaunay-Weber syndrome, Osler-Weber-Rendu syndrome, renal disease such as Autosomal dominant polycystic kidney

disease (ADPKD), and any other diseases or conditions that are related to or will respond to the levels of VEGF, VEGFr1, VEGFr2 and/or VEGFr3 in a cell or tissue, alone or in combination with other therapies. For example, a siNA molecule can comprise a delivery vehicle, including liposomes, for administration to a subject, carriers and diluents and their salts, and/or can be present in pharmaceutically acceptable formulations. Methods for the delivery of nucleic acid molecules are described in Akhtar *et al.*, 1992, *Trends Cell Bio.*, 2, 139; *Delivery Strategies for Antisense Oligonucleotide Therapeutics*, ed. Akhtar, 1995, Maurer *et al.*, 1999, *Mol. Membr. Biol.*, 16, 129-140; Hofland and Huang, 1999, *Handb. Exp. Pharmacol.*, 137, 165-192; and Lee *et al.*, 2000, *ACS Symp. Ser.*, 752, 184-192, all of which are incorporated herein by reference. Beigelman *et al.*, U.S. Pat. No. 6,395,713 and Sullivan *et al.*, PCT WO 94/02595 further describe the general methods for delivery of nucleic acid molecules. These protocols can be utilized for the delivery of virtually any nucleic acid molecule. Nucleic acid molecules can be administered to cells by a variety of methods known to those of skill in the art, including, but not restricted to, encapsulation in liposomes, by iontophoresis, or by incorporation into other vehicles, such as hydrogels, cyclodextrins (see for example Gonzalez *et al.*, 1999, *Bioconjugate Chem.*, 10, 1068-1074), biodegradable nanocapsules, and bioadhesive microspheres, or by proteinaceous vectors (O'Hare and Normand, International PCT Publication No. WO 00/53722). Alternatively, the nucleic acid/vehicle combination is locally delivered by direct injection or by use of an infusion pump. Direct injection of the nucleic acid molecules of the invention, whether subcutaneous, intramuscular, or intradermal, can take place using standard needle and syringe methodologies, or by needle-free technologies such as those described in Comry *et al.*, 1999, *Clin. Cancer Res.*, 5, 2330-2337 and Barry *et al.*, International PCT Publication No. WO 99/31262. The molecules of the instant invention can be used as pharmaceutical agents. Pharmaceutical agents prevent, modulate the occurrence, or treat (alleviate a symptom to some extent, preferably all of the symptoms) of a disease state in a subject.

Thus, the invention features a pharmaceutical composition comprising one or more nucleic acid(s) of the invention in an acceptable carrier, such as a stabilizer, buffer, and the like. The polynucleotides of the invention can be administered (*e.g.*, RNA, DNA or protein) and introduced into a subject by any standard means, with or without stabilizers, buffers, and

the like, to form a pharmaceutical composition. When it is desired to use a liposome delivery mechanism, standard protocols for formation of liposomes can be followed. The compositions of the present invention can also be formulated and used as tablets, capsules or elixirs for oral administration, suppositories for rectal administration, sterile solutions, suspensions for injectable administration, and the other compositions known in the art.

The present invention also includes pharmaceutically acceptable formulations of the compounds described. These formulations include salts of the above compounds, *e.g.*, acid addition salts, for example, salts of hydrochloric, hydrobromic, acetic acid, and benzene sulfonic acid.

A pharmacological composition or formulation refers to a composition or formulation in a form suitable for administration, *e.g.*, systemic administration, into a cell or subject, including for example a human. Suitable forms, in part, depend upon the use or the route of entry, for example oral, transdermal, or by injection. Such forms should not prevent the composition or formulation from reaching a target cell (*i.e.*, a cell to which the negatively charged nucleic acid is desirable for delivery). For example, pharmacological compositions injected into the blood stream should be soluble. Other factors are known in the art, and include considerations such as toxicity and forms that prevent the composition or formulation from exerting its effect.

By "systemic administration" is meant *in vivo* systemic absorption or accumulation of drugs in the blood stream followed by distribution throughout the entire body. Administration routes that lead to systemic absorption include, without limitation: intravenous, subcutaneous, intraperitoneal, inhalation, oral, intrapulmonary and intramuscular. Each of these administration routes exposes the siNA molecules of the invention to an accessible diseased tissue. The rate of entry of a drug into the circulation has been shown to be a function of molecular weight or size. The use of a liposome or other drug carrier comprising the compounds of the instant invention can potentially localize the drug, for example, in certain tissue types, such as the tissues of the reticular endothelial system (RES). A liposome formulation that can facilitate the association of drug with the surface of cells, such as, lymphocytes and macrophages is also useful. This approach can

provide enhanced delivery of the drug to target cells by taking advantage of the specificity of macrophage and lymphocyte immune recognition of abnormal cells, such as cells producing excess VEGF and/or VEGFr.

By "pharmaceutically acceptable formulation" is meant, a composition or formulation  
5 that allows for the effective distribution of the nucleic acid molecules of the instant invention in the physical location most suitable for their desired activity. Non-limiting examples of agents suitable for formulation with the nucleic acid molecules of the instant invention include: P-glycoprotein inhibitors (such as Pluronic P85), which can enhance entry of drugs into the CNS (Jolliet-Riant and Tillement, 1999, *Fundam. Clin. Pharmacol.*,  
10 13, 16-26); biodegradable polymers, such as poly (DL-lactide-coglycolide) microspheres for sustained release delivery after intracerebral implantation (Emerich, DF *et al.*, 1999, *Cell Transplant*, 8, 47-58) (Alkermes, Inc. Cambridge, MA); and loaded nanoparticles, such as those made of polybutylcyanoacrylate, which can deliver drugs across the blood brain barrier and can alter neuronal uptake mechanisms (*Prog Neuropsychopharmacol Biol*  
15 *Psychiatry*, 23, 941-949, 1999). Other non-limiting examples of delivery strategies for the nucleic acid molecules of the instant invention include material described in Boado *et al.*, 1998, *J. Pharm. Sci.*, 87, 1308-1315; Tyler *et al.*, 1999, *FEBS Lett.*, 421, 280-284; Pardridge *et al.*, 1995, *PNAS USA*, 92, 5592-5596; Boado, 1995, *Adv. Drug Delivery Rev.*, 15, 73-107; Aldrian-Herrada *et al.*, 1998, *Nucleic Acids Res.*, 26, 4910-4916; and Tyler *et al.*, 1999,  
20 *PNAS USA*, 96, 7053-7058.

The invention also features the use of the composition comprising surface-modified liposomes containing poly (ethylene glycol) lipids (PEG-modified, or long-circulating liposomes or stealth liposomes). These formulations offer a method for increasing the accumulation of drugs in target tissues. This class of drug carriers resists opsonization and  
25 elimination by the mononuclear phagocytic system (MPS or RES), thereby enabling longer blood circulation times and enhanced tissue exposure for the encapsulated drug (Lasic *et al.* *Chem. Rev.* 1995, 95, 2601-2627; Ishiwata *et al.*, *Chem. Pharm. Bull.* 1995, 43, 1005-1011). Such liposomes have been shown to accumulate selectively in tumors, presumably by extravasation and capture in the neovascularized target tissues (Lasic *et al.*, *Science* 1995,  
30 267, 1275-1276; Oku *et al.*, 1995, *Biochim. Biophys. Acta*, 1238, 86-90). The long-

circulating liposomes enhance the pharmacokinetics and pharmacodynamics of DNA and RNA, particularly compared to conventional cationic liposomes which are known to accumulate in tissues of the MPS (Liu *et al.*, *J. Biol. Chem.* 1995, 42, 24864-24870; Choi *et al.*, International PCT Publication No. WO 96/10391; Ansell *et al.*, International PCT Publication No. WO 96/10390; Holland *et al.*, International PCT Publication No. WO 96/10392). Long-circulating liposomes are also likely to protect drugs from nuclease degradation to a greater extent compared to cationic liposomes, based on their ability to avoid accumulation in metabolically aggressive MPS tissues such as the liver and spleen.

The present invention also includes compositions prepared for storage or administration that include a pharmaceutically effective amount of the desired compounds in a pharmaceutically acceptable carrier or diluent. Acceptable carriers or diluents for therapeutic use are well known in the pharmaceutical art, and are described, for example, in *Remington's Pharmaceutical Sciences*, Mack Publishing Co. (A.R. Gennaro edit. 1985), hereby incorporated by reference herein. For example, preservatives, stabilizers, dyes and flavoring agents can be provided. These include sodium benzoate, sorbic acid and esters of *p*-hydroxybenzoic acid. In addition, antioxidants and suspending agents can be used.

A pharmaceutically effective dose is that dose required to prevent, inhibit the occurrence, or treat (alleviate a symptom to some extent, preferably all of the symptoms) of a disease state. The pharmaceutically effective dose depends on the type of disease, the composition used, the route of administration, the type of mammal being treated, the physical characteristics of the specific mammal under consideration, concurrent medication, and other factors that those skilled in the medical arts will recognize. Generally, an amount between 0.1 mg/kg and 100 mg/kg body weight/day of active ingredients is administered dependent upon potency of the negatively charged polymer.

The nucleic acid molecules of the invention and formulations thereof can be administered orally, topically, parenterally, by inhalation or spray, or rectally in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and/or vehicles. The term parenteral as used herein includes percutaneous, subcutaneous, intravascular (*e.g.*, intravenous), intramuscular, or intrathecal injection or

infusion techniques and the like. In addition, there is provided a pharmaceutical formulation comprising a nucleic acid molecule of the invention and a pharmaceutically acceptable carrier. One or more nucleic acid molecules of the invention can be present in association with one or more non-toxic pharmaceutically acceptable carriers and/or diluents and/or  
5     adjuvants, and if desired other active ingredients. The pharmaceutical compositions containing nucleic acid molecules of the invention can be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsion, hard or soft capsules, or syrups or elixirs.

Compositions intended for oral use can be prepared according to any method known to  
10     the art for the manufacture of pharmaceutical compositions and such compositions can contain one or more such sweetening agents, flavoring agents, coloring agents or preservative agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients that are suitable for the manufacture of tablets. These excipients can  
15     be, for example, inert diluents; such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia; and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets can be uncoated or they can be coated by known techniques. In some cases such coatings can be  
20     prepared by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate can be employed.

Formulations for oral use can also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate,  
25     calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin or olive oil.

Aqueous suspensions contain the active materials in a mixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydropropyl-methylcellulose,

sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents can be a naturally-occurring phosphatide, for example, lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions can also contain one or more preservatives, for example ethyl, or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose or saccharin.

Oily suspensions can be formulated by suspending the active ingredients in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions can contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents and flavoring agents can be added to provide palatable oral preparations. These compositions can be preserved by the addition of an anti-oxidant such as ascorbic acid

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents or suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, can also be present.

Pharmaceutical compositions of the invention can also be in the form of oil-in-water emulsions. The oily phase can be a vegetable oil or a mineral oil or mixtures of these. Suitable emulsifying agents can be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol, anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for

example polyoxyethylene sorbitan monooleate. The emulsions can also contain sweetening and flavoring agents.

Syrups and elixirs can be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol, glucose or sucrose. Such formulations can also contain a demulcent, a preservative and flavoring and coloring agents. The pharmaceutical compositions can be in the form of a sterile injectable aqueous or oleaginous suspension. This suspension can be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents that have been mentioned above. The sterile injectable preparation can also be a sterile injectable solution or suspension in a non-toxic parentally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that can be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil can be employed including synthetic mono-or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

The nucleic acid molecules of the invention can also be administered in the form of suppositories, *e.g.*, for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient that is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials include cocoa butter and polyethylene glycols.

Nucleic acid molecules of the invention can be administered parenterally in a sterile medium. The drug, depending on the vehicle and concentration used, can either be suspended or dissolved in the vehicle. Advantageously, adjuvants such as local anesthetics, preservatives and buffering agents can be dissolved in the vehicle.

Dosage levels of the order of from about 0.1 mg to about 140 mg per kilogram of body weight per day are useful in the treatment of the above-indicated conditions (about 0.5 mg to about 7 g per subject per day). The amount of active ingredient that can be combined with the carrier materials to produce a single dosage form varies depending upon the host treated



and the particular mode of administration. Dosage unit forms generally contain between from about 1 mg to about 500 mg of an active ingredient.

It is understood that the specific dose level for any particular subject depends upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, and rate of excretion, drug combination and the severity of the particular disease undergoing therapy.

For administration to non-human animals, the composition can also be added to the animal feed or drinking water. It can be convenient to formulate the animal feed and drinking water compositions so that the animal takes in a therapeutically appropriate quantity of the composition along with its diet. It can also be convenient to present the composition as a premix for addition to the feed or drinking water.

The nucleic acid molecules of the present invention can also be administered to a subject in combination with other therapeutic compounds to increase the overall therapeutic effect. The use of multiple compounds to treat an indication can increase the beneficial effects while reducing the presence of side effects.

In one embodiment, the invention comprises compositions suitable for administering nucleic acid molecules of the invention to specific cell types. For example, the asialoglycoprotein receptor (ASGPr) (Wu and Wu, 1987, *J. Biol. Chem.* 262, 4429-4432) is unique to hepatocytes and binds branched galactose-terminal glycoproteins, such as asialoorosomucoid (ASOR). In another example, the folate receptor is overexpressed in many cancer cells. Binding of such glycoproteins, synthetic glycoconjugates, or folates to the receptor takes place with an affinity that strongly depends on the degree of branching of the oligosaccharide chain, for example, triantennary structures are bound with greater affinity than biantennary or monoantennary chains (Baenziger and Fiete, 1980, *Cell*, 22, 611-620; Connolly *et al.*, 1982, *J. Biol. Chem.*, 257, 939-945). Lee and Lee, 1987, *Glycoconjugate J.*, 4, 317-328, obtained this high specificity through the use of N-acetyl-D-galactosamine as the carbohydrate moiety, which has higher affinity for the receptor, compared to galactose. This "clustering effect" has also been described for the binding and uptake of mannosyl-

terminating glycoproteins or glycoconjugates (Ponpipom *et al.*, 1981, *J. Med. Chem.*, 24, 1388-1395). The use of galactose, galactosamine, or folate based conjugates to transport exogenous compounds across cell membranes can provide a targeted delivery approach to, for example, the treatment of liver disease, cancers of the liver, or other cancers. The use of  
5 bioconjugates can also provide a reduction in the required dose of therapeutic compounds required for treatment. Furthermore, therapeutic bioavailability, pharmacodynamics, and pharmacokinetic parameters can be modulated through the use of nucleic acid bioconjugates of the invention. Non-limiting examples of such bioconjugates are described in Vargeese *et al.*, USSN 10/201,394, filed August 13, 2001; and Matulic-Adamic *et al.*, USSN  
10 60/362,016, filed March 6, 2002.

Alternatively, certain siNA molecules of the instant invention can be expressed within cells from eukaryotic promoters (*e.g.*, Izant and Weintraub, 1985, *Science*, 229, 345; McGarry and Lindquist, 1986, *Proc. Natl. Acad. Sci.*, USA 83, 399; Scanlon *et al.*, 1991, *Proc. Natl. Acad. Sci. USA*, 88, 10591-5; Kashani-Sabet *et al.*, 1992, *Antisense Res. Dev.*, 2,  
15 3-15; Dropulic *et al.*, 1992, *J. Virol.*, 66, 1432-41; Weerasinghe *et al.*, 1991, *J. Virol.*, 65, 5531-4; Ojwang *et al.*, 1992, *Proc. Natl. Acad. Sci. USA*, 89, 10802-6; Chen *et al.*, 1992, *Nucleic Acids Res.*, 20, 4581-9; Sarver *et al.*, 1990 *Science*, 247, 1222-1225; Thompson *et al.*, 1995, *Nucleic Acids Res.*, 23, 2259; Good *et al.*, 1997, *Gene Therapy*, 4, 45. Those skilled in the art realize that any nucleic acid can be expressed in eukaryotic cells from the  
20 appropriate DNA/RNA vector. The activity of such nucleic acids can be augmented by their release from the primary transcript by a enzymatic nucleic acid (Draper *et al.*, PCT WO 93/23569, and Sullivan *et al.*, PCT WO 94/02595; Ohkawa *et al.*, 1992, *Nucleic Acids Symp. Ser.*, 27, 15-6; Taira *et al.*, 1991, *Nucleic Acids Res.*, 19, 5125-30; Ventura *et al.*, 1993, *Nucleic Acids Res.*, 21, 3249-55; Chowrira *et al.*, 1994, *J. Biol. Chem.*, 269, 25856.

25 In another aspect of the invention, RNA molecules of the present invention can be expressed from transcription units (see for example Couture *et al.*, 1996, *TIG.*, 12, 510) inserted into DNA or RNA vectors. The recombinant vectors can be DNA plasmids or viral vectors. siNA expressing viral vectors can be constructed based on, but not limited to, adeno-associated virus, retrovirus, adenovirus, or alphavirus. In another embodiment, pol  
30 III based constructs are used to express nucleic acid molecules of the invention (see for

example Thompson, U.S. Pats. Nos. 5,902,880 and 6,146,886). The recombinant vectors capable of expressing the siNA molecules can be delivered as described above, and persist in target cells. Alternatively, viral vectors can be used that provide for transient expression of nucleic acid molecules. Such vectors can be repeatedly administered as necessary. Once  
5 expressed, the siNA molecule interacts with the target mRNA and generates an RNAi response. Delivery of siNA molecule expressing vectors can be systemic, such as by intravenous or intra-muscular administration, by administration to target cells ex-planted from a subject followed by reintroduction into the subject, or by any other means that would allow for introduction into the desired target cell (for a review see Couture *et al.*, 1996,  
10 *TIG.*, 12, 510).

In one aspect the invention features an expression vector comprising a nucleic acid sequence encoding at least one siNA molecule of the instant invention. The expression vector can encode one or both strands of a siNA duplex, or a single self-complementary strand that self hybridizes into a siNA duplex. The nucleic acid sequences encoding the  
15 siNA molecules of the instant invention can be operably linked in a manner that allows expression of the siNA molecule (see for example Paul *et al.*, 2002, *Nature Biotechnology*, 19, 505; Miyagishi and Taira, 2002, *Nature Biotechnology*, 19, 497; Lee *et al.*, 2002, *Nature Biotechnology*, 19, 500; and Novina *et al.*, 2002, *Nature Medicine*, advance online publication doi:10.1038/nm725).

20 In another aspect, the invention features an expression vector comprising: a) a transcription initiation region (*e.g.*, eukaryotic pol I, II or III initiation region); b) a transcription termination region (*e.g.*, eukaryotic pol I, II or III termination region); and c) a nucleic acid sequence encoding at least one of the siNA molecules of the instant invention, wherein said sequence is operably linked to said initiation region and said  
25 termination region in a manner that allows expression and/or delivery of the siNA molecule. The vector can optionally include an open reading frame (ORF) for a protein operably linked on the 5' side or the 3'-side of the sequence encoding the siNA of the invention; and/or an intron (intervening sequences).

Transcription of the siNA molecule sequences can be driven from a promoter for eukaryotic RNA polymerase I (pol I), RNA polymerase II (pol II), or RNA polymerase III (pol III). Transcripts from pol II or pol III promoters are expressed at high levels in all cells; the levels of a given pol II promoter in a given cell type depends on the nature of the gene regulatory sequences (enhancers, silencers, etc.) present nearby. Prokaryotic RNA polymerase promoters are also used, providing that the prokaryotic RNA polymerase enzyme is expressed in the appropriate cells (Elroy-Stein and Moss, 1990, *Proc. Natl. Acad. Sci. U S A*, 87, 6743-7; Gao and Huang 1993, *Nucleic Acids Res.*, 21, 2867-72; Lieber *et al.*, 1993, *Methods Enzymol.*, 217, 47-66; Zhou *et al.*, 1990, *Mol. Cell. Biol.*, 10, 4529-37).

Several investigators have demonstrated that nucleic acid molecules expressed from such promoters can function in mammalian cells (e.g. Kashani-Sabet *et al.*, 1992, *Antisense Res. Dev.*, 2, 3-15; Ojwang *et al.*, 1992, *Proc. Natl. Acad. Sci. U S A*, 89, 10802-6; Chen *et al.*, 1992, *Nucleic Acids Res.*, 20, 4581-9; Yu *et al.*, 1993, *Proc. Natl. Acad. Sci. U S A*, 90, 6340-4; L'Huillier *et al.*, 1992, *EMBO J.*, 11, 4411-8; Lisiewicz *et al.*, 1993, *Proc. Natl. Acad. Sci. U. S. A*, 90, 8000-4; Thompson *et al.*, 1995, *Nucleic Acids Res.*, 23, 2259; Sullenger & Cech, 1993, *Science*, 262, 1566). More specifically, transcription units such as the ones derived from genes encoding U6 small nuclear (snRNA), transfer RNA (tRNA) and adenovirus VA RNA are useful in generating high concentrations of desired RNA molecules such as siNA in cells (Thompson *et al.*, *supra*; Couture and Stinchcomb, 1996, *supra*; Noonberg *et al.*, 1994, *Nucleic Acid Res.*, 22, 2830; Noonberg *et al.*, U.S. Pat. No. 5,624,803; Good *et al.*, 1997, *Gene Ther.*, 4, 45; Beigelman *et al.*, International PCT Publication No. WO 96/18736. The above siNA transcription units can be incorporated into a variety of vectors for introduction into mammalian cells, including but not restricted to, plasmid DNA vectors, viral DNA vectors (such as adenovirus or adeno-associated virus vectors), or viral RNA vectors (such as retroviral or alphavirus vectors) (for a review see Couture and Stinchcomb, 1996, *supra*).

In another aspect the invention features an expression vector comprising a nucleic acid sequence encoding at least one of the siNA molecules of the invention in a manner that allows expression of that siNA molecule. The expression vector comprises in one embodiment; a) a transcription initiation region; b) a transcription termination region; and c)

a nucleic acid sequence encoding at least one strand of the siNA molecule, wherein the sequence is operably linked to the initiation region and the termination region in a manner that allows expression and/or delivery of the siNA molecule.

5 In another embodiment the expression vector comprises: a) a transcription initiation region; b) a transcription termination region; c) an open reading frame; and d) a nucleic acid sequence encoding at least one strand of a siNA molecule, wherein the sequence is operably linked to the 3'-end of the open reading frame and wherein the sequence is operably linked to the initiation region, the open reading frame and the termination region in a manner that allows expression and/or delivery of the siNA molecule. In yet another embodiment, the  
10 expression vector comprises: a) a transcription initiation region; b) a transcription termination region; c) an intron; and d) a nucleic acid sequence encoding at least one siNA molecule, wherein the sequence is operably linked to the initiation region, the intron and the termination region in a manner which allows expression and/or delivery of the nucleic acid molecule.

15 In another embodiment, the expression vector comprises: a) a transcription initiation region; b) a transcription termination region; c) an intron; d) an open reading frame; and e) a nucleic acid sequence encoding at least one strand of a siNA molecule, wherein the sequence is operably linked to the 3'-end of the open reading frame and wherein the sequence is operably linked to the initiation region, the intron, the open reading frame and  
20 the termination region in a manner which allows expression and/or delivery of the siNA molecule.

#### VEGF/VEGFr biology and biochemistry

The following discussion is adapted from R&D Systems, Cytokine Mini Reviews, Vascular Endothelial Growth Factor (VEGF), Copyright ©2002 R&D Systems.  
25 Angiogenesis is a process of new blood vessel development from pre-existing vasculature. It plays an essential role in embryonic development, normal growth of tissues, wound healing, the female reproductive cycle (i.e., ovulation, menstruation and placental development), as well as a major role in many diseases. Particular interest has focused on cancer, since

tumors cannot grow beyond a few millimeters in size without developing a new blood supply. Angiogenesis is also necessary for the spread and growth of tumor cell metastases.

One of the most important growth and survival factors for endothelium is vascular endothelial growth factor (VEGF). VEGF induces angiogenesis and endothelial cell proliferation and plays an important role in regulating vasculogenesis. VEGF is a heparin-binding glycoprotein that is secreted as a homodimer of 45 kDa. Most types of cells, but usually not endothelial cells themselves, secrete VEGF. Since the initially discovered VEGF, VEGF-A, increases vascular permeability, it was known as vascular permeability factor. In addition, VEGF causes vasodilatation, partly through stimulation of nitric oxide synthase in endothelial cells. VEGF can also stimulate cell migration and inhibit apoptosis.

There are several splice variants of VEGF-A. The major ones include: 121, 165, 189 and 206 amino acids (aa), each one comprising a specific exon addition. VEGF165 is the most predominant protein, but transcripts of VEGF 121 may be more abundant. VEGF206 is rarely expressed and has been detected only in fetal liver. Recently, other splice variants of 145 and 183 aa have also been described. The 165, 189 and 206 aa splice variants have heparin-binding domains, which help anchor them in extracellular matrix and are involved in binding to heparin sulfate and presentation to VEGF receptors. Such presentation is a key factor for VEGF potency (i.e., the heparin-binding forms are more active). Several other members of the VEGF family have been cloned including VEGF-B, -C, and -D. Placenta growth factor (PlGF) is also closely related to VEGF-A. VEGF-A, -B, -C, -D, and PlGF are all distantly related to platelet-derived growth factors-A and -B. Less is known about the function and regulation of VEGF-B, -C, and -D, but they do not seem to be regulated by the major pathways that regulate VEGF-A.

VEGF-A transcription is potentiated in response to hypoxia and by activated oncogenes. The transcription factors, hypoxia inducible factor-1a (hif-1a) and -2a, are degraded by proteosomes in normoxia and stabilized in hypoxia. This pathway is dependent on the Von Hippel-Lindau gene product. Hif-1a and hif-2 heterodimerize with the aryl hydrocarbon nuclear translocator in the nucleus and bind the VEGF promoter/enhancer. This is a key pathway expressed in most types of cells. Hypoxia inducibility, in particular,

characterizes VEGF-A versus other members of the VEGF family and other angiogenic factors. VEGF transcription in normoxia is activated by many oncogenes, including H-ras and several transmembrane tyrosine kinases, such as the epidermal growth factor receptor and erbB2. These pathways together account for a marked upregulation of VEGF-A in  
5 tumors compared to normal tissues and are often of prognostic importance.

There are three receptors in the VEGF receptor family. They have the common properties of multiple IgG-like extracellular domains and tyrosine kinase activity. The enzyme domains of VEGF receptor 1 (VEGFr1, also known as Flt-1), VEGFr2 (also known as KDR or Flk-1), and VEGFr3 (also known as Flt-4) are divided by an inserted sequence.  
10 Endothelial cells also express additional VEGF receptors, Neuropilin-1 and Neuropilin-2. VEGF-A binds to VEGFr1 and VEGFr2 and to Neuropilin-1 and Neuropilin-2. PlGF and VEGF-B bind VEGFr1 and Neuropilin-1. VEGF-C and -D bind VEGFr3 and VEGFr2.

The VEGF-C/VEGFr3 pathway is important for lymphatic proliferation. VEGFr3 is specifically expressed on lymphatic endothelium. A soluble form of Flt-1 can be detected in  
15 peripheral blood and is a high affinity ligand for VEGF. Soluble Flt-1 can be used to antagonize VEGF function. VEGFr1 and VEGFr2 are upregulated in tumor and proliferating endothelium, partly by hypoxia and also in response to VEGF-A itself. VEGFr1 and VEGFr2 can interact with multiple downstream signaling pathways via proteins such as PLC-g, Ras, Shc, Nck, PKC and PI3-kinase. VEGFr1 is of higher affinity than VEGFr2 and  
20 mediates motility and vascular permeability. VEGFr2 is necessary for proliferation.

VEGF can be detected in both plasma and serum samples of patients, with much higher levels in serum. Platelets release VEGF upon aggregation and may be a major source of VEGF delivery to tumors. Several studies have shown that association of high serum levels of VEGF with poor prognosis in cancer patients may be correlated with an elevated  
25 platelet count. Many tumors release cytokines that can stimulate the production of megakaryocytes in the marrow and elevate the platelet count. This can result in an indirect increase of VEGF delivery to tumors.

VEGF is implicated in several other pathological conditions associated with enhanced angiogenesis. For example, VEGF plays a role in both psoriasis and rheumatoid arthritis. Diabetic retinopathy is associated with high intraocular levels of VEGF. Inhibition of VEGF function may result in infertility by blockade of corpus luteum function. Direct  
5 demonstration of the importance of VEGF in tumor growth has been achieved using dominant negative VEGF receptors to block in vivo proliferation, as well as blocking antibodies to VEGF39 or to VEGFr2.

The use of small interfering nucleic acid molecules targeting VEGF and corresponding receptors and ligands therefore provides a class of novel therapeutic agents that can be used  
10 in the diagnosis of and the treatment of cancer, proliferative diseases, or any other disease or condition that responds to modulation of VEGF and/or VEGFr genes.

Examples:

The following are non-limiting examples showing the selection, isolation, synthesis and activity of nucleic acids of the instant invention.

15 Example 1: Tandem synthesis of siNA constructs

Exemplary siNA molecules of the invention are synthesized in tandem using a cleavable linker, for example, a succinyl-based linker. Tandem synthesis as described herein is followed by a one-step purification process that provides RNAi molecules in high yield. This approach is highly amenable to siNA synthesis in support of high throughput  
20 RNAi screening, and can be readily adapted to multi-column or multi-well synthesis platforms.

After completing a tandem synthesis of a siNA oligo and its complement in which the 5'-terminal dimethoxytrityl (5'-O-DMT) group remains intact (trityl on synthesis), the oligonucleotides are deprotected as described above. Following deprotection, the siNA  
25 sequence strands are allowed to spontaneously hybridize. This hybridization yields a duplex in which one strand has retained the 5'-O-DMT group while the complementary strand comprises a terminal 5'-hydroxyl. The newly formed duplex behaves as a single molecule



during routine solid-phase extraction purification (Trityl-On purification) even though only one molecule has a dimethoxytrityl group. Because the strands form a stable duplex, this dimethoxytrityl group (or an equivalent group, such as other trityl groups or other hydrophobic moieties) is all that is required to purify the pair of oligos, for example, by  
5 using a C18 cartridge.

Standard phosphoramidite synthesis chemistry is used up to the point of introducing a tandem linker, such as an inverted deoxy abasic succinate or glyceryl succinate linker (see Figure 1) or an equivalent cleavable linker. A non-limiting example of linker coupling conditions that can be used includes a hindered base such as diisopropylethylamine (DIPA)  
10 and/or DMAP in the presence of an activator reagent such as Bromotripyrrolidinophosphoniumhexafluorophosphate (PyBrOP). After the linker is coupled, standard synthesis chemistry is utilized to complete synthesis of the second sequence leaving the terminal the 5'-O-DMT intact. Following synthesis, the resulting oligonucleotide is deprotected according to the procedures described herein and quenched  
15 with a suitable buffer, for example with 50mM NaOAc or 1.5M  $\text{NH}_4\text{H}_2\text{CO}_3$ .

Purification of the siNA duplex can be readily accomplished using solid phase extraction, for example using a Waters C18 SepPak 1g cartridge conditioned with 1 column volume (CV) of acetonitrile, 2 CV  $\text{H}_2\text{O}$ , and 2 CV 50mM NaOAc. The sample is loaded and then washed with 1 CV  $\text{H}_2\text{O}$  or 50mM NaOAc. Failure sequences are eluted with 1 CV  
20 14% ACN (Aqueous with 50mM NaOAc and 50mM NaCl). The column is then washed, for example with 1 CV  $\text{H}_2\text{O}$  followed by on-column detritylation, for example by passing 1 CV of 1% aqueous trifluoroacetic acid (TFA) over the column, then adding a second CV of 1% aqueous TFA to the column and allowing to stand for approximately 10 minutes. The remaining TFA solution is removed and the column washed with  $\text{H}_2\text{O}$  followed by 1 CV 1M  
25 NaCl and additional  $\text{H}_2\text{O}$ . The siNA duplex product is then eluted, for example, using 1 CV 20% aqueous CAN.

Figure 2 provides an example of MALDI-TOV mass spectrometry analysis of a purified siNA construct in which each peak corresponds to the calculated mass of an individual siNA strand of the siNA duplex. The same purified siNA provides three peaks

when analyzed by capillary gel electrophoresis (CGE), one peak presumably corresponding to the duplex siNA, and two peaks presumably corresponding to the separate siNA sequence strands. Ion exchange HPLC analysis of the same siNA construct only shows a single peak. Testing of the purified siNA construct using a luciferase reporter assay described below  
5 demonstrated the same RNAi activity compared to siNA constructs generated from separately synthesized oligonucleotide sequence strands.

Example 2: Identification of potential siNA target sites in any RNA sequence

The sequence of an RNA target of interest, such as a viral or human mRNA transcript, is screened for target sites, for example by using a computer folding algorithm. In a non-  
10 limiting example, the sequence of a gene or RNA gene transcript derived from a database, such as Genbank, is used to generate siNA targets having complementarity to the target. Such sequences can be obtained from a database, or can be determined experimentally as known in the art. Target sites that are known, for example, those target sites determined to be effective target sites based on studies with other nucleic acid molecules, for example  
15 ribozymes or antisense, or those targets known to be associated with a disease or condition such as those sites containing mutations or deletions, can be used to design siNA molecules targeting those sites. Various parameters can be used to determine which sites are the most suitable target sites within the target RNA sequence. These parameters include but are not limited to secondary or tertiary RNA structure, the nucleotide base composition of the target  
20 sequence, the degree of homology between various regions of the target sequence, or the relative position of the target sequence within the RNA transcript. Based on these determinations, any number of target sites within the RNA transcript can be chosen to screen siNA molecules for efficacy, for example by using *in vitro* RNA cleavage assays, cell culture, or animal models. In a non-limiting example, anywhere from 1 to 1000 target sites  
25 are chosen within the transcript based on the size of the siNA construct to be used. High throughput screening assays can be developed for screening siNA molecules using methods known in the art, such as with multi-well or multi-plate assays to determine efficient reduction in target gene expression.

Example 3: Selection of siNA molecule target sites in a RNA

The following non-limiting steps can be used to carry out the selection of siNAs targeting a given gene sequence or transcript.

1. The target sequence is parsed *in silico* into a list of all fragments or subsequences of a particular length, for example 23 nucleotide fragments, contained within the target sequence. This step is typically carried out using a custom Perl script, but commercial sequence analysis programs such as Oligo, MacVector, or the GCG Wisconsin Package can be employed as well.
2. In some instances the siNAs correspond to more than one target sequence; such would be the case for example in targeting different transcripts of the same gene, targeting different transcripts of more than one gene, or for targeting both the human gene and an animal homolog. In this case, a subsequence list of a particular length is generated for each of the targets, and then the lists are compared to find matching sequences in each list. The subsequences are then ranked according to the number of target sequences that contain the given subsequence; the goal is to find subsequences that are present in most or all of the target sequences. Alternately, the ranking can identify subsequences that are unique to a target sequence, such as a mutant target sequence. Such an approach would enable the use of siNA to target specifically the mutant sequence and not effect the expression of the normal sequence.
3. In some instances the siNA subsequences are absent in one or more sequences while present in the desired target sequence; such would be the case if the siNA targets a gene with a paralogous family member that is to remain untargeted. As in case 2 above, a subsequence list of a particular length is generated for each of the targets, and then the lists are compared to find sequences that are present in the target gene but are absent in the untargeted paralog.
4. The ranked siNA subsequences can be further analyzed and ranked according to GC content. A preference can be given to sites containing 30-70% GC, with a further preference to sites containing 40-60% GC.

5. The ranked siNA subsequences can be further analyzed and ranked according to self-folding and internal hairpins. Weaker internal folds are preferred; strong hairpin structures are to be avoided.
6. The ranked siNA subsequences can be further analyzed and ranked according to whether they have runs of GGG or CCC in the sequence. GGG (or even more Gs) in either strand can make oligonucleotide synthesis problematic and can potentially interfere with RNAi activity, so it is avoided whenever better sequences are available. CCC is searched in the target strand because that will place GGG in the antisense strand.
7. The ranked siNA subsequences can be further analyzed and ranked according to whether they have the dinucleotide UU (uridine dinucleotide) on the 3'-end of the sequence, and/or AA on the 5'-end of the sequence (to yield 3' UU on the antisense sequence). These sequences allow one to design siNA molecules with terminal TT thymidine dinucleotides.
8. Four or five target sites are chosen from the ranked list of subsequences as described above. For example, in subsequences having 23 nucleotides, the right 21 nucleotides of each chosen 23-mer subsequence are then designed and synthesized for the upper (sense) strand of the siNA duplex, while the reverse complement of the left 21 nucleotides of each chosen 23-mer subsequence are then designed and synthesized for the lower (antisense) strand of the siNA duplex (see Tables II and III). If terminal TT residues are desired for the sequence (as described in paragraph 7), then the two 3' terminal nucleotides of both the sense and antisense strands are replaced by TT prior to synthesizing the oligos.
9. The siNA molecules are screened in an *in vitro*, cell culture or animal model system to identify the most active siNA molecule or the most preferred target site within the target RNA sequence.

In an alternate approach, a pool of siNA constructs specific to a VEGF and/or VEGFr target sequence is used to screen for target sites in cells expressing VEGF and/or VEGFr RNA, such as HUVEC, HMVEC, or A375 cells. The general strategy used in this approach

is shown in Figure 9. A non-limiting example of such is a pool comprising sequences having any of SEQ ID NOS 1-2238. Cells expressing VEGF and/or VEGFr (e.g., HUVEC, HMVEC, or A375 cells) are transfected with the pool of siNA constructs and cells that demonstrate a phenotype associated with VEGF and/or VEGFr inhibition are sorted. The pool of siNA constructs can be expressed from transcription cassettes inserted into appropriate vectors (see for example Figure 7 and Figure 8). The siNA from cells demonstrating a positive phenotypic change (e.g., decreased proliferation, decreased VEGF and/or VEGFr mRNA levels or decreased VEGF and/or VEGFr protein expression), are sequenced to determine the most suitable target site(s) within the target VEGF and/or VEGFr RNA sequence.

Example 4: VEGF and/or VEGFr targeted siNA design

siNA target sites were chosen by analyzing sequences of the VEGF and/or VEGFr RNA target and optionally prioritizing the target sites on the basis of folding (structure of any given sequence analyzed to determine siNA accessibility to the target), by using a library of siNA molecules as described in Example 3, or alternately by using an *in vitro* siNA system as described in Example 6 herein. siNA molecules were designed that could bind each target and are optionally individually analyzed by computer folding to assess whether the siNA molecule can interact with the target sequence. Varying the length of the siNA molecules can be chosen to optimize activity. Generally, a sufficient number of complementary nucleotide bases are chosen to bind to, or otherwise interact with, the target RNA, but the degree of complementarity can be modulated to accommodate siNA duplexes or varying length or base composition. By using such methodologies, siNA molecules can be designed to target sites within any known RNA sequence, for example those RNA sequences corresponding to the any gene transcript.

Chemically modified siNA constructs are designed to provide nuclease stability for systemic administration *in vivo* and/or improved pharmacokinetic, localization, and delivery properties while preserving the ability to mediate RNAi activity. Chemical modifications as described herein are introduced synthetically using synthetic methods described herein and those generally known in the art. The synthetic siNA constructs are then assayed for

nuclease stability in serum and/or cellular/tissue extracts (e.g. liver extracts). The synthetic siNA constructs are also tested in parallel for RNAi activity using an appropriate assay, such as a luciferase reporter assay as described herein or another suitable assay that can quantify RNAi activity. Synthetic siNA constructs that possess both nuclease stability and RNAi activity can be further modified and re-evaluated in stability and activity assays. The chemical modifications of the stabilized active siNA constructs can then be applied to any siNA sequence targeting any chosen RNA and used, for example, in target screening assays to pick lead siNA compounds for therapeutic development (see for example Figure 11).

Example 5: Chemical Synthesis and Purification of siNA

siNA molecules can be designed to interact with various sites in the RNA message, for example, target sequences within the RNA sequences described herein. The sequence of one strand of the siNA molecule(s) is complementary to the target site sequences described above. The siNA molecules can be chemically synthesized using methods described herein. Inactive siNA molecules that are used as control sequences can be synthesized by scrambling the sequence of the siNA molecules such that it is not complementary to the target sequence. Generally, siNA constructs can be synthesized using solid phase oligonucleotide synthesis methods as described herein (see for example Usman *et al.*, US Patent Nos. 5,804,683; 5,831,071; 5,998,203; 6,117,657; 6,353,098; 6,362,323; 6,437,117; 6,469,158; Scaringe *et al.*, US Patent Nos. 6,111,086; 6,008,400; 6,111,086 all incorporated by reference herein in their entirety).

In a non-limiting example, RNA oligonucleotides are synthesized in a stepwise fashion using the phosphoramidite chemistry as is known in the art. Standard phosphoramidite chemistry involves the use of nucleosides comprising any of 5'-O-dimethoxytrityl, 2'-O-tert-butyldimethylsilyl, 3'-O-2-Cyanoethyl N,N-diisopropylphosphoroamidite groups, and exocyclic amine protecting groups (e.g. N6-benzoyl adenosine, N4 acetyl cytidine, and N2-isobutyryl guanosine). Alternately, 2'-O-Silyl Ethers can be used in conjunction with acid-labile 2'-O-orthoester protecting groups in the synthesis of RNA as described by Scaringe *supra*. Differing 2' chemistries can require different protecting groups, for example 2'-deoxy-2'-amino nucleosides can utilize N-phthaloyl

protection as described by Usman *et al.*, US Patent 5,631,360, incorporated by reference herein in its entirety).

During solid phase synthesis, each nucleotide is added sequentially (3'- to 5'- direction) to the solid support-bound oligonucleotide. The first nucleoside at the 3'-end of the chain is covalently attached to a solid support (e.g., controlled pore glass or polystyrene) using various linkers. The nucleotide precursor, a ribonucleoside phosphoramidite, and activator are combined resulting in the coupling of the second nucleoside phosphoramidite onto the 5'-end of the first nucleoside. The support is then washed and any unreacted 5'-hydroxyl groups are capped with a capping reagent such as acetic anhydride to yield inactive 5'-acetyl moieties. The trivalent phosphorus linkage is then oxidized to a more stable phosphate linkage. At the end of the nucleotide addition cycle, the 5'-O-protecting group is cleaved under suitable conditions (e.g., acidic conditions for trityl-based groups and Fluoride for silyl-based groups). The cycle is repeated for each subsequent nucleotide.

Modification of synthesis conditions can be used to optimize coupling efficiency, for example by using differing coupling times, differing reagent/phosphoramidite concentrations, differing contact times, differing solid supports and solid support linker chemistries depending on the particular chemical composition of the siNA to be synthesized. Deprotection and purification of the siNA can be performed as is generally described in Usman *et al.*, US 5,831,071, US 6,353,098, US 6,437,117, and Bellon *et al.*, US 6,054,576, US 6,162,909, US 6,303,773, or Scaringe *supra*, incorporated by reference herein in their entireties. Additionally, deprotection conditions can be modified to provide the best possible yield and purity of siNA constructs. For example, applicant has observed that oligonucleotides comprising 2'-deoxy-2'-fluoro nucleotides can degrade under inappropriate deprotection conditions. Such oligonucleotides are deprotected using aqueous methylamine at about 35°C for 30 minutes. If the 2'-deoxy-2'-fluoro containing oligonucleotide also comprises ribonucleotides, after deprotection with aqueous methylamine at about 35°C for 30 minutes, TEA-HF is added and the reaction maintained at about 65°C for an additional 15 minutes.

Example 6: RNAi *in vitro* assay to assess siNA activity

An *in vitro* assay that recapitulates RNAi in a cell-free system is used to evaluate siNA constructs targeting VEGF and/or VEGFr RNA targets. The assay comprises the system described by Tuschl *et al.*, 1999, *Genes and Development*, 13, 3191-3197 and Zamore *et al.*, 5 2000, *Cell*, 101, 25-33 adapted for use with VEGF and/or VEGFr target RNA. A *Drosophila* extract derived from syncytial blastoderm is used to reconstitute RNAi activity *in vitro*. Target RNA is generated via *in vitro* transcription from an appropriate VEGF and/or VEGFr expressing plasmid using T7 RNA polymerase or via chemical synthesis as described herein. Sense and antisense siNA strands (for example 20 uM each) are annealed 10 by incubation in buffer (such as 100 mM potassium acetate, 30 mM HEPES-KOH, pH 7.4, 2 mM magnesium acetate) for 1 min. at 90°C followed by 1 hour at 37°C, then diluted in lysis buffer (for example 100 mM potassium acetate, 30 mM HEPES-KOH at pH 7.4, 2mM magnesium acetate). Annealing can be monitored by gel electrophoresis on an agarose gel in TBE buffer and stained with ethidium bromide. The *Drosophila* lysate is prepared using 15 zero to two-hour-old embryos from Oregon R flies collected on yeasted molasses agar that are dechorionated and lysed. The lysate is centrifuged and the supernatant isolated. The assay comprises a reaction mixture containing 50% lysate [vol/vol], RNA (10-50 pM final concentration), and 10% [vol/vol] lysis buffer containing siNA (10 nM final concentration). The reaction mixture also contains 10 mM creatine phosphate, 10 ug/ml creatine 20 phosphokinase, 100 uM GTP, 100 uM UTP, 100 uM CTP, 500 uM ATP, 5 mM DTT, 0.1 U/uL RNasin (Promega), and 100 uM of each amino acid. The final concentration of potassium acetate is adjusted to 100 mM. The reactions are pre-assembled on ice and preincubated at 25° C for 10 minutes before adding RNA, then incubated at 25° C for an additional 60 minutes. Reactions are quenched with 4 volumes of 1.25 x Passive Lysis 25 Buffer (Promega). Target RNA cleavage is assayed by RT-PCR analysis or other methods known in the art and are compared to control reactions in which siNA is omitted from the reaction.

Alternately, internally-labeled target RNA for the assay is prepared by *in vitro* transcription in the presence of [ $\alpha$ -<sup>32</sup>P] CTP, passed over a G 50 Sephadex column by 30 spin chromatography and used as target RNA without further purification. Optionally,



target RNA is 5'-<sup>32</sup>P-end labeled using T4 polynucleotide kinase enzyme. Assays are performed as described above and target RNA and the specific RNA cleavage products generated by RNAi are visualized on an autoradiograph of a gel. The percentage of cleavage is determined by Phosphor Imager<sup>®</sup> quantitation of bands representing intact  
5 control RNA or RNA from control reactions without siNA and the cleavage products generated by the assay.

In one embodiment, this assay is used to determine target sites the VEGF and/or VEGFr RNA target for siNA mediated RNAi cleavage, wherein a plurality of siNA constructs are screened for RNAi mediated cleavage of the VEGF and/or VEGFr RNA  
10 target, for example, by analyzing the assay reaction by electrophoresis of labeled target RNA, or by northern blotting, as well as by other methodology well known in the art.

Example 7: Nucleic acid inhibition of VEGF and/or VEGFr target RNA *in vivo*

siNA molecules targeted to the huma VEGF and/or VEGFr RNA are designed and synthesized as described above. These nucleic acid molecules can be tested for cleavage  
15 activity *in vivo*, for example, using the following procedure. The target sequences and the nucleotide location within the VEGF and/or VEGFr RNA are given in **Table II and III**.

Two formats are used to test the efficacy of siNAs targeting VEGF and/or VEGFr. First, the reagents are tested in cell culture using, for example, HUVEC, HMVEC, or A375 cells to determine the extent of RNA and protein inhibition. siNA reagents (*e.g.*; see **Tables  
20 II and III**) are selected against the VEGF and/or VEGFr target as described herein. RNA inhibition is measured after delivery of these reagents by a suitable transfection agent to, for example, HUVEC, HMVEC, or A375 cells. Relative amounts of target RNA are measured versus actin using real-time PCR monitoring of amplification (*eg.*, ABI 7700 Taqman<sup>®</sup>). A comparison is made to a mixture of oligonucleotide sequences made to unrelated targets or  
25 to a randomized siNA control with the same overall length and chemistry, but randomly substituted at each position. Primary and secondary lead reagents are chosen for the target and optimization performed. After an optimal transfection agent concentration is chosen, a

RNA time-course of inhibition is performed with the lead siNA molecule. In addition, a cell-plating format can be used to determine RNA inhibition.

#### Delivery of siNA to Cells

Cells (e.g., HUVEC, HMVEC, or A375 cells) are seeded, for example, at  $1 \times 10^5$  cells per well of a six-well dish in EGM-2 (BioWhittaker) the day before transfection. siNA (final concentration, for example 20nM) and cationic lipid (e.g., final concentration  $2 \mu\text{g/ml}$ ) are complexed in EGM basal media (Biowhittaker) at  $37^\circ\text{C}$  for 30 mins in polystyrene tubes. Following vortexing, the complexed siNA is added to each well and incubated for the times indicated. For initial optimization experiments, cells are seeded, for example, at  $1 \times 10^3$  in 96 well plates and siNA complex added as described. Efficiency of delivery of siNA to cells is determined using a fluorescent siNA complexed with lipid. Cells in 6-well dishes are incubated with siNA for 24 hours, rinsed with PBS and fixed in 2% paraformaldehyde for 15 minutes at room temperature. Uptake of siNA is visualized using a fluorescent microscope.

#### 15 Taqman and Lightcycler quantification of mRNA

Total RNA is prepared from cells following siNA delivery, for example, using Qiagen RNA purification kits for 6-well or Rneasy extraction kits for 96-well assays. For Taqman analysis, dual-labeled probes are synthesized with the reporter dye, FAM or JOE, covalently linked at the 5'-end and the quencher dye TAMRA conjugated to the 3'-end. One-step RT-PCR amplifications are performed on, for example, an ABI PRISM 7700 Sequence Detector using 50  $\mu\text{l}$  reactions consisting of 10  $\mu\text{l}$  total RNA, 100 nM forward primer, 900 nM reverse primer, 100 nM probe, 1X TaqMan PCR reaction buffer (PE-Applied Biosystems), 5.5 mM  $\text{MgCl}_2$ , 300  $\mu\text{M}$  each dATP, dCTP, dGTP, and dTTP, 10U RNase Inhibitor (Promega), 1.25U AmpliTaq Gold (PE-Applied Biosystems) and 10U M-MLV Reverse Transcriptase (Promega). The thermal cycling conditions can consist of 30 min at  $48^\circ\text{C}$ , 10 min at  $95^\circ\text{C}$ , followed by 40 cycles of 15 sec at  $95^\circ\text{C}$  and 1 min at  $60^\circ\text{C}$ . Quantitation of mRNA levels is determined relative to standards generated from serially diluted total cellular RNA (300, 100, 33, 11 ng/txn) and normalizing to  $\beta$ -actin or GAPDH mRNA in

parallel TaqMan reactions. For each gene of interest an upper and lower primer and a fluorescently labeled probe are designed. Real time incorporation of SYBR Green I dye into a specific PCR product can be measured in glass capillary tubes using a lightcycler. A standard curve is generated for each primer pair using control cRNA. Values are represented as relative expression to GAPDH in each sample.

#### Western blotting

Nuclear extracts can be prepared using a standard micro preparation technique (see for example Andrews and Faller, 1991, *Nucleic Acids Research*, 19, 2499). Protein extracts from supernatants are prepared, for example using TCA precipitation. An equal volume of 20% TCA is added to the cell supernatant, incubated on ice for 1 hour and pelleted by centrifugation for 5 minutes. Pellets are washed in acetone, dried and resuspended in water. Cellular protein extracts are run on a 10% Bis-Tris NuPage (nuclear extracts) or 4-12% Tris-Glycine (supernatant extracts) polyacrylamide gel and transferred onto nitro-cellulose membranes. Non-specific binding can be blocked by incubation, for example, with 5% non-fat milk for 1 hour followed by primary antibody for 16 hour at 4°C. Following washes, the secondary antibody is applied, for example (1:10,000 dilution) for 1 hour at room temperature and the signal detected with SuperSignal reagent (Pierce).

#### Example 8: Animal Models useful to evaluate the down-regulation of VEGF and/or VEGFr gene expression

There are several animal models in which the anti-angiogenesis effect of nucleic acids of the present invention, such as siRNA, directed against VEGF, VEGFr1, VEGFr2 and/or VEGFr3 mRNAs can be tested. Typically a corneal model has been used to study angiogenesis in rat and rabbit since recruitment of vessels can easily be followed in this normally avascular tissue (Pandey *et al.*, 1995 *Science* 268: 567-569). In these models, a small Teflon or Hydron disk pretreated with an angiogenesis factor (e.g. bFGF or VEGF) is inserted into a pocket surgically created in the cornea. Angiogenesis is monitored 3 to 5 days later. siRNA directed against VEGF, VEGFr1, VEGFr2 and/or VEGFr3 mRNAs are delivered in the disk as well, or dropwise to the eye over the time course of the experiment.

In another eye model, hypoxia has been shown to cause both increased expression of VEGF and neovascularization in the retina (Pierce *et al.*, 1995 *Proc. Natl. Acad. Sci. USA.* 92: 905-909; Shweiki *et al.*, 1992 *J. Clin. Invest.* 91: 2235-2243).

5 In human glioblastomas, it has been shown that VEGF is at least partially responsible for tumor angiogenesis (Plate *et al.*, 1992 *Nature* 359, 845). Animal models have been developed in which glioblastoma cells are implanted subcutaneously into nude mice and the progress of tumor growth and angiogenesis is studied (Kim *et al.*, 1993 *supra*; Millauer *et al.*, 1994 *supra*).

10 Another animal model that addresses neovascularization involves Matrigel, an extract of basement membrane that becomes a solid gel when injected subcutaneously (Passaniti *et al.*, 1992 *Lab. Invest.* 67: 519-528). When the Matrigel is supplemented with angiogenesis factors such as VEGF, vessels grow into the Matrigel over a period of 3 to 5 days and angiogenesis can be assessed. Again, nucleic acids directed against VEGF mRNA are delivered in the Matrigel.

15 Several animal models exist for screening of anti-angiogenic agents. These include corneal vessel formation following corneal injury (Burger *et al.*, 1985 *Cornea* 4: 35-41; Lepri, *et al.*, 1994 *J. Ocular Pharmacol.* 10: 273-280; Ormerod *et al.*, 1990 *Am. J. Pathol.* 137: 1243-1252) or intracorneal growth factor implant (Grant *et al.*, 1993 *Diabetologia* 36: 282-291; Pandey *et al.* 1995 *supra*; Zieche *et al.*, 1992 *Lab. Invest.* 67: 711-715), vessel  
20 growth into Matrigel matrix containing growth factors (Passaniti *et al.*, 1992 *supra*), female reproductive organ neovascularization following hormonal manipulation (Shweiki *et al.*, 1993 *Clin. Invest.* 91: 2235-2243), several models involving inhibition of tumor growth in highly vascularized solid tumors (O'Reilly *et al.*, 1994 *Cell* 79: 315-328; Senger *et al.*, 1993 *Cancer and Metas. Rev.* 12: 303-324; Takahasi *et al.*, 1994 *Cancer Res.* 54: 4233-  
25 4237; Kim *et al.*, 1993 *supra*), and transient hypoxia-induced neovascularization in the mouse retina (Pierce *et al.*, 1995 *Proc. Natl. Acad. Sci. USA.* 92: 905-909).

The cornea model, described in Pandey *et al. supra*, is the most common and well characterized model for screening anti-angiogenic agent efficacy. This model involves an

5 avascular tissue into which vessels are recruited by a stimulating agent (growth factor, thermal or alkali burn, endotoxin). The corneal model utilizes the intrastromal corneal implantation of a Teflon pellet soaked in a VEGF-Hydron solution to recruit blood vessels toward the pellet, which can be quantitated using standard microscopic and image analysis techniques. To evaluate their anti-angiogenic efficacy, nucleic acids are applied topically to the eye or bound within Hydron on the Teflon pellet itself. This avascular cornea as well as the Matrigel (see below) provide for low background assays. While the corneal model has been performed extensively in the rabbit, studies in the rat have also been conducted.

The mouse model (Passaniti et al., *supra*) is a non-tissue model that utilizes Matrigel, an extract of basement membrane (Kleinman et al., 1986) or Millipore® filter disk, which can be impregnated with growth factors and anti-angiogenic agents in a liquid form prior to injection. Upon subcutaneous administration at body temperature, the Matrigel or Millipore® filter disk forms a solid implant. VEGF embedded in the Matrigel or Millipore® filter disk is used to recruit vessels within the matrix of the Matrigel or Millipore® filter disk which can be processed histologically for endothelial cell specific vWF (factor VIII antigen) immunohistochemistry, Trichrome-Masson stain, or hemoglobin content. Like the cornea, the Matrigel or Millipore® filter disk is avascular; however, it is not tissue. In the Matrigel or Millipore® filter disk model, nucleic acids are administered within the matrix of the Matrigel or Millipore® filter disk to test their anti-angiogenic efficacy. Thus, delivery issues in this model, as with delivery of nucleic acids by Hydron-coated Teflon pellets in the rat cornea model, may be less problematic due to the homogeneous presence of the nucleic acid within the respective matrix.

Other model systems to study tumor angiogenesis is reviewed by Folkman, 1985 *Adv. Cancer. Res.* 43, 175.

## 25 *Use of murine models*

For a typical systemic study involving 10 mice (20 g each) per dose group, 5 doses (1, 3, 10, 30 and 100 mg/kg daily over 14 days continuous administration), approximately 400

mg of siRNA, formulated in saline is used. A similar study in young adult rats (200 g) requires over 4 g. Parallel pharmacokinetic studies involve the use of similar quantities of siRNA further justifying the use of murine models.

*Lewis lung carcinoma and B-16 melanoma murine models*

- 5 Identifying a common animal model for systemic efficacy testing of nucleic acids is an efficient way of screening siRNA for systemic efficacy.

The Lewis lung carcinoma and B-16 murine melanoma models are well accepted models of primary and metastatic cancer and are used for initial screening of anti-cancer agents. These murine models are not dependent upon the use of immunodeficient mice, are  
10 relatively inexpensive, and minimize housing concerns. Both the Lewis lung and B-16 melanoma models involve subcutaneous implantation of approximately  $10^6$  tumor cells from metastatically aggressive tumor cell lines (Lewis lung lines 3LL or D122, LLc-LN7; B-16-BL6 melanoma) in C57BL/6J mice. Alternatively, the Lewis lung model can be produced by the surgical implantation of tumor spheres (approximately 0.8 mm in diameter).  
15 Metastasis also can be modeled by injecting the tumor cells directly intravenously. In the Lewis lung model, microscopic metastases can be observed approximately 14 days following implantation with quantifiable macroscopic metastatic tumors developing within 21-25 days. The B-16 melanoma exhibits a similar time course with tumor neovascularization beginning 4 days following implantation. Since both primary and  
20 metastatic tumors exist in these models after 21-25 days in the same animal, multiple measurements can be taken as indices of efficacy. Primary tumor volume and growth latency as well as the number of micro- and macroscopic metastatic lung foci or number of animals exhibiting metastases can be quantitated. The percent increase in lifespan can also be measured. Thus, these models provide suitable primary efficacy assays for screening  
25 systemically administered siRNA nucleic acids and siRNA nucleic acid formulations .

In the Lewis lung and B-16 melanoma models, systemic pharmacotherapy with a wide variety of agents usually begins 1-7 days following tumor implantation/inoculation with either continuous or multiple administration regimens. Concurrent pharmacokinetic studies

can be performed to determine whether sufficient tissue levels of siRNA can be achieved for pharmacodynamic effect to be expected. Furthermore, primary tumors and secondary lung metastases can be removed and subjected to a variety of *in vitro* studies (*i.e.* target RNA reduction).

5        In addition, animal models are useful in screening compounds, eg. siRNA molecules, for efficacy in treating renal failure, such as a result of autosomal dominant polycystic kidney disease (ADPKD). The Han:SPRD rat model, mice with a targeted mutation in the Pkd2 gene and congenital polycystic kidney (cpk) mice, closely resemble human ADPKD and provide animal models to evaluate the therapeutic effect of siRNA constructs that have  
10       the potential to interfere with one or more of the pathogenic elements of ADPKD mediated renal failure, such as angiogenesis. Angiogenesis may be necessary in the progression of ADPKD for growth of cyst cells as well as increased vascular permeability promoting fluid secretion into cysts. Proliferation of cystic epithelium is also a feature of ADPKD because cyst cells in culture produce soluble vascular endothelial growth factor (VEGF). VEGFr1  
15       has also been detected in epithelial cells of cystic tubules but not in endothelial cells in the vasculature of cystic kidneys or normal kidneys. VEGFr2 expression is increased in endothelial cells of cyst vessels and in endothelial cells during renal ischemia-reperfusion. It is proposed that inhibition of VEGF receptors with anti-VEGFr1 and anti-VEGFr2 siRNA molecules would attenuate cyst formation, renal failure and mortality in ADPKD. Anti-  
20       VEGFr2 siRNA molecules would therefore be designed to inhibit angiogenesis involved in cyst formation. As VEGFr1 is present in cystic epithelium and not in vascular endothelium of cysts, it is proposed that anti-VEGFr1 siRNA molecules would attenuate cystic epithelial cell proliferation and apoptosis which would in turn lead to less cyst formation. Further, it is proposed that VEGF produced by cystic epithelial cells is one of the stimuli for angiogenesis  
25       as well as epithelial cell proliferation and apoptosis. The use of Han:SPRD rats (see for example Kaspereit-Rittinghausen *et al.*, 1991, *Am.J.Pathol.* 139, 693-696), mice with a targeted mutation in the Pkd2 gene (Pkd2<sup>-/-</sup> mice, see for example Wu *et al.*, 2000, *Nat.Genet.* 24, 75-78) and cpk mice (see for example Woo *et al.*, 1994, *Nature*, 368, 750-753) all provide animal models to study the efficacy of siRNA molecules of the invention  
30       against VEGFr1 and VEGFr2 mediated renal failure.

VEGF, VEGFr1 VGFR2 and/or VEGFr3 protein levels can be measured clinically or experimentally by FACS analysis. VEGF, VEGFr1 VGFR2 and/or VEGFr3 encoded mRNA levels are assessed by Northern analysis, RNase-protection, primer extension analysis and/or quantitative RT-PCR. siRNA nucleic acids that block VEGF, VEGFr1  
5 VGFR2 and/or VEGFr3 protein encoding mRNAs and therefore result in decreased levels of VEGF, VEGFr1 VGFR2 and/or VEGFr3 activity by more than 20% *in vitro* can be identified.

Example 9: siNA-mediated inhibition of angiogenesis *in vivo*

The purpose of this study was to assess the anti-angiogenic activity of siNA targeted  
10 against VEGFr1 in the rat cornea model of VEGF induced angiogenesis (see above). The siNA molecules have matched inverted controls, which are inactive since they are not able to interact with the RNA target. The siNA molecules and VEGF were co-delivered using the filter disk method: Nitrocellulose filter disks (Millipore®) of 0.057 diameter were immersed in appropriate solutions and were surgically implanted in rat cornea as described  
15 by Pandey *et al.*, *supra*.

The stimulus for angiogenesis in this study was the treatment of the filter disk with 30  $\mu$ M VEGF, which is implanted within the cornea's stroma. This dose yields reproducible neovascularization stemming from the pericorneal vascular plexus growing toward the disk in a dose-response study 5 days following implant. Filter disks treated only with the vehicle  
20 for VEGF show no angiogenic response. The siNA were co-administered with VEGF on a disk in two different siNA concentrations. One concern with the simultaneous administration is that the siNA would not be able to inhibit angiogenesis since VEGF receptors could be stimulated. However, Applicant has observed that in low VEGF doses, the neovascular response reverts to normal, suggesting that the VEGF stimulus is essential  
25 for maintaining the angiogenic response. Blocking the production of VEGF receptors using simultaneous administration of anti-VEGF-R mRNA siNA could attenuate the normal neovascularization induced by the filter disk treated with VEGF.



## Materials and Methods:

### *Test Compounds and Controls*

R&D Systems VEGF, carrier free at 75  $\mu$ M in 82 mM Tris-Cl, pH 6.9

5 siNA, 1.67  $\mu$ G/ $\mu$ L, SITE 2340 (SEQ ID NO: 2; SEQ ID NO: 6) sense/antisense

siNA, 1.67  $\mu$ G/ $\mu$ L, INVERTED CONTROL FOR SITE 2340 (SEQ ID NO: 19; SEQ ID NO: 20) sense/antisense

siNA 1.67  $\mu$ g/ $\mu$ L, Site 2340 (SEQ ID NO: 419; SEQ ID NO: 420) sense/antisense

### *Animals*

10

Harlan Sprague-Dawley Rats, Approximately 225-250g

45 males, 5 animals per group.

### *Husbandry*

15

Animals are housed in groups of two. Feed, water, temperature and humidity are determined according to Pharmacology Testing Facility performance standards (SOP's) which are in accordance with the 1996 Guide for the Care and Use of Laboratory Animals (NRC). Animals are acclimated to the facility for at least 7 days prior to experimentation. During this time, animals are observed for overall health and sentinels are bled for baseline serology.

20

### *Experimental Groups*

Each solution (VEGF and siNAs) was prepared as a 1X solution for final concentrations shown in the experimental groups described in Table III.

25 *siNA Annealing Conditions*

siNA sense and antisense strands are annealed for 1 minute in H<sub>2</sub>O at 1.67mg/mL/strand followed by a 1 hour incubation at 37°C producing 3.34 mg/mL of duplexed siNA. For the 20µg/eye treatment, 6 µLs of the 3.34 mg/mL duplex is injected into the eye (see below). The 3.34 mg/mL duplex siNA can then be serially diluted for dose response assays.

#### *Preparation of VEGF Filter Disk*

For corneal implantation, 0.57 mm diameter nitrocellulose disks, prepared from 0.45 µm pore diameter nitrocellulose filter membranes (Millipore Corporation), were soaked for 30 min in 1 µL of 75 µM VEGF in 82 mM Tris·HCl (pH 6.9) in covered petri dishes on ice. Filter disks soaked only with the vehicle for VEGF (83 mM Tris-Cl pH 6.9) elicit no angiogenic response.

#### *Corneal surgery*

The rat corneal model used in this study was a modified from Koch *et al. Supra* and Pandey *et al., supra*. Briefly, corneas were irrigated with 0.5% povidone iodine solution followed by normal saline and two drops of 2% lidocaine. Under a dissecting microscope (Leica MZ-6), a stromal pocket was created and a presoaked filter disk (see above) was inserted into the pocket such that its edge was 1 mm from the corneal limbus.

#### *Intraconjunctival injection of test solutions*

Immediately after disk insertion, the tip of a 40-50 µm OD injector (constructed in our laboratory) was inserted within the conjunctival tissue 1 mm away from the edge of the corneal limbus that was directly adjacent to the VEGF-soaked filter disk. Six hundred nanoliters of test solution (siNA, inverted control or sterile water vehicle) were dispensed at a rate of 1.2 µL/min using a syringe pump (Kd Scientific). The injector was then removed, serially rinsed in 70% ethanol and sterile water and immersed in sterile water between each injection. Once the test solution was injected, closure of the eyelid was maintained using

microaneurism clips until the animal began to recover gross motor activity. Following treatment, animals were warmed on a heating pad at 37°C.

#### *Quantitation of angiogenic response*

5        Five days after disk implantation, animals were euthanized following administration of 0.4 mg/kg atropine and corneas were digitally imaged. The neovascular surface area (NSA, expressed in pixels) was measured *postmortem* from blood-filled corneal vessels using computerized morphometry (Image Pro Plus, Media Cybernetics, v2.0). The individual mean NSA was determined in triplicate from three regions of identical size in the area of  
10    maximal neovascularization between the filter disk and the limbus. The number of pixels corresponding to the blood-filled corneal vessels in these regions was summated to produce an index of NSA. A group mean NSA was then calculated. Data from each treatment group were normalized to VEGF/siNA vehicle-treated control NSA and finally expressed as percent inhibition of VEGF-induced angiogenesis.

#### 15    *Statistics*

After determining the normality of treatment group means, group mean percent inhibition of VEGF-induced angiogenesis was subjected to a one-way analysis of variance. This was followed by two post-hoc tests for significance including Dunnett's (comparison to  
20    VEGF control) and Tukey-Kramer (all other group mean comparisons) at alpha = 0.05. Statistical analyses were performed using JMP v.3.1.6 (SAS Institute).

Results are graphically represented in Figure 12. As shown in Figure 12, VEGFr1 site 4229 active siNA (RPI 29695/29699) at three concentrations were effective at inhibiting angiogenesis compared to the inverted siNA control (RPI 2983/29984) and the VEGF  
25    control. A chemically modified version of the VEGFr1 site 4229 active siNA comprising a sense strand having 2'-deoxy-2'-fluoro pyrimidines and ribo purines with 5' and 3' terminal inverted deoxybasic residues (RPI 30196) and an antisense strand having having 2'-deoxy-2'-fluoro pyrimidines and ribo purines with a terminal 3'-phosphorothioate internucleotide linkage (RPI 30416), showed similar inhibition. (Data not shown) This result shows siNA

molecules of differing chemically modified composition of the invention are capable of significantly inhibiting angiogenesis *in vivo*.

**Example 10: RNAi mediated inhibition of VEGF and/or VEGFr RNA expression**

siNA constructs (Table III) are tested for efficacy in reducing VEGF and/or VEGFr  
5 RNA expression in, for example, HUVEC, HMVEC, or A375 cells. Cells are plated  
approximately 24h before transfection in 96-well plates at 5,000-7,500 cells/well, 100  
μl/well, such that at the time of transfection cells are 70-90% confluent. For transfection,  
annealed siNAs are mixed with the transfection reagent (Lipofectamine 2000, Invitrogen) in  
a volume of 50 μl/well and incubated for 20 min. at room temperature. The siNA  
10 transfection mixtures are added to cells to give a final siNA concentration of 25 nM in a  
volume of 150 μl. Each siNA transfection mixture is added to 3 wells for triplicate siNA  
treatments. Cells are incubated at 37° for 24h in the continued presence of the siNA  
transfection mixture. At 24h, RNA is prepared from each well of treated cells. The  
supernatants with the transfection mixtures are first removed and discarded, then the cells  
15 are lysed and RNA prepared from each well. Target gene expression following treatment is  
evaluated by RT-PCR for the target gene and for a control gene (36B4, an RNA polymerase  
subunit) for normalization. The triplicate data is averaged and the standard deviations  
determined for each treatment. Normalized data are graphed and the percent reduction of  
target mRNA by active siNAs in comparison to their respective inverted control siNAs is  
20 determined.

Figure 13 shows a non-limiting example of reduction of VEGFr1 mRNA in A375  
cells mediated by chemically-modified siNAs that target VEGFr1 mRNA. A549 cells were  
transfected with 0.25 ug/well of lipid complexed with 25 nM siNA. A screen of siNA  
constructs (Stabilization "Stab" chemistries are shown in Table IV, constructs are referred  
25 to by RPI number, see Table III) comprising Stab 4/5 chemistry (RPI 31190/31193), Stab  
1/2 chemistry (RPI 31183/31186 and RPI 31184/31187), and unmodified RNA (RPI  
30075/30076) were compared to untreated cells, matched chemistry inverted control siNA  
constructs (RPI 31208/31211, RPI 31201/31204, RPI 31202/31205, and RPI 30077/30078),  
scrambled siNA control constructs (Scram1 and Scram2), and cells transfected with lipid

alone (transfection control). As shown in the figure, all of the siNA constructs significantly reduce VEGFr1 RNA expression. Additional stabilization chemistries as described in Table IV are similarly assayed for activity. These siNA constructs are compared to appropriate matched chemistry inverted controls. In addition, the siNA constructs are also compared to untreated cells, cells transfected with lipid and scrambled siNA constructs, and cells transfected with lipid alone (transfection control).

#### Example 11: Indications

The present body of knowledge in VEGF and/or VEGFr research indicates the need for methods to assay VEGF and/or VEGFr activity and for compounds that can regulate VEGF and/or VEGFr expression for research, diagnostic, and therapeutic use. As described herein, the nucleic acid molecules of the present invention can be used in assays to diagnose disease state related of VEGF and/or VEGFr levels. In addition, the nucleic acid molecules can be used to treat disease state related to VEGF and/or VEGFr levels.

Particular conditions and disease states that can be associated with VEGF and/or VEGFr expression modulation include, but are not limited to:

1) Tumor angiogenesis: Angiogenesis has been shown to be necessary for tumors to grow into pathological size (Folkman, 1971, *PNAS* 76, 5217-5221; Wellstein & Czubayko, 1996, *Breast Cancer Res and Treatment* 38, 109-119). In addition, it allows tumor cells to travel through the circulatory system during metastasis. Increased levels of gene expression of a number of angiogenic factors such as vascular endothelial growth factor (VEGF) have been reported in vascularized and edema-associated brain tumors (Berkman *et al.*, 1993 *J. Clin. Invest.* 91, 153). A more direct demonstration of the role of VEGF in tumor angiogenesis was demonstrated by Jim Kim *et al.*, 1993 *Nature* 362,841 wherein, monoclonal antibodies against VEGF were successfully used to inhibit the growth of rhabdomyosarcoma, glioblastoma multiforme cells in nude mice. Similarly, expression of a dominant negative mutated form of the flt-1 VEGF receptor inhibits vascularization induced by human glioblastoma cells in nude mice (Millauer *et al.*, 1994, *Nature* 367, 576). Specific

tumor/cancer types that can be targeted using the nucleic acid molecules of the invention include but are not limited to the tumor/cancer types described herein.

2) Ocular diseases: Neovascularization has been shown to cause or exacerbate ocular diseases including, but not limited to, macular degeneration, neovascular glaucoma, diabetic  
5 retinopathy, myopic degeneration, and trachoma (Norrby, 1997, *APMIS* 105, 417-437). Aiello *et al.*, 1994 *New Engl. J. Med.* 331, 1480, showed that the ocular fluid of a majority of patients suffering from diabetic retinopathy and other retinal disorders contains a high concentration of VEGF. Miller *et al.*, 1994 *Am. J. Pathol.* 145, 574, reported elevated levels of VEGF mRNA in patients suffering from retinal ischemia. These observations support a  
10 direct role for VEGF in ocular diseases. Other factors, including those that stimulate VEGF synthesis, may also contribute to these indications.

3) Dermatological Disorders: Many indications have been identified which may be angiogenesis dependent, including but not limited to, psoriasis, verruca vulgaris, angiofibroma of tuberous sclerosis, port-wine stains, Sturge Weber syndrome, Kippel-  
15 Trenaunay-Weber syndrome, and Osler-Weber-Rendu syndrome (Norrby, *supra*). Intradermal injection of the angiogenic factor b-FGF demonstrated angiogenesis in nude mice (Weckbecker *et al.*, 1992, *Angiogenesis: Key principles-Science-Technology-Medicine*, ed R. Steiner). Detmar *et al.*, 1994 *J. Exp. Med.* 180, 1141 reported that VEGF and its receptors were over-expressed in psoriatic skin and psoriatic dermal microvessels,  
20 suggesting that VEGF plays a significant role in psoriasis.

4) Rheumatoid arthritis: Immunohistochemistry and *in situ* hybridization studies on tissues from the joints of patients suffering from rheumatoid arthritis show an increased level of VEGF and its receptors (Fava *et al.*, 1994 *J. Exp. Med.* 180, 341). Additionally, Koch *et al.*, 1994 *J. Immunol.* 152, 4149, found that VEGF-specific antibodies were able to  
25 significantly reduce the mitogenic activity of synovial tissues from patients suffering from rheumatoid arthritis. These observations support a direct role for VEGF in rheumatoid arthritis. Other angiogenic factors including those of the present invention may also be involved in arthritis.

5) Endometriosis: Various studies indicate that VEGF is directly implicated in endometriosis. In one study, VEGF concentrations measured by ELISA in peritoneal fluid were found to be significantly higher in women with endometriosis than in women without endometriosis ( $24.1 \pm 15$  ng/ml vs  $13.3 \pm 7.2$  ng/ml in normals). In patients with endometriosis, higher concentrations of VEGF were detected in the proliferative phase of the menstrual cycle ( $33 \pm 13$  ng/ml) compared to the secretory phase ( $10.7 \pm 5$  ng/ml). The cyclic variation was not noted in fluid from normal patients (McLaren *et al.*, 1996, *Human Reprod.* 11, 220-223). In another study, women with moderate to severe endometriosis had significantly higher concentrations of peritoneal fluid VEGF than women without endometriosis. There was a positive correlation between the severity of endometriosis and the concentration of VEGF in peritoneal fluid. In human endometrial biopsies, VEGF expression increased relative to the early proliferative phase approximately 1.6-, 2-, and 3.6-fold in midproliferative, late proliferative, and secretory endometrium (Shifren *et al.*, 1996, *J. Clin. Endocrinol. Metab.* 81, 3112-3118). In a third study, VEGF-positive staining of human ectopic endometrium was shown to be localized to macrophages (double immunofluorescent staining with CD14 marker). Peritoneal fluid macrophages demonstrated VEGF staining in women with and without endometriosis. However, increased activation of macrophages (acid phosphatase activity) was demonstrated in fluid from women with endometriosis compared with controls. Peritoneal fluid macrophage conditioned media from patients with endometriosis resulted in significantly increased cell proliferation ( $[^3\text{H}]$  thymidine incorporation) in HUVEC cells compared to controls. The percentage of peritoneal fluid macrophages with VEGFr2 mRNA was higher during the secretory phase, and significantly higher in fluid from women with endometriosis ( $80 \pm 15\%$ ) compared with controls ( $32 \pm 20\%$ ). Flt-mRNA was detected in peritoneal fluid macrophages from women with and without endometriosis, but there was no difference between the groups or any evidence of cyclic dependence (McLaren *et al.*, 1996, *J. Clin. Invest.* 98, 482-489). In the early proliferative phase of the menstrual cycle, VEGF has been found to be expressed in secretory columnar epithelium (estrogen-responsive) lining both the oviducts and the uterus in female mice. During the secretory phase, VEGF expression was shown to have shifted to the underlying stroma composing the functional endometrium. In addition to examining the endometrium, neovascularization of ovarian

follicles and the corpus luteum, as well as angiogenesis in embryonic implantation sites have been analyzed. For these processes, VEGF was expressed in spatial and temporal proximity to forming vasculature (Shweiki *et al.*, 1993, *J. Clin. Invest.* 91, 2235-2243).

6) Kidney disease: Autosomal dominant polycystic kidney disease (ADPKD) is the most common life threatening hereditary disease in the USA. It affects about 1:400 to 1:1000 people and approximately 50% of people with ADPKD develop renal failure. ADPKD accounts for about 5-10% of end-stage renal failure in the USA, requiring dialysis and renal transplantation. Angiogenesis is implicated in the progression of ADPKD for growth of cyst cells, as well as increased vascular permeability promoting fluid secretion into cysts. Proliferation of cystic epithelium is a feature of ADPKD because cyst cells in culture produce soluble vascular endothelial growth factor (VEGF). VEGFr1 has been detected in epithelial cells of cystic tubules but not in endothelial cells in the vasculature of cystic kidneys or normal kidneys. VEGFr2 expression is increased in endothelial cells of cyst vessels and in endothelial cells during renal ischemia-reperfusion.

The use of radiation treatments and chemotherapeutics, such as Gemcytabine and cyclophosphamide, are non-limiting examples of chemotherapeutic agents that can be combined with or used in conjunction with the nucleic acid molecules (*e.g.* siNA molecules) of the instant invention. Those skilled in the art will recognize that other anti-cancer compounds and therapies can similarly be readily combined with the nucleic acid molecules of the instant invention (*e.g.* siNA molecules) and are hence within the scope of the instant invention. Such compounds and therapies are well known in the art (see for example *Cancer: Principles and Practice of Oncology*, Volumes 1 and 2, eds Devita, V.T., Hellman, S., and Rosenberg, S.A., J.B. Lippincott Company, Philadelphia, USA; incorporated herein by reference) and include, without limitation, folates, antifolates, pyrimidine analogs, fluoropyrimidines, purine analogs, adenosine analogs, topoisomerase I inhibitors, anthrapyrazoles, retinoids, antibiotics, anthacyclins, platinum analogs, alkylating agents, nitrosoureas, plant derived compounds such as vinca alkaloids, epipodophyllotoxins, tyrosine kinase inhibitors, taxols, radiation therapy, surgery, nutritional supplements, gene therapy, radiotherapy, for example 3D-CRT, immunotoxin therapy, for example ricin, and



monoclonal antibodies. Specific examples of chemotherapeutic compounds that can be combined with or used in conjunction with the nucleic acid molecules of the invention include, but are not limited to, Paclitaxel; Docetaxel; Methotrexate; Doxorubin; Edatrexate; Vinorelbine; Tomaxifen; Leucovorin; 5-fluoro uridine (5-FU); Ionotecan; Cisplatin; Carboplatin; Amsacrine; Cytarabine; Bleomycin; Mitomycin C; Dactinomycin; Mithramycin; Hexamethylmelamine; Dacarbazine; L-asparaginase; Nitrogen mustard; Melphalan, Chlorambucil; Busulfan; Ifosfamide; 4-hydroperoxycyclophosphamide; Thiotepe; Irinotecan (CAMPTOSAR®, CPT-11, Camptothecin-11, Campto) Tamoxifen; Herceptin; IMC C225; ABX-EGF; and combinations thereof. The above list of compounds are non-limiting examples of compounds and/or methods that can be combined with or used in conjunction with the nucleic acid molecules (e.g. siNA) of the instant invention. Those skilled in the art will recognize that other drug compounds and therapies can similarly be readily combined with the nucleic acid molecules of the instant invention (e.g., siNA molecules) are hence within the scope of the instant invention.

Example 12: Diagnostic uses

The siNA molecules of the invention can be used in a variety of diagnostic applications, such as in the identification of molecular targets (e.g., RNA) in a variety of applications, for example, in clinical, industrial, environmental, agricultural and/or research settings. Such diagnostic use of siNA molecules involves utilizing reconstituted RNAi systems, for example, using cellular lysates or partially purified cellular lysates. siNA molecules of this invention can be used as diagnostic tools to examine genetic drift and mutations within diseased cells or to detect the presence of endogenous or exogenous, for example viral, RNA in a cell. The close relationship between siNA activity and the structure of the target RNA allows the detection of mutations in any region of the molecule, which alters the base-pairing and three-dimensional structure of the target RNA. By using multiple siNA molecules described in this invention, one can map nucleotide changes, which are important to RNA structure and function *in vitro*, as well as in cells and tissues. Cleavage of target RNAs with siNA molecules can be used to inhibit gene expression and define the role of specified gene products in the progression of disease or infection. In this manner, other genetic targets can be defined as important mediators of the disease. These experiments will

lead to better treatment of the disease progression by affording the possibility of combination therapies (e.g., multiple siNA molecules targeted to different genes, siNA molecules coupled with known small molecule inhibitors, or intermittent treatment with combinations siNA molecules and/or other chemical or biological molecules). Other *in vitro* uses of siNA molecules of this invention are well known in the art, and include detection of the presence of mRNAs associated with a disease, infection, or related condition. Such RNA is detected by determining the presence of a cleavage product after treatment with a siNA using standard methodologies, for example, fluorescence resonance emission transfer (FRET).

10 In a specific example, siNA molecules that cleave only wild-type or mutant forms of the target RNA are used for the assay. The first siNA molecules (*i.e.*, those that cleave only wild-type forms of target RNA) are used to identify wild-type RNA present in the sample and the second siNA molecules (*i.e.*, those that cleave only mutant forms of target RNA) are used to identify mutant RNA in the sample. As reaction controls, synthetic substrates of both wild-type and mutant RNA are cleaved by both siNA molecules to demonstrate the relative siNA efficiencies in the reactions and the absence of cleavage of the "non-targeted" RNA species. The cleavage products from the synthetic substrates also serve to generate size markers for the analysis of wild-type and mutant RNAs in the sample population. Thus, each analysis requires two siNA molecules, two substrates and one unknown sample, which is combined into six reactions. The presence of cleavage products is determined using an RNase protection assay so that full-length and cleavage fragments of each RNA can be analyzed in one lane of a polyacrylamide gel. It is not absolutely required to quantify the results to gain insight into the expression of mutant RNAs and putative risk of the desired phenotypic changes in target cells. The expression of mRNA whose protein product is implicated in the development of the phenotype (*i.e.*, disease related or infection related) is adequate to establish risk. If probes of comparable specific activity are used for both transcripts, then a qualitative comparison of RNA levels is adequate and decreases the cost of the initial diagnosis. Higher mutant form to wild-type ratios are correlated with higher risk whether RNA levels are compared qualitatively or quantitatively.

All patents and publications mentioned in the specification are indicative of the levels of skill of those skilled in the art to which the invention pertains. All references cited in this disclosure are incorporated by reference to the same extent as if each reference had been incorporated by reference in its entirety individually.

5        One skilled in the art would readily appreciate that the present invention is well adapted to carry out the objects and obtain the ends and advantages mentioned, as well as those inherent therein. The methods and compositions described herein as presently representative of preferred embodiments are exemplary and are not intended as limitations on the scope of the invention. Changes therein and other uses will occur to those skilled in  
10    the art, which are encompassed within the spirit of the invention, are defined by the scope of the claims.

It will be readily apparent to one skilled in the art that varying substitutions and modifications can be made to the invention disclosed herein without departing from the scope and spirit of the invention. Thus, such additional embodiments are within the scope of  
15    the present invention and the following claims. The present invention teaches one skilled in the art to test various combinations and/or substitutions of chemical modifications described herein toward generating nucleic acid constructs with improved activity for mediating RNAi activity. Such improved activity can comprise improved stability, improved bioavailability, and/or improved activation of cellular responses mediating RNAi. Therefore, the specific  
20    embodiments described herein are not limiting and one skilled in the art can readily appreciate that specific combinations of the modifications described herein can be tested without undue experimentation toward identifying siNA molecules with improved RNAi activity.

The invention illustratively described herein suitably can be practiced in the absence  
25    of any element or elements, limitation or limitations that are not specifically disclosed herein. Thus, for example, in each instance herein any of the terms "comprising", "consisting essentially of", and "consisting of" may be replaced with either of the other two terms. The terms and expressions which have been employed are used as terms of description and not of limitation, and there is no intention that in the use of such terms and

expressions of excluding any equivalents of the features shown and described or portions thereof, but it is recognized that various modifications are possible within the scope of the invention claimed. Thus, it should be understood that although the present invention has been specifically disclosed by preferred embodiments, optional features, modification and  
5 variation of the concepts herein disclosed may be resorted to by those skilled in the art, and that such modifications and variations are considered to be within the scope of this invention as defined by the description and the appended claims.

In addition, where features or aspects of the invention are described in terms of Markush groups or other grouping of alternatives, those skilled in the art will recognize that  
10 the invention is also thereby described in terms of any individual member or subgroup of members of the Markush group or other group.

**Table I: VEGF and VEGFr Accession Numbers**

NM_005429	Homo sapiens vascular endothelial growth factor C (VEGFC), mRNA gi 19924300 ref NM_005429.2 [19924300]
NM_003376	Homo sapiens vascular endothelial growth factor (VEGF), mRNA gi 19923239 ref NM_003376.2 [19923239]
AF095785	Homo sapiens vascular endothelial growth factor (VEGF) gene, promoter region and partial cds gi 4154290 gb AF095785.1 [4154290]
NM_003377	Homo sapiens vascular endothelial growth factor B (VEGFB), mRNA gi 20070172 ref NM_003377.2 [20070172]
AF486837	Homo sapiens vascular endothelial growth factor isoform VEGF165 (VEGF) mRNA, complete cds gi 19909064 gb AF486837.1 [19909064]
AF468110	

Homo sapiens vascular endothelial growth factor B isoform (VEGFB) gene, complete cds, alternatively spliced  
gi|18766397|gb|AF468110.1|[18766397]

AF437895

Homo sapiens vascular endothelial growth factor (VEGF) gene, partial cds  
gi|16660685|gb|AF437895.1|AF437895[16660685]

AY047581

Homo sapiens vascular endothelial growth factor (VEGF) mRNA, complete cds  
gi|15422108|gb|AY047581.1|[15422108]

AF063657

Homo sapiens vascular endothelial growth factor receptor (FLT1) mRNA, complete cds  
gi|3132830|gb|AF063657.1|AF063657[3132830]

AF092127

Homo sapiens vascular endothelial growth factor (VEGF) gene, partial sequence  
gi|4139168|gb|AF092127.1|AF092127[4139168]

AF092126

Homo sapiens vascular endothelial growth factor (VEGF) gene, 5' UTR  
gi|4139167|gb|AF092126.1|AF092126[4139167]

AF092125  
Homo sapiens vascular endothelial growth factor (VEGF) gene, partial cds  
gi|4139165|gb|AF092125.1|AF092125[4139165]

E15157  
Human VEGF mRNA  
gi|5709840|dbj|E15157.1|pat|JP|1998052285|2[5709840]

E15156  
Human VEGF mRNA  
gi|5709839|dbj|E15156.1|pat|JP|1998052285|1[5709839]

E14233  
Human mRNA for vascular endothelial growth factor (VEGF), complete cds  
gi|5708916|dbj|E14233.1|pat|JP|1997286795|1[5708916]

AF024710  
Homo sapiens vascular endothelial growth factor (VEGF) mRNA, 3'UTR  
gi|2565322|gb|AF024710.1|AF024710[2565322]

AJ010438  
Homo sapiens mRNA for vascular endothelial growth factor, splicing variant  
VEGF183  
gi|3647280|emb|AJ010438.1|HSA010438[3647280]

AF098331  
Homo sapiens vascular endothelial growth factor (VEGF) gene, promoter, partial  
sequence  
gi|4235431|gb|AF098331.1|AF098331[4235431]

AF022375  
Homo sapiens vascular endothelial growth factor mRNA, complete cds  
gi|3719220|gb|AF022375.1|AF022375[3719220]

AH006909  
vascular endothelial growth factor {alternative splicing} [human, Genomic, 414  
nt 5 segments]  
gi|1680143|gb|AH006909.1||bbm|191843[1680143]

U01134  
Human soluble vascular endothelial cell growth factor receptor (sflt) mRNA,  
complete cds  
gi|451321|gb|U01134.1|U01134[451321]

E14000  
Human mRNA for FLT  
gi|3252767|dbj|E14000.1||pat|JP|1997255700|1[3252767]

E13332  
cDNA encoding vascular endothelial cell growth factor VEGF  
gi|3252137|dbj|E13332.1||pat|JP|1997173075|1[3252137]



E13256  
Human mRNA for FLT, complete cds  
gi|3252061|dbj|E13256.1||pat|JP|1997154588|1[3252061]

AF063658  
Homo sapiens vascular endothelial growth factor receptor 2 (KDR) mRNA, complete cds  
gi|3132832|gb|AF063658.1|AF063658[3132832]

AJ000185  
Homo Sapiens mRNA for vascular endothelial growth factor-D  
gi|2879833|emb|AJ000185.1|HSAJ185[2879833]

D89630  
Homo sapiens mRNA for VEGF-D, complete cds  
gi|2780339|dbj|D89630.1|[2780339]

AF035121  
Homo sapiens KDR/flk-1 protein mRNA, complete cds  
gi|2655411|gb|AF035121.1|AF035121[2655411]

AF020393  
Homo sapiens vascular endothelial growth factor C gene, partial cds and 5' upstream region  
gi|2582366|gb|AF020393.1|AF020393[2582366]

Y08736  
H.sapiens vegf gene, 3'UTR  
gi|1619596|emb|Y08736.1|HSVEGF3UT[1619596]

X62568  
H.sapiens vegf gene for vascular endothelial growth factor  
gi|37658|emb|X62568.1|HSVEGF[37658]

X94216  
H.sapiens mRNA for VEGF-C protein  
gi|1177488|emb|X94216.1|HSVEGFC[1177488]

NM\_002020  
Homo sapiens fms-related tyrosine kinase 4 (FLT4), mRNA  
gi|4503752|ref|NM\_002020.1|[4503752]

NM\_002253  
Homo sapiens kinase insert domain receptor (a type III receptor tyrosine kinase)  
(KDR), mRNA  
gi|11321596|ref|NM\_002253.1|[11321596]

Table II: VEGF and VEGFr siNA and Target Sequences

VEGFR1 gi4503748refNM_002019.1									
Pos	Target Sequence	Seq ID	UPos	Upper seq	Seq ID	LPos	Lower seq	Seq ID	
1	GCGGACACUCCUCCGCGC	1	1	GCGGACACUCCUCCGCGC	1	23	AGCCGAGAGGAGUGUCCGC	428	
18	UCCUCCCCGGCAGCGCGG	2	19	UCCUCCCCGGCAGCGCGG	2	41	CCGCCGUGCCGGGGAGGA	429	
37	GCGGCUCCGAGCGGGCUC	3	37	GCGGCUCCGAGCGGGCUC	3	59	GGAGCCCGUCCGAGCCGC	430	
55	CGGGGCUCCGGGUGCAGCGG	4	55	CGGGGCUCCGGGUGCAGCGG	4	77	CCGCUACCCGAGGCCCGC	431	
73	GCCAGCGGGCUGCGCGCG	5	73	GCCAGCGGGCUGCGCGCG	5	95	CGCCGCCAGGCCCGCUGGC	432	
91	GAGGAUUAACCGGGGAAGU	6	91	GAGGAUUAACCGGGGAAGU	6	113	ACUUCGCCGGGUAUCCUC	433	
109	UGGUUGUCCUCCGCGGGA	7	109	UGGUUGUCCUCCGCGGGA	7	131	UCCAGCCAGGAGACAACA	434	
127	AGCCGCGAGACGGCGCUC	8	127	AGCCGCGAGACGGCGCUC	8	149	GAGCGCCGUGUCGCGGCU	435	
145	CAGGGCGGGCGCGCGG	9	145	CAGGGCGGGCGCGCGG	9	167	CCGCCGGCCCGCGCCUG	436	
163	GCGGCGAACGAGGACGG	10	163	GCGGCGAACGAGGACGG	10	185	CCGUCUCUCUGUCCGCGC	437	
181	GACUCUGCGCGCGGGUCG	11	181	GACUCUGCGCGCGGGUCG	11	203	CGACCCGCGCCAGAGUC	438	
199	GUUGCGCGGGGAGCGCGG	12	199	GUUGCGCGGGGAGCGCGG	12	221	CGCGGUCGCCCGGCCAAC	439	
217	GGCACCGGGCGAGCAGGCC	13	217	GGCACCGGGCGAGCAGGCC	13	239	GGCCUGUCGCCCGGUGCC	440	
235	CGCGUCGCGCUCACCAUGG	14	235	CGCGUCGCGCUCACCAUGG	14	257	CCAUGGUGAGCGCGACGCG	441	
253	GUCAGCUACUGGACACCG	15	253	GUCAGCUACUGGACACCG	15	275	CGGUGUCCAGUAGCUGAC	442	
271	GGGGUCCUGUGGCGGCGC	16	271	GGGGUCCUGUGGCGGCGC	16	293	GGCGCACAGCAGGACCCC	443	
289	CUGCUCAGCUGUGCUCUUC	17	289	CUGCUCAGCUGUGCUCUUC	17	311	GAAGCAGACAGCUGAGCAG	444	
307	CUCACAGGAUCUAGUUCAG	18	307	CUCACAGGAUCUAGUUCAG	18	329	CUGAACUAGAUCCUGUGAG	445	
325	GGUUCAAAUAUAAAGAU	19	325	GGUUCAAAUAUAAAGAU	19	347	GAUCUUUAUUUUGAACCC	446	
343	CCUGAACUGAGUUUAAAG	20	343	CCUGAACUGAGUUUAAAG	20	365	CUUUUAAACUCAGUUCAGG	447	
361	GGCACCAGCACAUAUGC	21	361	GGCACCAGCACAUAUGC	21	383	GCAUGAUGUGUGGGUGGCC	448	
379	CAAGCAGGCCAGACACUJC	22	379	CAAGCAGGCCAGACACUJC	22	401	GCAGUGUCUGGCCUGCUUG	449	
397	CAUCUCCAUGCAGGGGGG	23	397	CAUCUCCAUGCAGGGGGG	23	419	CCCCCUGCAUUGGAGUAG	450	
415	GAAGCAGCCCAUAAUUGU	24	415	GAAGCAGCCCAUAAUUGU	24	437	ACCAUUUUGGGCUGCUUC	451	
433	UCUUUGCCUGAAUUGGUGA	25	433	UCUUUGCCUGAAUUGGUGA	25	455	UCACCAUUUCAGGCAAGA	452	
451	AGUAAGGAAAGCGAAAGGC	26	451	AGUAAGGAAAGCGAAAGGC	26	473	GCCUUUCGCUUCCUUAU	453	
469	CUGAGCAUAAUAAUUCUG	27	469	CUGAGCAUAAUAAUUCUG	27	491	CAGAUUUAUUAUUGCUCAG	454	
487	GCCUGUGGAAGAAUUGGCA	28	487	GCCUGUGGAAGAAUUGGCA	28	509	UGCCAUUUUCCUCCACAGGC	455	
505	AAACAUAUUCGACUACUUC	29	505	AAACAUAUUCGACUACUUC	29	527	AAGUACUGCAGAAUUGUU	456	
523	UUAAACCUUGAACACAGCUC	30	523	UUAAACCUUGAACACAGCUC	30	545	GAGCUGUGUUAACAGGUUAA	457	

541	CAAGCAAAACCCACACUGGCU	31	541	CAAGCAAAACCCACACUGGCU	31	563	AGCCAGUGUGGUUUUGCUUG	458
559	UUCUACAGCUGCAAAUAUC	32	559	CUAGCAGCUGCAAAUAUC	32	581	GAUAUUUGCAGCUGUAGAA	459
577	CUAGCUGUACCUACUCAA	33	577	CUAGCUGUACCUACUCAA	33	599	UUGAAGUAGGUACACUAG	460
595	AAGAAGAGGAAACAGAAU	34	595	AAGAAGAGGAAACAGAAU	34	617	AUUCUGUUUCCUUCUUCU	461
613	UCUGCAUCUAUAUAUUA	35	613	UCUGCAUCUAUAUAUUA	35	635	UAAUAUAUAGAUUGCAGA	462
631	AUUGAGUAGAGGUAGAC	36	631	AUUGAGUAGAGGUAGAC	36	653	GUUACCUUGUAUCACUAAU	463
649	CCUUUCGUAAGAGUACA	37	649	CCUUUCGUAAGAGUACA	37	671	UGUACAUCUCUACGAAAGG	464
667	AGUGAAUCCCGAAUUA	38	667	AGUGAAUCCCGAAUUA	38	689	UAUUUCGGGGAUUUUCACU	465
685	AGUGAAUCCCGAAUUA	39	685	AUACAUAGACUGAAGGAA	39	707	UUCUUUCAGUACUGUGUAU	466
703	AGGAGCUGUACUAGGAA	40	703	AGGAGCUGUACUAGGAA	40	725	AGGGAUAGCAGAGCUCUCCU	467
721	UGCCGGUACGUCACCUA	41	721	UGCCGGUACGUCACCUA	41	743	UAGGAGCAGUACCCGGCA	468
739	AACAUCACUUAUUA	42	739	AACAUCACUUAUUA	42	761	UUAAAGUAACAGUAGUUU	469
757	AAAAAGUUCCACUUGACA	43	757	AAAAAGUUCCACUUGACA	43	779	UUAAGUAGGAAACUUUUU	470
775	ACUUUGAUCCUUGGAA	44	775	ACUUUGAUCCUUGGAA	44	797	UUCCAUCAGGGAUCAAAGU	471
793	AAACGCAUAUCUGGACA	45	793	AAACGCAUAUCUGGACA	45	815	UGUCCAGAUUAUGCGUUU	472
811	AGUAGAAAGGCUUCAUA	46	811	AGUAGAAAGGCUUCAUA	46	833	UGAUGAAGCCCUUUUCUACU	473
829	AUAUCAAUGCAACGUACA	47	829	AUAUCAAUGCAACGUACA	47	851	UGUACGUUGCAUUUUGAUU	474
847	AAAGAAUAGGGCUUCUGA	48	847	AAAGAAUAGGGCUUCUGA	48	869	UCAGAAAGCCUUAUUUCUUU	475
865	ACCUGUGAAGCAACAGUCA	49	865	ACCUGUGAAGCAACAGUCA	49	887	UGACUUGUUGCUUCACAGGU	476
883	AUUGGCAUUUUGUAAGA	50	883	AUUGGCAUUUUGUAAGA	50	905	UCUUAACAAUUGCCCAUU	477
901	ACAAACUACUCACACAUC	51	901	ACAAACUACUCACACAUC	51	923	GAUGUGAGAGUAGUUUGU	478
919	CGACAAACCAUAACAUA	52	919	CGACAAACCAUAACAUA	52	941	UGAUUGUAUUUGGUUUGUCG	479
937	AUAGAUGUCCAAUAAGCA	53	937	AUAGAUGUCCAAUAAGCA	53	959	UGCUUAUUUGGACACUUAU	480
955	ACACCACGCCAGUCAAU	54	955	ACACCACGCCAGUCAAU	54	977	AUUUGACUGGGCGUGUGU	481
973	UUACUUAAGAGGCCAUACUC	55	973	UUACUUAAGAGGCCAUACUC	55	995	GAGUAGGCCUUCUAAGUAA	482
991	CUUGUCCUCAAUUUGUACUG	56	991	CUUGUCCUCAAUUUGUACUG	56	1013	CAGUACAAUUGAGGACAAG	483
1009	GCUACCAUCCCUUGAACA	57	1009	GCUACCAUCCCUUGAACA	57	1031	UGUACAAGGGAGUGGUAGC	484
1027	ACGAGAGUCAAUUGACCU	58	1027	ACGAGAGUCAAUUGACCU	58	1049	AGGUCAUUUUGAACUCUCGU	485
1045	UGGAGUUACCCUGAUGAAA	59	1045	UGGAGUUACCCUGAUGAAA	59	1067	UUUACAUCAGGGUAACUCCA	486
1063	AAAAUAAGAGAGCUUCCG	60	1063	AAAAUAAGAGAGCUUCCG	60	1085	CGGAAGCUCUCUUAUUUUU	487
1081	GUAGGGGACGAAUUGACC	61	1081	GUAGGGGACGAAUUGACC	61	1103	GGUCAUUUCGUCGCCUUAAC	488
1099	CAAAGCAUUUCCCAUGCCA	62	1099	CAAAGCAUUUCCCAUGCCA	62	1121	UGGCAUGGGAAUUGCUUUG	489
1117	AACAUAUUCUACAGUGUUC	63	1117	AACAUAUUCUACAGUGUUC	63	1139	GAACACUGUAGAAUUAUUGU	490
1135	CUUACUUAUGACAAAUGC	64	1135	CUUACUUAUGACAAAUGC	64	1157	GCAUUUUUGCAAAUUAUAG	491
1153	CAGAACAAAGACAAAGGAC	65	1153	CAGAACAAAGACAAAGGAC	65	1175	GUCCUUUGUCUUUUGUUCUG	492
1171	CUUUUAUACUUGUGUGUAA	66	1171	CUUUUAUACUUGUGUGUAA	66	1193	UUACACGACAAAGUUAUAAAG	493

1189	AGGAGUGGACCAUUAUUA	67	1189	AGGAGUGGACCAUUAUUA	67	1211	UGAAUGAUGGUCCACUCCU	494
1207	AAUUCUGUUAACACUCCAG	68	1207	AAUUCUGUUAACACUCCAG	68	1229	CUGAGGUGUUAACACAGAUUU	495
1225	GUGCAUUAUUAUUAUUAAG	69	1225	GUGCAUUAUUAUUAUUAAG	69	1247	CUUUAUUAUUAUUAUUAUUA	496
1243	GCAUUAUUAUUAUUAUUA	70	1243	GCAUUAUUAUUAUUAUUA	70	1265	GUUUCACAGUGAUGAUGAUG	497
1261	CAUCGAAACAGAGUGGUC	71	1261	CAUCGAAACAGAGUGGUC	71	1283	GCACUUGUGUUAUUAUUAUUA	498
1279	CUUGAAACAGAGUGGUC	72	1279	CUUGAAACAGAGUGGUC	72	1301	UGCCAGCUACGGUUAUUAUUA	499
1297	AAGCGGUCUUAUUAUUAUUA	73	1297	AAGCGGUCUUAUUAUUAUUA	73	1319	AGAGCCGUAAGAGCCGCUU	500
1315	UCUUAUUAUUAUUAUUAUUA	74	1315	UCUUAUUAUUAUUAUUAUUA	74	1337	AUGCCUUAUUAUUAUUAUUA	501
1333	UUUCCUCCGCGGAGUUG	75	1333	UUUCCUCCGCGGAGUUG	75	1355	CAACUCCGCGGAGGGAUA	502
1351	GUUUGUUAUUAUUAUUAUUA	76	1351	GUUUGUUAUUAUUAUUAUUA	76	1373	AGCCAUUUUAUUAUUAUUA	503
1369	UUACCUUGCGACUGAGAAU	77	1369	UUACCUUGCGACUGAGAAU	77	1391	AUUUCACAGUCGAGGUAUA	504
1387	UCUGCUCGUAUUAUUAUUA	78	1387	UCUGCUCGUAUUAUUAUUA	78	1409	GAGUCAAUUAUUAUUAUUA	505
1405	CGUGGCUACUCGUUAUUA	79	1405	CGUGGCUACUCGUUAUUA	79	1427	UAUUAACGAGUAGCCACG	506
1423	AUCAAGGACGUUAUUAUUA	80	1423	AUCAAGGACGUUAUUAUUA	80	1445	CUUCAGUUAUUAUUAUUA	507
1441	GAGGAGGAGGUAUUAUUA	81	1441	GAGGAGGAGGUAUUAUUA	81	1463	UAUUAUUAUUAUUAUUAUUA	508
1459	ACAUCUUGCUGAGCAUUA	82	1459	ACAUCUUGCUGAGCAUUA	82	1481	UUUAGCUCAGCAAGAUUGU	509
1477	AAACAGUCAAUUAUUAUUA	83	1477	AAACAGUCAAUUAUUAUUA	83	1499	UAAACACAUUAUUAUUAUUA	510
1495	AAAAACCUACUGCCACUC	84	1495	AAAAACCUACUGCCACUC	84	1517	GAGUGGACGAGGAGUUAUUA	511
1513	CUAAUUGUUAUUAUUAUUA	85	1513	CUAAUUGUUAUUAUUAUUA	85	1535	GUUUCACAUUAUUAUUAUUA	512
1531	CCCCAGUUAUUAUUAUUA	86	1531	CCCCAGUUAUUAUUAUUA	86	1553	CCUUAUUAUUAUUAUUAUUA	513
1549	GCCGUGUUAUUAUUAUUA	87	1549	GCCGUGUUAUUAUUAUUA	87	1571	CUGGAAACGAGUAGCCACG	514
1567	GACCCGCGUUAUUAUUAUUA	88	1567	GACCCGCGUUAUUAUUAUUA	88	1589	GUGGUAAGAGAGCCGCGGUC	515
1585	CUGGGCAGCAGCAUUAUUA	89	1585	CUGGGCAGCAGCAUUAUUA	89	1607	CAUUAUGCGGUUAUUAUUA	516
1603	CUGACUUGUUAUUAUUAUUA	90	1603	CUGACUUGUUAUUAUUAUUA	90	1625	GAUUAUGCGGUUAUUAUUA	517
1621	GGUAUCCCUUAUUAUUAUUA	91	1621	GGUAUCCCUUAUUAUUAUUA	91	1643	UUUAGGUAUUAUUAUUAUUA	518
1639	AUCAAGUGGUGUUAUUAUUA	92	1639	AUCAAGUGGUGUUAUUAUUA	92	1661	GGUGCCAGCAUUAUUAUUA	519
1657	CCUUGUAUUAUUAUUAUUA	93	1657	CCUUGUAUUAUUAUUAUUA	93	1679	AAUGAUUAUUAUUAUUAUUA	520
1675	UCCGAAGCAAGGUGUUAUUA	94	1675	UCCGAAGCAAGGUGUUAUUA	94	1697	AGUCACCUUUAUUAUUAUUA	521
1693	UUUUGUUAUUAUUAUUAUUA	95	1693	UUUUGUUAUUAUUAUUAUUA	95	1715	CUUUAUUAUUAUUAUUAUUA	522
1711	GAGUCCUUAUUAUUAUUAUUA	96	1711	GAGUCCUUAUUAUUAUUAUUA	96	1733	CAUCCAGGAUUAUUAUUAUUA	523
1729	GCUGACAGCAUUAUUAUUA	97	1729	GCUGACAGCAUUAUUAUUA	97	1751	UUUCCAUUAUUAUUAUUAUUA	524
1747	AACAGAAUUAUUAUUAUUA	98	1747	AACAGAAUUAUUAUUAUUA	98	1769	UGAUGCUCUUAUUAUUAUUA	525
1765	ACUCAGCGCAUUAUUAUUA	99	1765	ACUCAGCGCAUUAUUAUUA	99	1787	UUUUAUUAUUAUUAUUAUUA	526
1783	AUAGAAGGAAUUAUUAUUA	100	1783	AUAGAAGGAAUUAUUAUUA	100	1805	UCUUAUUAUUAUUAUUAUUA	527
1801	AUGGCUAGCACCUUUGUUA	101	1801	AUGGCUAGCACCUUUGUUA	101	1823	CAACCAAGGUGUUAUUAUUA	528
1819	GUGGCUAGCACCUUUGUUA	102	1819	GUGGCUAGCACCUUUGUUA	102	1841	AAUUAUUAUUAUUAUUAUUA	529

1837	UCUGGAUUCUACAUUUUGCA	103	1837	UCUGGAUUCUACAUUUUGCA	103	1859	UGCAAAUUGUAGAUUCCAGA	530
1855	AUAGCUUCCAAUAAAGUUG	104	1855	AUAGCUUCCAAUAAAGUUG	104	1877	CAACUUAUUGGAAGCUAU	531
1873	GGGACUGUGGGAAGAAACA	105	1873	GGGACUGUGGGAAGAAACA	105	1895	UGUUUUUUUUUUUUUUUUUU	532
1891	AUAAGCUUUUAUAUCACAG	106	1891	AUAAGCUUUUAUAUCACAG	106	1913	CUGUGAUUAAAGCUUAU	533
1909	GAUGUGCCAAUUGGUUUC	107	1909	GAUGUGCCAAUUGGUUUC	107	1931	GAAACCAUUGGCAACUC	534
1927	CAUGUUAACUUGGAAAAA	108	1927	CAUGUUAACUUGGAAAAA	108	1949	UUUUUUUUUUUUUUUUUUUU	535
1945	AUGCCGACGGAAGGAGG	109	1945	AUGCCGACGGAAGGAGG	109	1967	CCUCUCCUUCGUGGGCAU	536
1963	GACCUGAAACUGUCUUGCA	110	1963	GACCUGAAACUGUCUUGCA	110	1985	UGCAAGACAGUUCAGGUC	537
1981	ACAGUUAACAAGUUCUUAU	111	1981	ACAGUUAACAAGUUCUUAU	111	2003	AUAAGAACUUGUUAACUGU	538
1999	UACAGAGACGUUAACUUGGA	112	1999	UACAGAGACGUUAACUUGGA	112	2021	UCCAAGUAAACGUCUCUGUA	539
2017	AUUUUAACUGGGACAUUA	113	2017	AUUUUAACUGGGACAUUA	113	2039	UACUGUCCGCGAGUAAAAU	540
2035	AUAACAGAAACUAGCUCU	114	2035	AUAACAGAAACUAGCUCU	114	2057	AGUGCAUUGUUCUGUUAU	541
2053	UACAGUUAUAGCAAGCAAA	115	2053	UACAGUUAUAGCAAGCAAA	115	2075	UUUGCUUGCUAAUACUGUA	542
2071	AAAAGGCCAUACACUUAAG	116	2071	AAAAGGCCAUACACUUAAG	116	2093	CCUUAUGUAGUGGCCAUUUU	543
2089	GAGCACUCCAUACACUCUUA	117	2089	GAGCACUCCAUACACUCUUA	117	2111	UAAGAGUGAUGGAGUGCUC	544
2107	AUUCUUAACCAUCAUGAAUG	118	2107	AUUCUUAACCAUCAUGAAUG	118	2129	CAUUAUGAUGGUAAAGAU	545
2125	GUUUCUUGCAAGAUUCAG	119	2125	GUUUCUUGCAAGAUUCAG	119	2147	CUGAAUCUUGCAGGGAAC	546
2143	GGCACCUAUUGCCUGCAGAG	120	2143	GGCACCUAUUGCCUGCAGAG	120	2165	CUCUGCAGGCAUAGGUGCC	547
2161	GCCAGGAUUAUACACAG	121	2161	GCCAGGAUUAUACACAG	121	2183	CUGUGUAUACAUUCCUGGC	548
2179	GGGGAAGAAUCCUCCAGA	122	2179	GGGGAAGAAUCCUCCAGA	122	2201	UCUGGAGGAUUUUCUCCCC	549
2197	AAGAAAGAAUUAACAUA	123	2197	AAGAAAGAAUUAACAUA	123	2219	UGAUUUGUAUUUUCUUCUU	550
2215	AGAGUACAGGAAGCAGCAU	124	2215	AGAGUACAGGAAGCAGCAU	124	2237	AUGGUGCUUCCUGAUCUCU	551
2233	UACCUCCUGCGAAACCUCA	125	2233	UACCUCCUGCGAAACCUCA	125	2255	UGAGGUUUUCGAGGAGGUA	552
2251	AGUGAUCACACAGUGGCCA	126	2251	AGUGAUCACACAGUGGCCA	126	2273	UGGCCACUUGUGAUCACU	553
2269	AUCAGCAGUUCACACCAU	127	2269	AUCAGCAGUUCACACCAU	127	2291	AAGUGGGAACUGCUGAU	554
2287	UUAGACUGUUAUGCUUAUG	128	2287	UUAGACUGUUAUGCUUAUG	128	2309	CAUUAAGCAUGACAGUCUAA	555
2305	GGUGUCCCGAGCCUACAGA	129	2305	GGUGUCCCGAGCCUACAGA	129	2327	UCUGAGGCUUGGAGGACAC	556
2323	AUCACUUGGUUUAACAA	130	2323	AUCACUUGGUUUAACAA	130	2345	UGUUUUUAAACCAAGUGAU	557
2341	AACACAAAAUACAACAAG	131	2341	AACACAAAAUACAACAAG	131	2363	CUUGUUGAUUUUUGGUUU	558
2359	GAGCCUGGAUUUUUUUAG	132	2359	GAGCCUGGAUUUUUUUAG	132	2381	CUAAAAAUUCCAGGCUC	559
2377	GGACCAAGGAAGCAGCAGC	133	2377	GGACCAAGGAAGCAGCAGC	133	2399	GCGUGCGUUCUCCUGGUCC	560
2395	CUGUUUAUUGAAAGAGUCA	134	2395	CUGUUUAUUGAAAGAGUCA	134	2417	UGACUCUUUCAAUAAACAG	561
2413	ACAGAAGAGGAAGAGGUG	135	2413	ACAGAAGAGGAAGAGGUG	135	2435	CACCUUACUCCUUCUCUG	562
2431	GUCUUAUCACUGCAAGGCCA	136	2431	GUCUUAUCACUGCAAGGCCA	136	2453	UGGCUUUGCAGUGAUAGAC	563
2449	ACCAACCAAGAGGCGUCUG	137	2449	ACCAACCAAGAGGCGUCUG	137	2471	CAGAGCCCUUCUGGUUGGU	564
2467	GUGGAAAGUUCAGCAUACC	138	2467	GUGGAAAGUUCAGCAUACC	138	2489	GGUAGUCUGAACUUCUCCAG	565

2485	CUCACUGUUCAGGAACCU	139	2485	CUCACUGUUCAGGAACCU	139	2507	AGGUUCCUUGAACAGUGAG	566
2503	UCGGACAAGUCUAAUCUGG	140	2503	UCGGACAAGUCUAAUCUGG	140	2525	CCAGAUUAGACUUGUCCGA	567
2521	GAGCUGAUCACUCUAAACAU	141	2521	GAGCUGAUCACUCUAAACAU	141	2543	AUGUUAGAGUGAUCAGCUC	568
2539	UGCACCUGUGGUCGCGA	142	2539	UGCACCUGUGGUCGCGA	142	2561	UCGCAGCCACACAGGUGCA	569
2557	ACUCUCUUGGUCUCCU	143	2557	ACUCUCUUGGUCUCCU	143	2579	AUAGGAGCCAGAGAGAGU	570
2575	UUAACCCUCCUUAUCCGAA	144	2575	UUAACCCUCCUUAUCCGAA	144	2597	UUCGGAUAAAGGAGGUA	571
2593	AAAUGAAAGGUCUCCU	145	2593	AAAUGAAAGGUCUCCU	145	2615	AAGAAGACCUUUAUUAUUU	572
2611	UCUGAAUAAAGACUGACU	146	2611	UCUGAAUAAAGACUGACU	146	2633	AGUCAGUCUUAUUAUUCAGA	573
2629	UACCUAUCUUAUUAUUGG	147	2629	UACCUAUCUUAUUAUUGG	147	2651	CCAUUAUUAUUAUUAUAGGUA	574
2647	GACCCAGAUAGGUUCCU	148	2647	GACCCAGAUAGGUUCCU	148	2669	AAGGAACUUCACUCUGGGUC	575
2665	UUGGAUGAGCAGUGAGC	149	2665	UUGGAUGAGCAGUGAGC	149	2687	GCUCACACUUCACUCUCCAA	576
2683	CGGCUCCUUAUGAUGCCA	150	2683	CGGCUCCUUAUGAUGCCA	150	2705	UGGCAUCUUAAGGAGCGG	577
2701	AGCAAGUGGGAGUUGCCC	151	2701	AGCAAGUGGGAGUUGCCC	151	2723	GGGCAACUCCACUUGCU	578
2719	CGGAGAGACUUAACUGG	152	2719	CGGAGAGACUUAACUGG	152	2741	CCAGUUAUAGUCUCUCCGG	579
2737	GGCAAAUCACUUGGAAGAG	153	2737	GGCAAAUCACUUGGAAGAG	153	2759	CUCUCCAAAGUGAUUUGCC	580
2755	GGGGCUUUGGAAAGUGG	154	2755	GGGGCUUUGGAAAGUGG	154	2777	CCACUUAUCCAAAGCCCC	581
2773	GUUCAAGCAUCAGCAUUG	155	2773	GUUCAAGCAUCAGCAUUG	155	2795	CAAAUGCUGAUGCUUGAAC	582
2791	GGCAUUAAGAAUACCCUA	156	2791	GGCAUUAAGAAUACCCUA	156	2813	UAGGUAUUAUUAUUAUGCC	583
2809	ACUGCCGACUGUGGCG	157	2809	ACUGCCGACUGUGGCG	157	2831	CAGCCACAGUCCGGCAGCU	584
2827	GUGAAUUGCUAAGAGG	158	2827	GUGAAUUGCUAAGAGG	158	2849	CCUCUUAUUAUUAUUAUAC	585
2845	GGGGCCACGGCAGCGAGU	159	2845	GGGGCCACGGCAGCGAGU	159	2867	ACUCGUGGCGGUGGCGCC	586
2863	UACAAAGCUCUGAUGACUG	160	2863	UACAAAGCUCUGAUGACUG	160	2885	CAGUCAUCAGAGCUUUGUA	587
2881	GAGCUAAAACUUGACCC	161	2881	GAGCUAAAACUUGACCC	161	2903	GGGUAAGAUUUUUAAGCUC	588
2899	CACAUUGGCCACCAUCUGA	162	2899	CACAUUGGCCACCAUCUGA	162	2921	UCAGAUGGUGGCCAAUGUG	589
2917	AACGUGUUAACCUUGG	163	2917	AACGUGUUAACCUUGG	163	2939	CAGCAGGUUAACCAACGUU	590
2935	GGAGCCUGCACCAGCAAG	164	2935	GGAGCCUGCACCAGCAAG	164	2957	CUUGCUUGGUGCAGGCUCC	591
2953	GGAGGGCCUUGAUGGUGA	165	2953	GGAGGGCCUUGAUGGUGA	165	2975	UCACCAUCAGAGGCGCCUCC	592
2971	AUUGUUAUUAUCUGCAAU	166	2971	AUUGUUAUUAUCUGCAAU	166	2993	AUUUGCAGUUAUUAACAAU	593
2989	UAUGGAAUUCUCUCCAAU	167	2989	UAUGGAAUUCUCUCCAAU	167	3011	AGUUGGAGAGAUUUUCCAU	594
3007	UACCUCAAGAGCAACCGUG	168	3007	UACCUCAAGAGCAACCGUG	168	3029	CACGUUUGCUUUGAGGUA	595
3025	GACUUAUUUUUUCUGAACA	169	3025	GACUUAUUUUUUCUGAACA	169	3047	UGUUGAGAAAAAUUAAGUC	596
3043	AAGGAUGCAGCAGCAGACACA	170	3043	AAGGAUGCAGCAGCAGACACA	170	3065	UGUUGAUGGUGGCAUCCUU	597
3061	AUGGAGCCUUAAGAAAGAAA	171	3061	AUGGAGCCUUAAGAAAGAAA	171	3083	UUUCUUUUAUUAAGGCUCCAU	598
3079	AAAUGGAGCAGGCGCUGG	172	3079	AAAUGGAGCAGGCGCUGG	172	3101	CCAGGCCUGGCGCAGGCUUUU	599
3097	GAACAAGGCAAGAAACCAA	173	3097	GAACAAGGCAAGAAACCAA	173	3119	UUGGUUUUUGGCGUUGUUC	600
3115	AGACUAGAUAGCGUCAGCA	174	3115	AGACUAGAUAGCGUCAGCA	174	3137	UGGUGACGCUAUCUAGUCU	601

3133	AGCAGCGAAAGCUUUGCGA	175	3133	AGCAGCGAAAGCUUUGCGA	175	3155	UCGCAAGCUUUGCGUCU	602
3151	AGCUCCGGCUUUCAGGAAG	176	3151	AGCUCCGGCUUUCAGGAAG	176	3173	CUUCCUGAAAGCCGGAGCU	603
3169	GAUAAAGUCUCAGGUAUG	177	3169	GAUAAAGUCUCAGGUAUG	177	3191	CAUCACUCAGACUUUUAUC	604
3187	GUUGAGGAAGAGGAGUAU	178	3187	GUUGAGGAAGAGGAGUAU	178	3209	AAUCCUCCUCUUCUCAAAC	605
3205	UCUGACGGUUCUACAAGG	179	3205	UCUGACGGUUCUACAAGG	179	3227	CCUUGUAGAAACCCGUCAGA	606
3223	GAGCCCAUCAGUAUGGAAG	180	3223	GAGCCCAUCAGUAUGGAAG	180	3245	CUUCCAUAUGUAGGGCUC	607
3241	GAUCUGAUUUCUACAGUU	181	3241	GAUCUGAUUUCUACAGUU	181	3263	AACUGUAAGAAUUCAGAU	608
3259	UUUCAAGUGGCCAGAGGCA	182	3259	UUUCAAGUGGCCAGAGGCA	182	3281	UGCCUCUGGCCACUUGAAA	609
3277	AUGGAGUCCUGUCUCCA	183	3277	AUGGAGUCCUGUCUCCA	183	3299	UGGAAGACAGGAACUCCAU	610
3295	AGAAAGUGCAUUCUCCGG	184	3295	AGAAAGUGCAUUCUCCGG	184	3317	CCCGAUGAAUGCACUUUCU	611
3313	GACUUGGAGCGAGAAACA	185	3313	GACUUGGAGCGAGAAACA	185	3335	UGUUUCUGCGUGCCAGGUC	612
3331	AUUCUUUUAUCUGAGAACA	186	3331	AUUCUUUUAUCUGAGAACA	186	3353	UGUUCUCAGAUAAAAGAAU	613
3349	AACGUGGUGAAGAUUUGUG	187	3349	AACGUGGUGAAGAUUUGUG	187	3371	CACAAAUCUACCCACGCUU	614
3367	GAUUUUGGCCUUGCCCGGG	188	3367	GAUUUUGGCCUUGCCCGGG	188	3389	CCCGGCAAGGCCAAAAUUC	615
3385	GAUAAUUAUAAAGAACCCCG	189	3385	GAUAAUUAUAAAGAACCCCG	189	3407	CGGGGUUCUUAUAAUUAUC	616
3403	GAUUAUGUGAGAAAAGGAG	190	3403	GAUUAUGUGAGAAAAGGAG	190	3425	CUCCUUUUCUCACAUAAUC	617
3421	GAUACUGGACUUCUCCUGA	191	3421	GAUACUGGACUUCUCCUGA	191	3443	UCAGAGGAAGUCGAGUAUC	618
3439	AAUUGGAUGGCUCCCGAAU	192	3439	AAUUGGAUGGCUCCCGAAU	192	3461	AUUCGGGAGCCAUCCAUUU	619
3457	UCUAUCUUUGACAAAUCU	193	3457	UCUAUCUUUGACAAAUCU	193	3479	AGAUUUUGUCAAGAUAGA	620
3475	UACAGCAACCAAGAGCGACG	194	3475	UACAGCAACCAAGAGCGACG	194	3497	CGUCGCUUUGGUGCUGUA	621
3493	GUUGUGUCUUUACGGAGUAU	195	3493	GUUGUGUCUUUACGGAGUAU	195	3515	AUACUCCGUAAGACCACAC	622
3511	UUGCUGUGGGAUUCUUCU	196	3511	UUGCUGUGGGAUUCUUCU	196	3533	AGAAGAUUUCACACAGCAA	623
3529	UCCUAGGUGGGUCCUCCAU	197	3529	UCCUAGGUGGGUCCUCCAU	197	3551	AUGGAGACCCACCUCUAGGA	624
3547	UACCCAGGAGUACAAAUGG	198	3547	UACCCAGGAGUACAAAUGG	198	3569	CCAUUUUGUACUCCUUGGUA	625
3565	GAUGAGGACUUUUGCAGUC	199	3565	GAUGAGGACUUUUGCAGUC	199	3587	GACUGCAAAAGUCCUUAUC	626
3583	CGCCUGAGGGAAGGCAUGA	200	3583	CGCCUGAGGGAAGGCAUGA	200	3605	UCAUGCCUCCUCCAGGCG	627
3601	AGGAUGAGAGCUCCUAGAU	201	3601	AGGAUGAGAGCUCCUAGAU	201	3623	ACUACGAGCUCUCAUCCU	628
3619	UACUCUACUCCUGAAAUCU	202	3619	UACUCUACUCCUGAAAUCU	202	3641	AGAUUUCAGGAGUAGAGUA	629
3637	UAUCAGAUCAUGCUGGACU	203	3637	UAUCAGAUCAUGCUGGACU	203	3659	AGUCCAGCAUGAUUCUGUA	630
3655	UGCUGGCACAGAGACCCAA	204	3655	UGCUGGCACAGAGACCCAA	204	3677	UUGGGUCUCUGUGGCCAGCA	631
3673	AAAGAAAGGCCAAGAUUUG	205	3673	AAAGAAAGGCCAAGAUUUG	205	3695	CAAAUUCUUGCCUUCUUCU	632
3691	GCAGAACUUGUGGAAAAC	206	3691	GCAGAACUUGUGGAAAAC	206	3713	GUUUUCCACAAGUUCUGC	633
3709	CUAGGUGAUUUGCUUCAAG	207	3709	CUAGGUGAUUUGCUUCAAG	207	3731	CUUGAAGCAAAUCCACUAG	634
3727	GCAAUGUUAACACAGGAUG	208	3727	GCAAUGUUAACACAGGAUG	208	3749	CAUCCUGUUGUACAUUUGC	635
3745	GGUAAAGACUACAUCCCAA	209	3745	GGUAAAGACUACAUCCCAA	209	3767	UUGGGAUGUAGUCCUUAACC	636
3763	AUCAUUGCCAUACUGACAG	210	3763	AUCAUUGCCAUACUGACAG	210	3785	CUGUCAGUAUGGCAUUGAU	637



3781	GGAAUAGUGGGUUUACAU	211	3781	GGAAUAGUGGGUUUACAU	211	3803	AUGUAAACCCACAUUUCC	638
3789	UACUCAACUCCUGCCUUCU	212	3789	UACUCAACUCCUGCCUUCU	212	3821	AGAAAGCAGGAGUUGAGUA	639
3817	UCUGAGGACUUCUUAAGG	213	3817	UCUGAGGACUUCUUAAGG	213	3839	CCUUGAAGAGUCCUCAGA	640
3835	GAAAGUUAUUCAGCUCGGA	214	3835	GAAAGUUAUUCAGCUCGGA	214	3857	UCGGAGCUGAAAUACUUC	641
3853	AAGUUUAUUCAGGAAGCU	215	3853	AAGUUUAUUCAGGAAGCU	215	3875	AGCUCCUUGAAUUAACU	642
3871	UCUGAUGAUGUCAGAUUG	216	3871	UCUGAUGAUGUCAGAUUG	216	3893	CAUAUCUGACAUCAUCAGA	643
3889	GUAAUUGCUUUAAGUUA	217	3889	GUAAUUGCUUUAAGUUA	217	3911	UGAACUUGAAAGCAUUUAC	644
3907	AUGAGCCUGGAAAGAUCA	218	3907	AUGAGCCUGGAAAGAUCA	218	3929	UGAUUCUUCAGGCUCAU	645
3925	AAAACCUUUGAAGACU	219	3925	AAAACCUUUGAAGACU	219	3947	AAAGUUCUCAAAGGUUU	646
3943	UUACCGAUGCCACCUCCA	220	3943	UUACCGAUGCCACCUCCA	220	3965	UGGAGGUGGCAUUCGGUAA	647
3961	AUGUUUGAUGACUACCGG	221	3961	AUGUUUGAUGACUACCGG	221	3983	CCUGGUGGCUUGUCGCC	648
3979	GGCGACAGCAGCACUCUG	222	3979	GGCGACAGCAGCACUCUG	222	4001	ACAGAGUGGUGGAGGCCAA	649
3997	UUGGCCUCUCCCAUGCUGA	223	3997	UUGGCCUCUCCCAUGCUGA	223	4018	UCAGCAUGGGAGAGGCCAA	650
4015	AAGCGCUUACCCUGGACUG	224	4015	AAGCGCUUACCCUGGACUG	224	4037	CAGUCCAGGUGAAGCGCUU	651
4033	GACAGCAACCCCAAGGCCU	225	4033	GACAGCAACCCCAAGGCCU	225	4055	AGGCCUUGGUUUGCUGUC	652
4051	UCGCUCAAGAUUGACUUGA	226	4051	UCGCUCAAGAUUGACUUGA	226	4073	UCAAGUCAUUCUUGAGCGA	653
4069	AGAGUAACAGUAAAGUA	227	4069	AGAGUAACAGUAAAGUA	227	4091	UACUUUACUUGGUUACUCU	654
4087	AAGGAGUCGGGCGUCUG	228	4087	AAGGAGUCGGGCGUCUG	228	4109	CAGACAGCCCGACUCU	655
4105	GAUGUACAGGCCCAGUU	229	4105	GAUGUACAGGCCCAGUU	229	4127	AACUGGGCCUGCUGACAU	656
4123	UUCUGCAUUCAGCUGUG	230	4123	UUCUGCAUUCAGCUGUG	230	4145	CACAGCUGGAUUGGCAGAA	657
4141	GGGCACGUCAGCGAAGGCA	231	4141	GGGCACGUCAGCGAAGGCA	231	4163	UGCCUUCGUCAGCGUCCC	658
4159	AAGCGCAGGUUACCCUACG	232	4159	AAGCGCAGGUUACCCUACG	232	4181	CGUAGGUGAACCCUGCGCUU	659
4177	GACCAGCGUGAGCUGGAAA	233	4177	GACCAGCGUGAGCUGGAAA	233	4199	UUUCCAGCUCAGCGUGGUC	660
4195	AGGAAAUUCGUGUGCU	234	4195	AGGAAAUUCGUGUGCU	234	4217	AGCAGCAGCGAUUUUCCU	661
4213	UCCCCGCCCCAGACUACA	235	4213	UCCCCGCCCCAGACUACA	235	4235	UGUAGUCUGGGGGCGGGGA	662
4231	AACUCGGUUGUCCUGUACU	236	4231	AACUCGGUUGUCCUGUACU	236	4253	AGUACAGGACCCAGAGUU	663
4249	UCCACCCACCCAUUAGA	237	4249	UCCACCCACCCAUUAGA	237	4271	UCUAGUUGGUGGGGUGGA	664
4267	AGUUUGACACGAAGCCUUA	238	4267	AGUUUGACACGAAGCCUUA	238	4289	UAAGGCUUCGUGUCAAAACU	665
4285	AUUUCUAGAGCACAUGUG	239	4285	AUUUCUAGAGCACAUGUG	239	4307	CACAUUGGCUUCUAGAAAU	666
4303	GUUUUUAUACCCAGGAA	240	4303	GUUUUUAUACCCAGGAA	240	4325	UUCUUGGGGUAUAAAUAC	667
4321	AACUAGCUUUUUGCCAGUUA	241	4321	AACUAGCUUUUUGCCAGUUA	241	4343	AUACUGGCAAAAGCUAGUU	668
4339	UUUAGCAUUAUUAAGUUA	242	4339	UUUAGCAUUAUUAAGUUA	242	4361	UAAACUUUAUUAUGCAUAA	669
4357	ACACCUUUAUUCUCCAUUG	243	4357	ACACCUUUAUUCUCCAUUG	243	4379	CAUGGAAAGCAUAAAGGUGU	670
4375	GGAGCCAGCUGCUUUUUG	244	4375	GGAGCCAGCUGCUUUUUG	244	4397	CAAAAAGCAGCUGGCUCCC	671
4393	GUGAUUUUUUUUAUAGUGC	245	4393	GUGAUUUUUUUUAUAGUGC	245	4415	GCACUUAUAAAAAUACAC	672
4411	CUUUUUUUUUUUUAGACUAC	246	4411	CUUUUUUUUUUUUAGACUAC	246	4433	GUUAGUCAAAAAAAG	673

4429	CAAGAAUGUAACUCCAGAU	247	4429	CAAGAAUGUAACUCCAGAU	247	4451	AUCUGGAGUUAUAUUCUUG	674
4447	UAGAGAAUAGUGACAAGU	248	4447	UAGAGAAUAGUGACAAGU	248	4469	ACUUGUCACUAUUCUCUA	675
4465	UGAAGAACACUACUGCUAA	249	4465	UGAAGAACACUACUGCUAA	249	4487	UUAGCAGUAGUGUUCUUA	676
4483	AAUCCUACUGUUAUUCUAGU	250	4483	AAUCCUACUGUUAUUCUAGU	250	4505	ACUGAGUAAACUAGAGAUU	677
4501	UGUAGAGAAUUCUUCU	251	4501	UGUAGAGAAUUCUUCU	251	4523	AGGAAGAUUUCUUAACA	678
4519	UAAACCAUAGAUUCU	252	4519	UAAACCAUAGAUUCU	252	4541	AGGAAGUUAUUGGUUUA	679
4537	UGUCCAAACCCCGCCACC	253	4537	UGUCCAAACCCCGCCACC	253	4559	GGUGCGGGGUUGGAGCA	680
4555	CUAGGACACGAGGACCA	254	4555	CUAGGACACGAGGACCA	254	4577	UGGUCCUGCGUGCCCGAG	681
4573	AGUUUAUAGGAGCUGC	255	4573	AGUUUAUAGGAGCUGC	255	4595	GCAGCUCCUCAAUCAAACU	682
4591	CACUGAUACCCCAUUGCAU	256	4591	CACUGAUACCCCAUUGCAU	256	4613	AUGCAUUGGGUGAUCAGUG	683
4609	UCACGUACCCACUUGGCC	257	4609	UCACGUACCCACUUGGCC	257	4631	GGCCAGUGGGUGAUCAGUG	684
4627	CAGCCUUGCAGCCCAAAAC	258	4627	CAGCCUUGCAGCCCAAAAC	258	4649	GUUUGGGUGCAGGGCUG	685
4645	CCCAGGGCAACAGCCCGU	259	4645	CCCAGGGCAACAGCCCGU	259	4687	ACGGGUUGUUGCCUUGG	686
4663	UUAGCCAGGGGAUCACU	260	4663	UUAGCCAGGGGAUCACU	260	4685	AGUGAUCCCUUGGGGCUAA	687
4681	UGGCUGCCUGAGCAACAU	261	4681	UGGCUGCCUGAGCAACAU	261	4703	AUGUUGCUCAGGCCAGCCA	688
4699	UCUCGGGAGUCCUAGCA	262	4699	UCUCGGGAGUCCUAGCA	262	4721	UGUAGAGGACUCCCGAGA	689
4717	AGGCCUAGACAUUGAGG	263	4717	AGGCCUAGACAUUGAGG	263	4739	CCUCACAUUGUUAAGCCU	690
4735	GAGGAAAGGAAAAAAGC	264	4735	GAGGAAAGGAAAAAAGC	264	4757	GCUUUUUUCCUUAUCCUC	691
4753	CAAAAAGCAGGAGAAAA	265	4753	CAAAAAGCAGGAGAAAA	265	4775	UUUUCUCCUUGCUUUUUG	692
4771	AGAGAAACCGGAGAGGC	266	4771	AGAGAAACCGGAGAGGC	266	4793	GCUIUCUCCCGGUUUUCUCU	693
4789	CAUGAGAAAGAAUUUGAGA	267	4789	CAUGAGAAAGAAUUUGAGA	267	4811	UCUCAAUUUCUUCUCAUG	694
4807	ACGCACCAUUGGGCACGG	268	4807	ACGCACCAUUGGGCACGG	268	4829	CCGUGCCCAUUGGUGCGU	695
4825	GAGGGGACGGGCUACAGC	269	4825	GAGGGGACGGGCUACAGC	269	4847	GCUGAGCCCGUCCCGCUC	696
4843	CAUUGCCAUUUCAGUGGCU	270	4843	CAUUGCCAUUUCAGUGGCU	270	4865	AGCCACUGAAUUGGCAUUG	697
4861	UUCCCAGCUUGACCCUUC	271	4861	UUCCCAGCUUGACCCUUC	271	4883	GAAGGUGACAGCUGGGAA	698
4879	CUACAUUUGAGGGCCAGC	272	4879	CUACAUUUGAGGGCCAGC	272	4901	GCUGGGCCUCAAUUGUAG	699
4897	CCAGGACAGUUGGACAGC	273	4897	CCAGGACAGUUGGACAGC	273	4919	GCUGUCCAUUGCUCUCCUGG	700
4915	CGAUGAGGGACAUUUUCU	274	4915	CGAUGAGGGACAUUUUCU	274	4937	AGAAAUGUCCCGCUCAUCG	701
4933	UGGAUUCUGGGAGGCAAGA	275	4933	UGGAUUCUGGGAGGCAAGA	275	4955	UCUUGCCUCCAGAAUCCA	702
4951	AAAAGGACAAUUAUUCUUU	276	4951	AAAAGGACAAUUAUUCUUU	276	4973	AAAAGAUUUUGUCCUUUU	703
4969	UUUGGAACUAAAGCAAAU	277	4969	UUUGGAACUAAAGCAAAU	277	4991	AUUUGCUUUAUUAUCCAAA	704
4987	UUUAGACCUUUUACCUAUGG	278	4987	UUUAGACCUUUUACCUAUGG	278	5009	CCAUAGGUAAAGGUUUA	705
5005	GAAGUGGUUUAUUGUCCAU	279	5005	GAAGUGGUUUAUUGUCCAU	279	5027	AUGGACAUAGAACCAUUC	706
5023	UUCUCAUUCGUGGCAUGUU	280	5023	UUCUCAUUCGUGGCAUGUU	280	5045	AACAUGCCACGAAUGAGAA	707
5041	UUUGAUUUUGUAGCACUGAG	281	5041	UUUGAUUUUGUAGCACUGAG	281	5063	CUCAGUGCUACAAUCAA	708
5059	GGGUGGCACUACACUCUGA	282	5059	GGGUGGCACUACACUCUGA	282	5081	UCAGAGUUGAGUGCCACCC	709

5077	AGCCCAUACUUIUGGCUCC	283	5077	AGCCCAUACUUIUGGCUCC	283	5099	GGAGCCAAAGUAUGGGCU	710
5095	CUCUAGUAAGACGACUGA	284	5085	CUCUAGUAAGACGACUGA	284	5117	UCAGUGCAUCUUAUAGAG	711
5113	AAAACUUAAGCCAGAGUAG	285	5113	AAAACUUAAGCCAGAGUAG	285	5135	CUAACUCUGGCUAAGUUUU	712
5131	GGUUGUCUCCAGGCCAUGA	286	5131	GGUUGUCUCCAGGCCAUGA	286	5153	UCAUGGCCUGGAGACAACC	713
5149	AUGGCCUUAACUAGAAAU	287	5149	AUGGCCUUAACUAGAAAU	287	5171	AUUUUCAGUGUAAGGCCAU	714
5167	UGUCACAUUACUUAUUUGG	288	5167	UGUCACAUUACUUAUUUGG	288	5189	CCCAAAUAGAAUGUGACA	715
5185	GUUUUAUUAUAGUCCAG	289	5185	GUUUUAUUAUAGUCCAG	289	5207	CUGGACUUAUUAUUAUAC	716
5203	GACACUUAACUUAUUUCU	290	5203	GACACUUAACUUAUUUCU	290	5225	AGAAUUUGAGUUAAGUGUC	717
5221	UUGGUUAUUAUCUGUUUUG	291	5221	UUGGUUAUUAUCUGUUUUG	291	5243	CAAAACAGAAUUAUACCAA	718
5239	GCACAGUUAAGUUGAAAG	292	5239	GCACAGUUAAGUUGAAAG	292	5261	CUUUCACACUUAACUGUGC	719
5257	GAAAGCUGAGAGAAUGAA	293	5257	GAAAGCUGAGAGAAUGAA	293	5279	UUCAUUCUUCUCAGCUUUC	720
5275	AAUUGCAGUCCUGAGGAGA	294	5275	AAUUGCAGUCCUGAGGAGA	294	5297	UCUCCUCAGGACUGCAUUU	721
5293	AGUUUUCUCCAUUACAAA	295	5293	AGUUUUCUCCAUUACAAA	295	5315	UUUUGAUUGGAGAAAACU	722
5311	ACGAGGGCUGAUGGAGAA	296	5311	ACGAGGGCUGAUGGAGAA	296	5333	UUCUCCAUUAGGCCUUGU	723
5329	AAAAGGUCAUUAAGGUCAA	297	5329	AAAAGGUCAUUAAGGUCAA	297	5351	UUGACCUUAUUGACCUUUU	724
5347	AGGGAAGACCCGUCUCUA	298	5347	AGGGAAGACCCGUCUCUA	298	5369	UAGAGACGGGUCUUCUUU	725
5365	AUACCAACCAACCAUUC	299	5365	AUACCAACCAACCAUUC	299	5387	GAUUUGUUUGGUUGUAU	726
5383	CACCAACACAGUUGGGACC	300	5383	CACCAACACAGUUGGGACC	300	5405	GGUCCAAACUGUGUUGGUG	727
5401	CCAAAACACAGGAUGUCAG	301	5401	CCAAAACACAGGAUGUCAG	301	5423	CUGACUUCUUGUGUUUUGG	728
5419	GUCACGUUUCUUAUUCAUU	302	5419	GUCACGUUUCUUAUUCAUU	302	5441	AAUGAAAGGAAACGUGAC	729
5437	UUAUUGGGGAUUCACUAU	303	5437	UUAUUGGGGAUUCACUAU	303	5459	AUAGUGGAAUUCUCCAUUA	730
5455	UCUCACACUUAUCUGAAAG	304	5455	UCUCACACUUAUCUGAAAG	304	5477	CUUUCAGAUUAUGUGUGAGA	731
5473	GGAUGUGGAAGAGCAUUAU	305	5473	GGAUGUGGAAGAGCAUUAU	305	5495	CUAAUGCUCUUCACAUCC	732
5491	GCUGGCGCAUUAUAGCAC	306	5491	GCUGGCGCAUUAUAGCAC	306	5513	GUGCUUAUUAUUGCGCCAGC	733
5509	CUUUAAAGCUCCUUGAGUAA	307	5509	CUUUAAAGCUCCUUGAGUAA	307	5531	UUACUCAGGAGCUUUAAG	734
5527	AAAAGGUGUAUUAUUAUUA	308	5527	AAAAGGUGUAUUAUUAUUA	308	5549	AAUUAACUACUCCUUAU	735
5545	UAUGCAAGGUUAUUAUUAU	309	5545	UAUGCAAGGUUAUUAUUAU	309	5567	UGGAGAAUACCUUGCAUA	736
5563	AGUUGGACUACAGGAUUAU	310	5563	AGUUGGACUACAGGAUUAU	310	5585	AAUUAUCCUGAGUCCCAACU	737
5581	UAGUUAUAGCCCAUUAU	311	5581	UAGUUAUAGCCCAUUAU	311	5603	AGUGAUGGCUCAUUAACUA	738
5599	UAGAAGAAAAGCCCAUUAU	312	5599	UAGAAGAAAAGCCCAUUAU	312	5621	AAAAUGGGCUUUUUCUUA	739
5617	UCAACUGCUUUUGAAACUUG	313	5617	UCAACUGCUUUUGAAACUUG	313	5639	CAAGUUUCAAAGCAGUUA	740
5635	GCCUGGGUCUGAGCAUGA	314	5635	GCCUGGGUCUGAGCAUGA	314	5657	UCAUGCUCAGACCCAGGC	741
5653	AUGGAAUAGGGAGACAGG	315	5653	AUGGAAUAGGGAGACAGG	315	5675	CCUGUCUCCUUAUUCUUA	742
5671	GGUAGAAAGGGCGCCUAC	316	5671	GGUAGAAAGGGCGCCUAC	316	5693	GUAGGCGCCUUAUUCUUA	743
5689	CUCUACAGGGUUAUUAAGAU	317	5689	CUCUACAGGGUUAUUAAGAU	317	5711	AUCUUUAGACCCUUAAGAG	744
5707	UCAAGUGGGCCUUGGAUCG	318	5707	UCAAGUGGGCCUUGGAUCG	318	5729	CGAUCCAAGGCCACUUA	745

5725	GCUAAGCUGGCUUCUUG	319	5725	GCUAAGCUGGCUUCUUG	319	5747	CAAACAGAGCCAGCUUAGC	746
5743	GAUGCUAUUUUAUGCAAGUU	320	5743	GAUGCUAUUUUAUGCAAGUU	320	5765	AACUUGCAUAAAUAGCAUC	747
5761	UAGGGUCUUAUUAUUAAGG	321	5761	UAGGGUCUUAUUAUUAAGG	321	5783	CCUAAUUAUUAUUAAGCCUA	748
5779	GAUGCGCCUACUUCUAGG	322	5779	GAUGCGCCUACUUCUAGG	322	5801	CCUGAAGAGUAGGCGCAUC	749
5797	GGUCUAAAGCAUUAAGGG	323	5797	GGUCUAAAGCAUUAAGGG	323	5819	CCCACUUGAUUUUAGACC	750
5815	GCCUUGGAUCGUAAGCUG	324	5815	GCCUUGGAUCGUAAGCUG	324	5837	CAGCUUAGCGAUCCAAAGGC	751
5833	GGCUCUGUUUAUGCUAUU	325	5833	GGCUCUGUUUAUGCUAUU	325	5855	AUAGCAUCAAAACAGAGCC	752
5851	UUUAGCAAGUUAGGGUCUA	326	5851	UUUAGCAAGUUAGGGUCUA	326	5873	UAGACCCUAAACUUGCAUAA	753
5869	AUGUAUUUAGGAUGUCG	327	5869	AUGUAUUUAGGAUGUCG	327	5891	GCAGACAUCUUAUUAAGAU	754
5887	CACCUUCUGCAGCCAGUCA	328	5887	CACCUUCUGCAGCCAGUCA	328	5909	UGACUGGCGUCAGAGGUG	755
5905	AGAAAGCUGGAGAGGCAACA	329	5905	AGAAAGCUGGAGAGGCAACA	329	5927	UGUUGCCUUCUCCAGCUUCU	756
5923	AGUGGAUUGCUUCUUC	330	5923	AGUGGAUUGCUUCUUC	330	5945	CAAGAAGCAGCAUCCACU	757
5941	GGGGAGAAAGAUUGCUUC	331	5941	GGGGAGAAAGAUUGCUUC	331	5963	GAAGCAUACUUCUUCUCCC	758
5959	CCUUUUUAUCCAUUAUUA	332	5959	CCUUUUUAUCCAUUAUUA	332	5981	AAAUACAUGGUAUAAAGG	759
5977	UAACUGUAGAACCCUGAGCU	333	5977	UAACUGUAGAACCCUGAGCU	333	5999	AGCUCAGGUUCUACAGUUA	760
5995	UCUAAAGUAACCGAAGAUU	334	5995	UCUAAAGUAACCGAAGAUU	334	6017	CAUUCUUCGGUUAUUAUA	761
6013	GUUAGCCUUCUUCUUAUG	335	6013	GUUAGCCUUCUUCUUAUG	335	6035	CAUAAAGAACAGAGGCAUAC	762
6031	GUGCCACAUCUUAUUA	336	6031	GUGCCACAUCUUAUUA	336	6053	UUAACAAGGAUUGGCGAC	763
6049	AAGGCUUCUUAUUAAGA	337	6049	AAGGCUUCUUAUUAAGA	337	6071	UCUUCUACAGAGAGCCUU	764
6067	AGAUGGACCGUUAUUAAGC	338	6067	AGAUGGACCGUUAUUAAGC	338	6089	GCUGAUAGCGGCUCCAUUCU	765
6085	CACAUUCCUUAUUAAGCCU	339	6085	CACAUUCCUUAUUAAGCCU	339	6107	AGGCUCACUAGGGAUUG	766
6103	UACUGGCUUCUUGCAGCGG	340	6103	UACUGGCUUCUUGCAGCGG	340	6125	CCGCUGCCAGGAGCCAGUA	767
6121	GCUUUUGUGGAAGACUAC	341	6121	GCUUUUGUGGAAGACUAC	341	6143	GUGAGUUCUCCACAAAGC	768
6139	CUAGCCAGAAAGAGAGGAGU	342	6139	CUAGCCAGAAAGAGAGGAGU	342	6161	ACUCCUUCUUCUUGGCUAG	769
6157	UGGGACAGUCCUCCACAC	343	6157	UGGGACAGUCCUCCACAC	343	6179	GGUGGAGAGGACUGUCCCA	770
6175	CAAGAUUAUUAUUAUUA	344	6175	CAAGAUUAUUAUUAUUA	344	6197	UGUUUGGAUUAUUAUUAU	771
6193	AAAAGCAGGCUAGAGCCAG	345	6193	AAAAGCAGGCUAGAGCCAG	345	6215	CUGGCUUAGCCUUCUUCU	772
6211	GAAGAGAGGACAAUUAU	346	6211	GAAGAGAGGACAAUUAU	346	6233	AAAGAUUUGUCCUUCUUC	773
6229	UGUUUGUCCUUCUUAU	347	6229	UGUUUGUCCUUCUUAU	347	6251	GUAAAGAGAGGAAACAACA	774
6247	CACAUACGCAAAACCCUG	348	6247	CACAUACGCAAAACCCUG	348	6269	CAGGUGUUGCGUAUUG	775
6265	GUGACAGCUGGCAUUAU	349	6265	GUGACAGCUGGCAUUAU	349	6287	UAAAUIUCCAGCUGUCCAC	776
6283	AUAAUACAGGUUAUUAU	350	6283	AUAAUACAGGUUAUUAU	350	6305	UUCAGAUUACCUUUAUUAU	777
6301	AGAGGUUAUUAUUAUUA	351	6301	AGAGGUUAUUAUUAUUA	351	6323	UUUCUGAGUUUAUUAUUAU	778
6319	AAAAGAGACCUUUAUUA	352	6319	AAAAGAGACCUUUAUUA	352	6341	UUGACUGAGGUUUAUUAU	779
6337	AUUCUCUUAUUAUUAU	353	6337	AUUCUCUUAUUAUUAU	353	6359	AAAAAAAGUUAUUAUUA	780
6355	UUUUUUUCCAAUUAUUA	354	6355	UUUUUUUCCAAUUAUUA	354	6377	UAUCUGAUUUGGAAAAAA	781

6373	AAUAGCCCAGCAAAUJAGUG	355	6373	AAUAGCCCAGCAAAUJAGUG	355	6395	CACUUAUUUGCGGCUAUU	782
6391	GAUAAACAAUUAACCCUUA	356	6391	GAUAAACAAUUAACCCUUA	356	6413	UAAGGUUUUAUUUGUUAUC	783
6409	AGCUGUUCAUGUCUUGAUU	357	6409	AGCUGUUCAUGUCUUGAUU	357	6431	AUACAAGACAGAACAGCU	784
6427	UUCAAUAAUUAUUUCUUA	358	6427	UUCAAUAAUUAUUUCUUA	358	6449	UUAAGAAUUAUUAUUGAA	785
6445	AUAAUUAAGAGACCAUAAU	359	6445	AUAAUUAAGAGACCAUAAU	359	6467	AUUAUGGUCUCUUAUUGAU	786
6463	UAAAUACUCCUUAUUAAGA	360	6463	UAAAUACUCCUUAUUAAGA	360	6485	UCUUGAAAAGGAGUUAUUA	787
6481	AGAAAACCAAAACCAUJAG	361	6481	AGAAAACCAAAACCAUJAG	361	6503	CUAAUGGUUUUGCUUUUCU	788
6499	GAUUUUAUACUCAGCUCCU	362	6499	GAUUUUAUACUCAGCUCCU	362	6521	AGGAGCUGAGUAACAUIUC	789
6517	UUCAAACUCAGGUUUUGUAG	363	6517	UUCAAACUCAGGUUUUGUAG	363	6539	CUACAACCCUGAGUUGAA	790
6535	GCAUACAUAGAGUCCAUCCA	364	6535	GCAUACAUAGAGUCCAUCCA	364	6557	UGGAUGGACUAGUUAUGC	791
6553	AUCAGUCAAGAAUUGGUUC	365	6553	AUCAGUCAAGAAUUGGUUC	365	6575	GAACCAUUCUUAUGACUGAU	792
6571	CCAUUCGGAGUCUUAUUGU	366	6571	CCAUUCGGAGUCUUAUUGU	366	6593	ACAUUAAGACUCCAGAUUG	793
6589	UAGAAAGAAAUAUGGAGAC	367	6589	UAGAAAGAAAUAUGGAGAC	367	6611	GUCUCCAUUUUUUCUUUCUA	794
6607	CUUGUAUAUAGAGCUAGU	368	6607	CUUGUAUAUAGAGCUAGU	368	6629	ACUAGCUCAUUAUUACAAG	795
6625	UUAACAAGUGCUUUGUICAU	369	6625	UUAACAAGUGCUUUGUICAU	369	6647	AUGAACAGCACUUAUUGUAA	796
6643	UUAUAAUAGCACUGAAAAU	370	6643	UUAUAAUAGCACUGAAAAU	370	6665	AUUUUCAGGUCUUAUUUAA	797
6661	UUGAAACAUGAAUUAACUG	371	6661	UUGAAACAUGAAUUAACUG	371	6683	CAGUUAUUAUGUUAUCAA	798
6679	GAUAAUUAUCCAUAUUAU	372	6679	GAUAAUUAUCCAUAUUAU	372	6701	AAUUGAUUGGAUUAUUAUC	799
6697	UGCCAUUAUUGACAAAAU	373	6697	UGCCAUUAUUGACAAAAU	373	6719	AUUUUUGUCAUAAUUGGCA	800
6715	UGGUUGGCACUACAAAGA	374	6715	UGGUUGGCACUACAAAGA	374	6737	UCUUUUGUUGUGCCCAACCA	801
6733	AACGAGACUUCUUAUCAG	375	6733	AACGAGACUUCUUAUCAG	375	6755	CUGAAAGGAAGUGCUCUGU	802
6751	GAGUUAUCUGAGAUUAUGUA	376	6751	GAGUUAUCUGAGAUUAUGUA	376	6773	UACAUUAUCUCAGAAACUC	803
6769	ACGUGAACAGUCUGGGUG	377	6769	ACGUGAACAGUCUGGGUG	377	6791	CACCCAGACUGUUCACGU	804
6787	GGAAUGGGGCUAGAAACCAU	378	6787	GGAAUGGGGCUAGAAACCAU	378	6809	AUGGUUUCAGCCCCAUUCC	805
6805	UGUGCAAGUCUGUGUCUUG	379	6805	UGUGCAAGUCUGUGUCUUG	379	6827	CAAGACACAGACUUGCACA	806
6823	GUACAGUCCAAGAGUGACA	380	6823	GUACAGUCCAAGAGUGACA	380	6845	UGUCACUUCUUGGACUGAC	807
6841	ACCGAGAUUAUUAUUUAG	381	6841	ACCGAGAUUAUUAUUUAG	381	6863	CUAAAUAUACAUCUCUGU	808
6859	GGGACCCGUGCCUUGUUAUC	382	6859	GGGACCCGUGCCUUGUUAUC	382	6881	GAACAAGGCACCGGUGCC	809
6877	CCUAGCCCACAGAAUGCA	383	6877	CCUAGCCCACAGAAUGCA	383	6899	UGCAUUCUUGUGGGCUAGG	810
6895	AAACAUCAAACAGAUACUC	384	6895	AAACAUCAAACAGAUACUC	384	6917	GAGUAUCUGUUAUGUUAU	811
6913	CGUAGCCUUAUUAUUAU	385	6913	CGUAGCCUUAUUAUUAU	385	6935	AAUUAAAUAUGAGGCUAGCG	812
6931	UGAUUAAAGGAGGAGUGCA	386	6931	UGAUUAAAGGAGGAGUGCA	386	6953	UGCACUCCUCCUUUAUCA	813
6949	AUCUUUGGCCGACAGUGGU	387	6949	AUCUUUGGCCGACAGUGGU	387	6971	ACCACUGUGGCCAAAGAU	814
6967	UGUAACUGUGUGUGUGUGU	388	6967	UGUAACUGUGUGUGUGUGU	388	6989	ACACACACACACAGUUA	815
6985	UGUGUGUGUGUGUGUGUGU	389	6985	UGUGUGUGUGUGUGUGUGU	389	7007	ACACACACACACACACACA	816
7003	UGUGUGUGUGUGGGUGUGG	390	7003	UGUGUGUGUGUGGGUGUGG	390	7025	CCACACCCACACACACACA	817

7021	GGUGUAUGUGUUUUUGUG	391	7021	GGUGUAUGUGUUUUUGUG	391	7043	CACAAAACACACAUACACC	818
7039	GCAUAAUAUUUAAGGAAA	392	7039	GCAUAAUAUUUAAGGAAA	392	7061	UUUCCUUUUUUUAAGUAGC	819
7057	ACUGGAAUUUUUAAGUUAU	393	7057	ACUGGAAUUUUUAAGUUAU	393	7079	GUACUUUUUUUUUUUCCAGU	820
7075	CUUUUAUACAAACCAAGAA	394	7075	CUUUUAUACAAACCAAGAA	394	7097	UUUUGGUUUUGUAUAAAAG	821
7093	AUAUAUGCUACAGAUUAUA	395	7093	AUAUAUGCUACAGAUUAUA	395	7115	UUUAUUCUGUAGCAUAUAU	822
7111	AGACAGACAUUGUUUGGUC	396	7111	AGACAGACAUUGUUUGGUC	396	7133	GACCAACCAUGUCUGUCU	823
7129	CCUAUAUUUCUAGCUAUA	397	7129	CCUAUAUUUCUAGCUAUA	397	7151	UCAUGACUAGAAUAUAGG	824
7147	AUGAAUGUAUUUUUAUAU	398	7147	AUGAAUGUAUUUUUAUAU	398	7169	GUUAUUAUAUGAUAUAU	825
7165	CCAUCUUAUAUAUAUAU	399	7165	CCAUCUUAUAUAUAUAU	399	7187	GUUAUUAUAUGAUAUAU	826
7183	CUUAAAUAUUUUUUUAUAU	400	7183	CUUAAAUAUUUUUUUAUAU	400	7205	AUUAAGAAUAUUUUUAAG	827
7201	UUUGGAAUUUGUAUUCGUAC	401	7201	UUUGGAAUUUGUAUUCGUAC	401	7223	GUACGAUUAUUAUCCCAA	828
7219	CCAACUUAUUUAUAUAUAU	402	7219	CCAACUUAUUUAUAUAUAU	402	7241	AGUUUAUCAUUUAAGUUGG	829
7237	UUGGCAACUGCUUUUAUAU	403	7237	UUGGCAACUGCUUUUAUAU	403	7259	ACAUAUAAGCAGUUUGCAA	830
7255	UUCUGUCUCCUUGCAUAAA	404	7255	UUCUGUCUCCUUGCAUAAA	404	7277	UUUAUGGAAGGAGACAGAA	831
7273	AUUUUUCAAUAUAUAUAU	405	7273	AUUUUUCAAUAUAUAUAU	405	7295	AUAUAUAUUUUUGAAAAAU	832
7291	UCAACAAAGAAAGCUCU	406	7291	UCAACAAAGAAAGCUCU	406	7313	AGAGCUUUUUUUUUUUUGA	833
7309	UUUUUUUCCUUAUAUAUA	407	7309	UUUUUUUCCUUAUAUAUA	407	7331	UUUAUUUUUAAGAAAAAAA	834
7327	ACUCAAUUUUAUCCUUGUU	408	7327	ACUCAAUUUUAUCCUUGUU	408	7349	AACAAGGAUAUUUUUUGAGU	835
7345	UUAGAGCAGAGAAAAUAUA	409	7345	UUAGAGCAGAGAAAAUAUA	409	7367	UAAUUUUUCUCUGCUCUAA	836
7363	AAGAAAAACUUUGAAUUGG	410	7363	AAGAAAAACUUUGAAUUGG	410	7385	CCAUUUAAGUUUUUUUUUU	837
7381	GUUCUCAAUAUUUGCUAAA	411	7381	GUUCUCAAUAUUUGCUAAA	411	7403	UUUAAGCAUUUUUUUGAGAC	838
7399	AUAUUUCAAUUGGAAUAUA	412	7399	AUAUUUCAAUUGGAAUAUA	412	7421	AGUUUUCCAUUGAAAAUAU	839
7417	UAAUUGUAUUUAAGCUAUA	413	7417	UAAUUGUAUUUAAGCUAUA	413	7439	UCAGCUAAACUAACAUUAUA	840
7435	AUUGUAUUGGGUUUUUGGAA	414	7435	AUUGUAUUGGGUUUUUGGAA	414	7457	UUCGAAAAACCCCAUACAAU	841
7453	ACCUUUCACUUUUUUUGUUUG	415	7453	ACCUUUCACUUUUUUUGUUUG	415	7475	CAACAAAAAGUGAAAGGU	842
7471	GUUUUACCUAAUUUACAAC	416	7471	GUUUUACCUAAUUUACAAC	416	7493	GUUGUGAAUAUAGGUAAAC	843
7489	CUGUGUAUUUUGCCAAUAUA	417	7489	CUGUGUAUUUUGCCAAUAUA	417	7511	UUUAUUGGCAUUUUACACAG	844
7507	AUUCCUGUCCAUUGAAAAUG	418	7507	AUUCCUGUCCAUUGAAAAUG	418	7529	CAUUUUAUGGACAGGAUAU	845
7525	GCAAAUUUAUCCAGUGUAGA	419	7525	GCAAAUUUAUCCAGUGUAGA	419	7547	UCUACACUGGGAUAAUUUGC	846
7543	AUAUAUUUAGCCAUACCC	420	7543	AUAUAUUUAGCCAUACCC	420	7565	GGUGAUGGCUCAAAUAUAU	847
7561	CUAUGGAUAUUUGGUAGUU	421	7561	CUAUGGAUAUUUGGUAGUU	421	7583	AACUAGCCAAUAUCCAUAG	848
7579	UUUGCCUUUAUAUAAGCAA	422	7579	UUUGCCUUUAUAUAAGCAA	422	7601	UUUGCUUUAUAUAAAGGCAA	849
7597	AUUAUUAUCCAGCCUGAAUG	423	7597	AUUAUUAUCCAGCCUGAAUG	423	7619	CAUUCAGGCUGAAUUAUAU	850
7615	GUCUGCCUAUAUAUUCUCU	424	7615	GUCUGCCUAUAUAUUCUCU	424	7637	AGAGAAUAUAUAGGACAG	851
7633	UGCUCUUUGUAUUUCUCCUU	425	7633	UGCUCUUUGUAUUUCUCCUU	425	7655	AAGGAGAAUAUAAAGAGCA	852
7651	UUGAACCCGUUAAACAUUC	426	7651	UUGAACCCGUUAAACAUUC	426	7673	GAUGUUUUUAACGGGUUCAA	853

7662	AAAACAUCUUGGCACUC	427	7682	AAAACAUCUUGGCACUC	427	7684	GAGUGCCACAGGAUGUUUU	854
------	-------------------	-----	------	-------------------	-----	------	---------------------	-----

VEGFR2 gi|11321596|ref|NM\_002253.1

Pos	Target Sequence	Seq ID	UPos	Upper seq	Seq ID	LPpos	Lower seq	Seq ID
1	ACUGAGUCCCGGACCCCG	855	1	ACUGAGUCCCGGACCCCG	855	23	CGGGGUCCCGGACUCAGU	1179
19	GGGAGAGCGGUCAGUGUGU	856	19	GGGAGAGCGGUCAGUGUGU	856	41	ACACACUGACCGCUCUCCC	1180
37	UGGUCGUCGCUUUCUCU	857	37	UGGUCGUCGCUUUCUCU	857	59	AGAGGAAACGCAGGACCA	1181
55	UGCCUGCGCCGGCAUCAC	858	55	UGCCUGCGCCGGCAUCAC	858	77	GUGAUGCCCGCGCGCAAG	1182
73	CUUGCGCGCCGAGAAAGU	859	73	CUUGCGCGCCGAGAAAGU	859	95	ACUUUCUGCGCGCGCAAG	1183
91	UCCGUCUGGAGCCUGGAU	860	91	UCCGUCUGGAGCCUGGAU	860	113	AUCCAGGCGCGCGCAAG	1184
109	UAUCCUUCUACCGGCAC	861	109	UAUCCUUCUACCGGCAC	861	131	GUGCCGUGGAGAGGAUA	1185
127	CCCGCAGACGCCUUCAG	862	127	CCCGCAGACGCCUUCAG	862	149	CUGCCGUGGAGAGGAUA	1186
145	GGCGCGGUCGGCGCCCGG	863	145	GGCGCGGUCGGCGCCCGG	863	167	CCGGCGCGGACCGCGGC	1187
163	GGCUCUCCUAGCCUUGCG	864	163	GGCUCUCCUAGCCUUGCG	864	185	CGCACAGGCGUAGGAGCC	1188
181	GCUCACUCUGCCGCGCUG	865	181	GCUCACUCUGCCGCGCUG	865	203	CAGCGCAGGACGAGGAGC	1189
199	GCGGGUGCGCGAGUUC	866	199	GCGGGUGCGCGAGUUC	866	221	GGAACUCGCGGACCCCGC	1190
217	CACUCCGCGCCUUCUUC	867	217	CACUCCGCGCCUUCUUC	867	239	AGAAGGAGCGCGGAGGUG	1191
235	UCUAGACAGCGCGGAG	868	235	UCUAGACAGCGCGGAG	868	257	CUCCAGCGCGGAGGAGG	1192
253	GAAAGAACCGCGCCCGAG	869	253	GAAAGAACCGCGCCCGAG	869	275	CUCCAGCGCGGAGGAGG	1193
271	GUUCUGGCAUUCGCCCCG	870	271	GUUCUGGCAUUCGCCCCG	870	293	CGGGCGAAUUGCCCAAGC	1194
289	GGCUCGAGGUGCAGGAUC	871	289	GGCUCGAGGUGCAGGAUC	871	311	GCAUCCUGCACCUCGAGCC	1195
307	CAGAGCAAGGUGCUGCUG	872	307	CAGAGCAAGGUGCUGCUG	872	329	CCAGCAGCACCUCGAGCC	1196
325	GCCGUCGCGCUGGCGUCU	873	325	GCCGUCGCGCUGGCGUCU	873	347	AGAGCCACAGGCGGACGCG	1197
343	UGCGUGGAGACCGGCGCG	874	343	UGCGUGGAGACCGGCGCG	874	365	CGGCCGCGGUCUCCACGCA	1198
361	GCCUCUGUGGGUUGCCUA	875	361	GCCUCUGUGGGUUGCCUA	875	383	UAGGCAAAACCCACAGAGC	1199
379	AGUGUUCUCUUGAUCUGC	876	379	AGUGUUCUCUUGAUCUGC	876	401	GCAGAUCAAGAGAAACACU	1200
397	CCCAGGCUCAGCAUACAA	877	397	CCCAGGCUCAGCAUACAA	877	419	UUUGUAUGCUGAGCCUGGG	1201
415	AAAGACAUACUACAAUUA	878	415	AAAGACAUACUACAAUUA	878	437	UAAUUGUAAGUAUGUUAU	1202
433	AAGGCUAAUACACUCUUC	879	433	AAGGCUAAUACACUCUUC	879	455	GAAGAGUUGUAUAGCCUU	1203
451	CAAAUUAUUGGAGGCGAC	880	451	CAAAUUAUUGGAGGCGAC	880	473	GUCCCCUGCAAGUAUUAUG	1204
469	CAGAGGACUUGGAGGCGC	881	469	CAGAGGACUUGGAGGCGC	881	491	GCCAGUCCAAAGUCCUUCUG	1205
487	CUUUGGCCCCAAUUAUCAGA	882	487	CUUUGGCCCCAAUUAUCAGA	882	509	UCUGAUUAUUGGGCCAAAG	1206
505	AGUGGCAGUGAGCAAAAGG	883	505	AGUGGCAGUGAGCAAAAGG	883	527	CCCUUGCUCACUGCCACU	1207
523	GUGGAGGUGACUGAGUGCA	884	523	GUGGAGGUGACUGAGUGCA	884	545	UGCACUCAGUACCCUCCAC	1208



541	AGCGAUGGCCUUCUGUA	885	541	AGCGAUGGCCUUCUGUA	885	563	UACAGAAGAGGCCAUCGCU	1208
559	AAGACACUCACAAUCCAA	886	559	AAGACACUCACAAUCCAA	886	581	UUGGAUUUGAGUGUCUU	1210
577	AAAGUGUCGGAAUAGACA	887	577	AAAGUGUCGGAAUAGACA	887	589	UGCAUUUCCGAGUCUUU	1211
595	ACUGAGCCUACAAGUCU	888	595	ACUGAGCCUACAAGUCU	888	617	AGCAUUUGUAGGCCUCCAGU	1212
613	UUCUACCGGAAACUACU	889	613	UUCUACCGGAAACUACU	889	635	AGUCAGUUIUCCCGGUAGAA	1213
631	UUGGCCUUGGUCAUUUAG	890	631	UUGGCCUUGGUCAUUUAG	890	653	CAUAAUAGACCGAGGCCAA	1214
649	GUCUAGUUCUACAAGAUACA	891	649	GUCUAGUUCUACAAGAUACA	891	671	UGUAUUCUUGAAACAUAAGAC	1215
667	AGAUCUCCAUUUUUAUUGCUU	892	667	AGAUCUCCAUUUUUAUUGCUU	892	689	AAGCAUAAUUGGAGAUUCU	1216
685	UCUGUUAUGACCAACAUG	893	685	UCUGUUAUGACCAACAUG	893	707	CAUGUUGGUCACUAAACAGA	1217
703	GGAGUCGUGUACAUAUACUG	894	703	GGAGUCGUGUACAUAUACUG	894	725	CAGUAUUGUACACGACUCC	1218
721	GAGAACAAAACAAACUG	895	721	GAGAACAAAACAAACUG	895	743	CAGUUUUGUUIUUGUUCUC	1219
739	GUGGUGAUUCCAUGUCUCG	896	739	GUGGUGAUUCCAUGUCUCG	896	761	CGAGACAUUGGAUACCCAC	1220
757	GGGUCCAUAUCAAUUCUCA	897	757	GGGUCCAUAUCAAUUCUCA	897	779	UGAGAUUUGAAUUGGACCC	1221
775	AACGUGUCACUUGUGCAA	898	775	AACGUGUCACUUGUGCAA	898	797	UUGCACAAAGUGACACGUU	1222
793	AGAUACCCAGAAAAGAGAU	899	793	AGAUACCCAGAAAAGAGAU	899	815	AUCUCUUUUCUGGGUUAUCU	1223
811	UUUGUUCUUGAUGGUAAACA	900	811	UUUGUUCUUGAUGGUAAACA	900	833	UGUACCAUCAGGAAACAAA	1224
829	AGAAUUUCCUGGGACAGCA	901	829	AGAAUUUCCUGGGACAGCA	901	851	UGUUGCCAGGAAAUUUCU	1225
847	AAGAAGGCUUUAUUAUUC	902	847	AAGAAGGCUUUAUUAUUC	902	869	GAUAGUAAGGCCUUCUUU	1226
865	CCCAGCUACAUGACAGCU	903	865	CCCAGCUACAUGACAGCU	903	887	AGCUGAUCUAGUAGCUGGG	1227
883	UAUGCUGGCAUGGUCUUCU	904	883	UAUGCUGGCAUGGUCUUCU	904	905	AGAAGACCAUGCCAGCAUA	1228
901	UGUGAAGCAAAAUAUUAUG	905	901	UGUGAAGCAAAAUAUUAUG	905	923	CAUUAUUUUUUGCUUUCACA	1229
919	GAUGAAAGUUAACAGUCUA	906	919	GAUGAAAGUUAACAGUCUA	906	941	UAGACUGGUAACUUAUUC	1230
937	AUUAUGUACAUAUAGUUGUCG	907	937	AUUAUGUACAUAUAGUUGUCG	907	959	CGACAACUAUUAUUAUUAU	1231
955	GUUGUAGGGUUAUAGGAUUU	908	955	GUUGUAGGGUUAUAGGAUUU	908	977	AAAUCCUAUACCCUACAAC	1232
973	UAUGAUUGGUUUCUGAGUC	909	973	UAUGAUUGGUUUCUGAGUC	909	985	GACUCAGAACCACAUCAUA	1233
991	CCGUCUCAUGGAUUAUUAAC	910	991	CCGUCUCAUGGAUUAUUAAC	910	1013	GUUCAAUUCCAUAGAGACGG	1234
1009	CUAUCUGUUGGAGAAAAGC	911	1009	CUAUCUGUUGGAGAAAAGC	911	1031	GCUUUUUCCCAACAGAUAG	1235
1027	CUUGUCUUAUUUUAUUAACAG	912	1027	CUUGUCUUAUUUUAUUAACAG	912	1049	CUUUAUUAUUUAAGACAAG	1236
1045	GCAAGAACUGAACUAUUAUUG	913	1045	GCAAGAACUGAACUAUUAUUG	913	1067	CAUUUAGUUCAGUUCUUGC	1237
1063	GUGGGAUUUGACUUAACU	914	1063	GUGGGAUUUGACUUAACU	914	1085	AGUUAGAGUCAAUCCCCAC	1238
1081	UGGGAUAACCUUUCUUGCA	915	1081	UGGGAUAACCUUUCUUGCA	915	1103	UCGAAGAGGGUUAUUCCCA	1239
1099	AAGCAUCAGCAUAGAAAC	916	1099	AAGCAUCAGCAUAGAAAC	916	1121	GUUUCUUAUUGCUGAUGCUU	1240
1117	CUUGUAACCCGAGACCUAA	917	1117	CUUGUAACCCGAGACCUAA	917	1139	UUAGGUCUCGGUUUUAACAAG	1241
1135	AAACCCAGUCUGGGAGUG	918	1135	AAACCCAGUCUGGGAGUG	918	1157	CACUCCAGACUGGGUUAU	1242
1153	GAGAUAGAAGAAUUUUAUGA	919	1153	GAGAUAGAAGAAUUUUAUGA	919	1175	UCAAAAUIUUCUUAUCUUC	1243
1171	AGCACCUUAACUAUAGAUUG	920	1171	AGCACCUUAACUAUAGAUUG	920	1193	CAUCUAUAGUUAAGGUGCU	1244



1189	GGUGUAACCCGGAGUGACC	921	1189	GGUGUAACCCGGAGUGACC	921	1211	GGUCACUCGGGUGUACACC	1245
1207	CAAGGAUUGUACACCUUG	922	1207	CAAGGAUUGUACACCUUG	922	1229	CACAGGUGUACAAUCCUUG	1246
1225	GCAGCAUCCAGUGGGCUGA	923	1225	GCAGCAUCCAGUGGGCUGA	923	1247	UCAGCCCACUGGAUGCUGC	1247
1243	AUGACCAAGAAAGAACAGCA	924	1243	AUGACCAAGAAAGAACAGCA	924	1265	UGCUGUUCUUCUUGGUGAU	1248
1261	ACAUUUGUCAGGUUCCUAG	925	1261	ACAUUUGUCAGGUUCCUAG	925	1283	CAUGGACCCUGACAAAUGU	1249
1279	GAUAAACUUUUGUUGCUU	926	1279	GAUAAACUUUUGUUGCUU	926	1301	AAGCAACAAAAGGUUUUUC	1250
1297	UUUGGAUUGGCAUGGAU	927	1297	UUUGGAUUGGCAUGGAU	927	1319	AUUCUAGCCACUUCCAA	1251
1315	UCUCUGGUGAAGCCACGG	928	1315	UCUCUGGUGAAGCCACGG	928	1337	CCGUGGUUCCACCAGAGA	1252
1333	GUGGGGAGCGUGUCAGAA	929	1333	GUGGGGAGCGUGUGAGAA	929	1355	UUUCGACAGCUCUCCCCAC	1253
1351	AUCCUGCGAAGUACCUUG	930	1351	AUCCUGCGAAGUACCUUG	930	1373	CAAGGUACUUCGACGGGAU	1254
1369	GGUACCCACCCGAGAAA	931	1369	GGUACCCACCCGAGAAA	931	1391	UUUCUGGGGUGGGUAAOC	1255
1387	AUAAAUGGUUAAAAUUG	932	1387	AUAAAUGGUUAAAAUUG	932	1409	CAUUUUUUAUACCAUUUAU	1256
1405	GGAUACCCUUGAGUCCA	933	1405	GGAUACCCUUGAGUCCA	933	1427	UGGACUCAAGGGGUUUC	1257
1423	AUACACAAUUAAGCGG	934	1423	AUACACAAUUAAGCGG	934	1445	CCGCUUUAUUGUGUGAUU	1258
1441	GGGAUGUACUGACGAUUA	935	1441	GGGAUGUACUGACGAUUA	935	1463	UAUUCUGUACUACUAGCCC	1259
1459	AUGGAAGUGAGUAAAGAG	936	1459	AUGGAAGUGAGUAAAGAG	936	1481	CUCUUUCAGUCACUUCCAU	1260
1477	GACACAGGAUUAUACACUG	937	1477	GACACAGGAUUAUACACUG	937	1499	CAGUGAAUUUCCUUGUGUC	1261
1495	GUCAUCCUUAACCAUCCCA	938	1495	GUCAUCCUUAACCAUCCCA	938	1517	UGGGAUUGGUAAGGAUGAC	1262
1513	AUUUCAAAGGAGAGCAGA	939	1513	AUUUCAAAGGAGAGCAGA	939	1535	UCUGCUUCUCCUUUGAAU	1263
1531	AGCCAUGUGUUCUUCUGG	940	1531	AGCCAUGUGUUCUUCUGG	940	1553	CCAGAGAGACCACAUIGGCU	1264
1549	GUUGUGUAGUCCCAUCC	941	1549	GUUGUGUAGUCCCAUCC	941	1571	GGGUGGGGACAUACACAAC	1265
1567	CAGAUUGGUGAGAAUUCUC	942	1567	CAGAUUGGUGAGAAUUCUC	942	1589	GAGAUUUCUCACCAUUCUG	1266
1585	CUAAUCUCUCCUGUGGAU	943	1585	CUAAUCUCUCCUGUGGAU	943	1607	AUUCACAGGAGAGAUUAG	1267
1603	UCCUACCAUACGGCAGCA	944	1603	UCCUACCAUACGGCAGCA	944	1625	UGGUGCGGUACUGGUAGGA	1268
1621	ACUCAAACGUGACAUUGUA	945	1621	ACUCAAACGUGACAUUGUA	945	1643	UACAUUCAGCGUUUGAGU	1269
1639	ACGGUCUAUCCAUUCCUC	946	1639	ACGGUCUAUCCAUUCCUC	946	1661	GAGGAUUGGCAUAGACCGU	1270
1657	CCCCGCAUCACAUCCACU	947	1657	CCCCGCAUCACAUCCACU	947	1679	AGUGGAUUGAUUGCGGGG	1271
1675	UGGUUUGGCAUUGGAGG	948	1675	UGGUUUGGCAUUGGAGG	948	1697	CCUCCAACUGCCAAUACCA	1272
1693	GAAGAGUGCGCCACGAGC	949	1693	GAAGAGUGCGCCACGAGC	949	1715	GCUCGUUGGGCGCACUUC	1273
1711	CCAGCCCAAGCUGUCUCAG	950	1711	CCAGCCCAAGCUGUCUCAG	950	1733	CUGAGACAGCUUGGCGGG	1274
1729	GUGACAAACCCAUACCCU	951	1729	GUGACAAACCCAUACCCU	951	1751	AAGGUAUGGGGUUUGUCAC	1275
1747	UGUGAAGAUUGGAGAGUG	952	1747	UGUGAAGAUUGGAGAGUG	952	1769	CACUUCUCCAUUCUUCACA	1276
1765	GUGGAGGACUUCAGGGAG	953	1765	GUGGAGGACUUCAGGGAG	953	1787	CUCCUGGAGUCCUCCAG	1277
1783	GGAAUAAAUUAGAGUUA	954	1783	GGAAUAAAUUAGAGUUA	954	1805	UAACUCAAUUUUUUUCC	1278
1801	AAUAAAUUUAUUGCUC	955	1801	AAUAAAUUUAUUGCUC	955	1823	GAGCAAUUUGAUUUUUUUAU	1279
1819	CUAAUUGAAGGAAAAACA	956	1819	CUAAUUGAAGGAAAAACA	956	1841	UGUUUUUUUCCUCAAUUAUAG	1280

1837	AAAACUGUAAGUACCCUUG	957	1837	AAAACUGUAAGUACCCUUG	957	1859	CAAGGGUACUACAGUUUU	1281
1855	GUUAUCCAAGCGGCAAAUG	958	1855	GUUAUCCAAGCGGCAAAUG	958	1877	CAUUGCCGCUUGGAUAC	1282
1873	GUGUCAGCUUUGUACAAAU	959	1873	GUGUCAGCUUUGUACAAAU	959	1895	AUUUGUACAAAGCGUACAC	1283
1891	UGUGAAGCGGUCAACAAAG	960	1891	UGUGAAGCGGUCAACAAAG	960	1913	CUUUGUUGACCGCUUCACA	1284
1909	GUCGGGAGAGGAGAGAGGG	961	1909	GUCGGGAGAGGAGAGAGGG	961	1931	CCCUUCUCCUCCUCCGAC	1285
1927	GUGAUUCUUCUCCACGUGA	962	1927	GUGAUUCUUCUCCACGUGA	962	1949	UCACGUGGAAGGAGAUAC	1286
1945	ACAGGGGUCUUGAAUUA	963	1945	ACAGGGGUCUUGAAUUA	963	1967	UAUUUCAGGAGCCUUGU	1287
1963	ACUUUGCAACCUGACAUUC	964	1963	ACUUUGCAACCUGACAUUC	964	1985	GCAUGUCAGGUUGCAAAGU	1288
1981	CAGCCACUUGAGCAGGAGA	965	1981	CAGCCACUUGAGCAGGAGA	965	2003	UCUCCUGCUAGUGGGCUG	1289
1999	AGCGUGUCUUGUGGUGCA	966	1999	AGCGUGUCUUGUGGUGCA	966	2021	UGCACCACAAAGACACGU	1290
2017	ACUGCAGACAGAUUCUACGU	967	2017	ACUGCAGACAGAUUCUACGU	967	2039	ACGUAGAUUCUGUCGACGU	1291
2035	UUUGAGAACCUCAUUGGU	968	2035	UUUGAGAACCUCAUUGGU	968	2057	ACCAUGUGAGGUUCUCAAA	1292
2053	UACAAGCUUGGCCCCACAGC	969	2053	UACAAGCUUGGCCCCACAGC	969	2075	GCUGUGGGCCAAAGCUUGUA	1293
2071	CCUCUGCCAAUCCAUGUGG	970	2071	CCUCUGCCAAUCCAUGUGG	970	2093	CCCAUGGUUUGGCAGAGG	1294
2089	GGAGAGUUGCCACACACUUG	971	2089	GGAGAGUUGCCACACACUUG	971	2111	CAGGUGGGGCAACUCC	1295
2107	GUUUGCAAGAAUUGGAUA	972	2107	GUUUGCAAGAAUUGGAUA	972	2129	UAUCCAAGUUCUUGCAAAC	1296
2125	ACUCUUGGAAAUUGAAUG	973	2125	ACUCUUGGAAAUUGAAUG	973	2147	CAUUCAAUUCCAAAGAGU	1297
2143	GCCACCAUGUUCUUAUA	974	2143	GCCACCAUGUUCUUAUA	974	2165	UAUUGAGAGAACAUUGGUGC	1298
2161	AGCACAAUAGACAUUUUGA	975	2161	AGCACAAUAGACAUUUUGA	975	2183	UCAAAUUGUUAUUUGUCU	1299
2179	AUCAUGGAGCUUAAGAAUG	976	2179	AUCAUGGAGCUUAAGAAUG	976	2201	CAUUCUUAAGCUCCAUGAU	1300
2197	GCAUCCUUGCAGGACCAAG	977	2197	GCAUCCUUGCAGGACCAAG	977	2219	CUUUGUCCUGCAAGGAGC	1301
2215	GGAGACUUAUGUCUCCUUG	978	2215	GGAGACUUAUGUCUCCUUG	978	2237	CAAGGCAGACAUAGUCUCC	1302
2233	GCUCAAGACAGGAAGACCA	979	2233	GCUCAAGACAGGAAGACCA	979	2255	UGGUCUCCUGUCUUGAGC	1303
2251	AAGAAAGACAUUGCGUGG	980	2251	AAGAAAGACAUUGCGUGG	980	2273	CCACGCAUUGUCUUAUUCU	1304
2269	GUCAGGCAGCUACAGUCC	981	2269	GUCAGGCAGCUACAGUCC	981	2291	GGACUGUGAGCUGCCUGAC	1305
2287	CUAGAGCGUGUGGCACCCA	982	2287	CUAGAGCGUGUGGCACCCA	982	2309	UGGGUCCACACGCUCUAG	1306
2305	ACGAUCACAGGAACCUUG	983	2305	ACGAUCACAGGAACCUUG	983	2327	CCAGGUUUCUUGUAGUCCU	1307
2323	GAGAAUCAGACGACAAAGUA	984	2323	GAGAAUCAGACGACAAAGUA	984	2345	UACUUGUCUGUUAUUCUC	1308
2341	AUUGGGAAAGCAUCGAAAG	985	2341	AUUGGGAAAGCAUCGAAAG	985	2363	CUUCGAUGCUUUCUCCAAU	1309
2359	GUCUCAUGCACGGCAUCUG	986	2359	GUCUCAUGCACGGCAUCUG	986	2381	CAGAUGCCGUGCAUGAGAC	1310
2377	GGGAUCCCCUCCACAGA	987	2377	GGGAUCCCCUCCACAGA	987	2399	UCUGUGGAGGGGGAUUCUCC	1311
2395	AUCAUGUGGUUUAAGUA	988	2395	AUCAUGUGGUUUAAGUA	988	2417	UAUCUUUAACCCACAUAGU	1312
2413	AUUGAGACCCUUGUAGAG	989	2413	AUUGAGACCCUUGUAGAG	989	2435	CUUCUACAAAGGUCUCAU	1313
2431	GACUCAGGCAUUGUAUGA	990	2431	GACUCAGGCAUUGUAUGA	990	2453	UCAUUAUAGCCUGAGUC	1314
2449	AAGGAUGGGAACCGGAACC	991	2449	AAGGAUGGGAACCGGAACC	991	2471	GGUCCGCUUCCCAUCCU	1315
2467	CUCACUUAUCCGCAGAGUGA	992	2467	CUCACUUAUCCGCAGAGUGA	992	2489	UCACUCUGCGGAUAGUGAG	1316

2485	AGGAAGGAGGACGAAGGCC	993	2485	AGGAAGGAGGACGAAGGCC	993	2507	GGCCUUCGUCUCCUUCU	1317
2503	CUCUACACCUGCCAGGCAU	994	2503	CUCUACACCUGCCAGGCAU	994	2525	AUCCUUCGUCUCCUUCU	1318
2521	UGCAGUGUUCUUGGCUUG	995	2521	UGCAGUGUUCUUGGCUUG	995	2543	CACAGCCAAGAACACUGCA	1319
2539	GCAAAAGUGGAGGCAUUCU	996	2539	GCAAAAGUGGAGGCAUUCU	996	2561	AAAUGCCUCCACUUCU	1320
2557	UUCAUAAUAGAAAGGUGCCC	997	2557	UUCAUAAUAGAAAGGUGCCC	997	2579	GGGCACCUUCUUAUUAUAA	1321
2575	CAGAAAGACGAACUUGG	998	2575	CAGAAAGACGAACUUGG	998	2597	CCAAGUUCGUCUUCUUCU	1322
2593	GAAUACAUUAUUCUAG	999	2593	GAAUACAUUAUUCUAG	999	2615	CUACUAGAAUUAUUAUUC	1323
2611	GGCAGGCGGUGAUUGCCA	1000	2611	GGCAGGCGGUGAUUGCCA	1000	2633	UGGCAUACACGCGGUGCC	1324
2629	AUGUUCUUCUGGCUACUUC	1001	2629	AUGUUCUUCUGGCUACUUC	1001	2651	GAAGUAGCCAGAAACAU	1325
2647	CUUGUCAUACUCCUACGGA	1002	2647	CUUGUCAUACUCCUACGGA	1002	2669	UCCGUAGGAUGAUGACAAG	1326
2665	ACCGUUAAGCGGCGCAUUG	1003	2665	ACCGUUAAGCGGCGCAUUG	1003	2687	CAUUGCCCGCUUAACGGU	1327
2683	GGAGGGGAACUGAAGACAG	1004	2683	GGAGGGGAACUGAAGACAG	1004	2705	CUGUCUUCAGUUCGCCUCC	1328
2701	GGCUACUUGUCCUACUGCA	1005	2701	GGCUACUUGUCCUACUGCA	1005	2723	UGACGAUGGACAAAGUAGCC	1329
2719	AUGGAUCCAGAUAAACUCC	1006	2719	AUGGAUCCAGAUAAACUCC	1006	2741	GGAGUUCUUCGGAUCCAU	1330
2737	CCAUUGGAUAAACAUUGUG	1007	2737	CCAUUGGAUAAACAUUGUG	1007	2759	CACAUGUUCUUCGGAUUGG	1331
2755	GAACGACUCCUUAUGAUG	1008	2755	GAACGACUCCUUAUGAUG	1008	2777	CAUCAUAGGCGAGUCCUUG	1332
2773	GCCAGCAAUUGGGAUUC	1009	2773	GCCAGCAAUUGGGAUUC	1009	2795	GGAAUCCCAUUCUUGCUGGC	1333
2791	CCAGAGACCGGCUUGAAGC	1010	2791	CCAGAGACCGGCUUGAAGC	1010	2813	GCUUCAGCGGUCUUCUGGC	1334
2809	CUAGGUAGCCUUCUUGGCC	1011	2809	CUAGGUAGCCUUCUUGGCC	1011	2831	GGCCAAGAGGCUUACCUAG	1335
2827	CGUGGUGCCUUCUGGCAAG	1012	2827	CGUGGUGCCUUCUGGCAAG	1012	2849	CUUGGCCAAAGGCCACCACG	1336
2845	GUGAUUGAAGCAGAUGCCU	1013	2845	GUGAUUGAAGCAGAUGCCU	1013	2867	AGGCAUCUGCUCAUUCAC	1337
2863	UUUGGAAUUGACAAGACAG	1014	2863	UUUGGAAUUGACAAGACAG	1014	2885	CUUGCUUUGCAUUCUCCAA	1338
2881	GCAACUUGCAGGACAGUAG	1015	2881	GCAACUUGCAGGACAGUAG	1015	2903	CUACUGUCCUGCAAGUUGC	1339
2899	GCAGUCAAAUUGUUGAAAG	1016	2899	GCAGUCAAAUUGUUGAAAG	1016	2921	CUUUCACCAUUCUUGACUUC	1340
2917	GAAGGAGCAACACACAGUG	1017	2917	GAAGGAGCAACACACAGUG	1017	2939	CACUGUGUUGUCCUUC	1341
2935	GAGCAUCGAGCUCUACU	1018	2935	GAGCAUCGAGCUCUACU	1018	2957	ACAUGAGAGCUGCAUGCUC	1342
2953	UCUGAACUCAAGAUCCUCA	1019	2953	UCUGAACUCAAGAUCCUCA	1019	2975	UGAGGAUUCUUGAGUUCAGA	1343
2971	AUUCAUUUGGUCACCAUC	1020	2971	AUUCAUUUGGUCACCAUC	1020	2993	GAUGGUGACCAUUAUGAAU	1344
2989	CUCAUUGGUCUACCUUC	1021	2989	CUCAUUGGUCUACCUUC	1021	3011	GAAGGUGACCAUUAUGAG	1345
3007	CUAGGUGCCUUGUACCAAGC	1022	3007	CUAGGUGCCUUGUACCAAGC	1022	3029	GCUUGGUGACAGGCCUAG	1346
3025	CCAGGAGGCGCACUACUGG	1023	3025	CCAGGAGGCGCACUACUGG	1023	3047	CCAUAGUGGCCUCCUUGG	1347
3043	GUGAUUGGAAUUCUGCA	1024	3043	GUGAUUGGAAUUCUGCA	1024	3065	UGCAGAUUCCACAUCAC	1348
3061	AAAUUGGAAACCUUGCCA	1025	3061	AAAUUGGAAACCUUGCCA	1025	3083	UGGACAGGUUCCAAAUUU	1349
3079	ACUUAUUGGAGGAGCAAGA	1026	3079	ACUUAUUGGAGGAGCAAGA	1026	3101	UCUUGCUCUCCAGGUAAGU	1350
3097	AGAAUUGAAUUGUCCCU	1027	3097	AGAAUUGAAUUGUCCCU	1027	3119	AGGGGACAAUUCAUUUCU	1351
3115	UACAAGACCAAAAGGGGCAC	1028	3115	UACAAGACCAAAAGGGGCAC	1028	3137	GUGCCCUUUGGUCUUGUA	1352

3133	CGAUUCCGUCUAAAGGAAAG	1029	3133	CGAUUCCGUCUAAAGGAAAG	1029	3155	CUUCCUUGACGGAAUCG	1353
3151	GACUACGUUGGAGCAUCC	1030	3151	GACUACGUUGGAGCAUCC	1030	3173	GGAUUGCUCCAGGUAGUC	1354
3169	CCUGUGGAUCUGAAACGGC	1031	3169	CCUGUGGAUCUGAAACGGC	1031	3191	GCCGUUUCAGAUCCACAGG	1355
3187	CGCUUGGACAGCAUACCCA	1032	3187	CGCUUGGACAGCAUACCCA	1032	3209	UGGUUGAGCUUGUCCAAAGCG	1356
3205	AGUAGCCAGAGCUCAGCCA	1033	3205	AGUAGCCAGAGCUCAGCCA	1033	3227	UGGCUAGAGCUCUGGCUACU	1357
3223	AGCUUGGAAUUUGGAGG	1034	3223	AGCUUGGAAUUUGGAGG	1034	3245	CCUCCACAAUCCAGAGCU	1358
3241	GAGAAGUCCUCAGUGAUG	1035	3241	GAGAAGUCCUCAGUGAUG	1035	3263	CAUCACUAGGGAGCUUCUC	1359
3259	GUAGAAGAGGAGGAGCUC	1036	3259	GUAGAAGAGGAGGAGCUC	1036	3281	GAGCUUCCUUCUUCUAC	1360
3277	CCUGAAGUCUGUAUAGG	1037	3277	CCUGAAGUCUGUAUAGG	1037	3299	CCUUUACAGAUUCUACAGG	1361
3295	GACUUCUUGACCUUGGAGC	1038	3295	GACUUCUUGACCUUGGAGC	1038	3317	GCUCCAAGGUCAGGAAAGUC	1362
3313	CAUCUACUCUGUACAGCU	1039	3313	CAUCUACUCUGUACAGCU	1039	3335	AGCUGUACAGAUAGAGAU	1363
3331	UUCCAAGUGGCUAAGGGCA	1040	3331	UUCCAAGUGGCUAAGGGCA	1040	3353	UGCCCUUAGCCACUUGGAA	1364
3349	AUGGAGUUCUUGGCAUCGC	1041	3349	AUGGAGUUCUUGGCAUCGC	1041	3371	GCGAUGCCAAAGAACUCCAU	1365
3367	CGAAAGUGUAUCCACAGGG	1042	3367	CGAAAGUGUAUCCACAGGG	1042	3389	CCUGUGGAUACACUUCUG	1366
3385	GACCUUGGCGCAGCAAAUA	1043	3385	GACCUUGGCGCAGCAAAUA	1043	3407	UAUUUCGUGCCGCCAGGUC	1367
3403	AUCCUUCUUAUCCGAGAGA	1044	3403	AUCCUUCUUAUCCGAGAGA	1044	3425	UCUUCUCCGAUAAGAGGAU	1368
3421	AACGUGGUUAAAUCUGUG	1045	3421	AACGUGGUUAAAUCUGUG	1045	3443	CACAGAUUUUAAACCAGUU	1369
3439	GACUUGGCUUUGCCCGGG	1046	3439	GACUUGGCUUUGCCCGGG	1046	3461	CCCGGCCAAAGCCAAAGUC	1370
3457	GAUUAUUUAUAAAGAUCCAG	1047	3457	GAUUAUUUAUAAAGAUCCAG	1047	3479	CUGGAUCUUUUAUAAUUC	1371
3475	GAUUAUGUCAGAAAAGGAG	1048	3475	GAUUAUGUCAGAAAAGGAG	1048	3497	CUCUUUUCUGACAUAAUC	1372
3493	GAUGCUCGCCUCCCUUUGA	1049	3493	GAUGCUCGCCUCCCUUUGA	1049	3515	UCAAAGGAGGCCGAGCAUC	1373
3511	AAUUGGAUGGCCCCAGAAA	1050	3511	AAUUGGAUGGCCCCAGAAA	1050	3533	UUUCUGGGGCCAUCCAUUU	1374
3529	ACAAUUUUUAGACAGAGUGU	1051	3529	ACAAUUUUUAGACAGAGUGU	1051	3551	ACACUUCUGUCAAUAAUUGU	1375
3547	UACACAAUCCAGAGUAGCG	1052	3547	UACACAAUCCAGAGUAGCG	1052	3569	CGUCACUCUGGAUUUGUGUA	1376
3565	GUUCUGGCUUUUGGUGUUU	1053	3565	GUUCUGGCUUUUGGUGUUU	1053	3587	AAACACCAAAAGACCAGAC	1377
3583	UUGCUUGGGGAAAUUUUU	1054	3583	UUGCUUGGGGAAAUUUUU	1054	3605	AAAAUUAUUUCCACAGCAA	1378
3601	UCCUUAAGGUGCUUCUCCAU	1055	3601	UCCUUAAGGUGCUUCUCCAU	1055	3623	AUGGAGAAGCACCUAAGGA	1379
3619	UAUCCUGGGUAAAGAUUG	1056	3619	UAUCCUGGGUAAAGAUUG	1056	3641	CAUUCUUUACCCCGAGGAUA	1380
3637	GAUGAAGAAUUUUGUAGGC	1057	3637	GAUGAAGAAUUUUGUAGGC	1057	3659	GCCUACAAAUUUCUUAUC	1381
3655	CGAUUGAAAGAAAGGAACUA	1058	3655	CGAUUGAAAGAAAGGAACUA	1058	3677	UAGUUCUUCUUCUUAUC	1382
3673	AGAUUGAGGGCCCCUGAUU	1059	3673	AGAUUGAGGGCCCCUGAUU	1059	3695	AUUCAGGGGCCUUAUCUUC	1383
3691	UAUACUACACCCAGAAUUG	1060	3691	UAUACUACACCCAGAAUUG	1060	3713	ACAUUCUGGUGUAGUAUA	1384
3709	UACAGACCAUUGCUGGACU	1061	3709	UACAGACCAUUGCUGGACU	1061	3731	AGUCCAGCAUGGUCUGUA	1385
3727	UGCUGGCACGGGGAGCCCA	1062	3727	UGCUGGCACGGGGAGCCCA	1062	3749	UGGGCCCCCGUGCCAGCA	1386
3745	AGUCAGAGACCCACGUUUU	1063	3745	AGUCAGAGACCCACGUUUU	1063	3767	AAAACGUGGUGUCUGACU	1387
3763	UCAGAGUUGGUGGAACAUU	1064	3763	UCAGAGUUGGUGGAACAUU	1064	3785	AAUGUCCACCAACUCUGA	1388

3781	UUGGAAAUUCUUGCAAG	1065	3781	UUGGAAAUUCUUGCAAG	1065	3803	CUUGCAAGAGAUUUCCAA	1389
3789	GCUAUUGCUCAGCAGGAUG	1066	3789	GCUAUUGCUCAGCAGGAUG	1066	3821	GAUCCUUGCUGAGCAUUGCC	1390
3817	GGCAAAGACUACAUUGUUC	1067	3817	GGCAAAGACUACAUUGUUC	1067	3839	GAACAAUGUAGUUCUUGCC	1391
3835	CUUCCGAUUAUCAGAGACUU	1068	3835	CUUCCGAUUAUCAGAGACUU	1068	3857	AAGUCUCUGAUUUCGGAAG	1392
3853	UUGAGCAUGGAAGAGGAUU	1069	3853	UUGAGCAUGGAAGAGGAUU	1069	3875	AAUCCUUCUCCAUUCGCAA	1393
3871	UCUGGACUCUCUUGCCUA	1070	3871	UCUGGACUCUCUUGCCUA	1070	3893	UAGGCAGAGAGAGUCCAGA	1394
3889	ACCUCACCGUUGUCCUGUA	1071	3889	ACCUCACCGUUGUCCUGUA	1071	3911	UACAGGAACAGGUGAGGU	1395
3907	AUGGAGGAGGAGGAGGAUU	1072	3907	AUGGAGGAGGAGGAGGAUU	1072	3929	AUACUUCUCCUCCUCCAU	1396
3925	UGUGACCCCAAUUCCAUU	1073	3925	UGUGACCCCAAUUCCAUU	1073	3947	AAUGGAUUUUGGGUCACA	1397
3943	UAUGACAACACAGCAGGAA	1074	3943	UAUGACAACACAGCAGGAA	1074	3965	UUCUUGCUGUGUUGUCAUA	1398
3961	AUCAGUCAGUUAUCUGAGA	1075	3961	AUCAGUCAGUUAUCUGAGA	1075	3983	UCUGCAGAUACUGACUGAU	1399
3979	AACAGUAAGCGAAAGAGCC	1076	3979	AACAGUAAGCGAAAGAGCC	1076	4001	GGCUCUUUCGCUUACUGUU	1400
3997	CGGCCUGUGAGUGUAAAA	1077	3997	CGGCCUGUGAGUGUAAAA	1077	4019	UUUUACACUCACAGGCCG	1401
4015	ACAUUUGAAGAUUUCGCGU	1078	4015	ACAUUUGAAGAUUUCGCGU	1078	4037	ACGGGAUUAUUCUAAUUA	1402
4033	UUAAGAAGAACAGAGUAA	1079	4033	UUAAGAAGAACAGAGUAA	1079	4055	UUACUUCUGGUUUCUUA	1403
4051	AAAGUAUCCAGAUAGACA	1080	4051	AAAGUAUCCAGAUAGACA	1080	4073	UGUACUUGGGAUUACUUA	1404
4069	AACAGACGGACAGUGGUA	1081	4069	AACAGACGGACAGUGGUA	1081	4091	UACACUUGCCUGUGUU	1405
4087	AUGGUUCUUGCCUCAGAG	1082	4087	AUGGUUCUUGCCUCAGAG	1082	4109	CUUCUGAGGCAAGAACCAU	1406
4105	GAGCUGAAAACUUUGGAAG	1083	4105	GAGCUGAAAACUUUGGAAG	1083	4127	CUUCCAAAGUUUUCAGCUC	1407
4123	GACAGAACCAAUUAUCUC	1084	4123	GACAGAACCAAUUAUCUC	1084	4145	GAGUAUUUUGGUUCUGUC	1408
4141	CCAUCUUUUGGUGGAUUG	1085	4141	CCAUCUUUUGGUGGAUUG	1085	4163	CCAUCCACCAAAAGAUUG	1409
4159	GUGCCCAACAAAGCAGGG	1086	4159	GUGCCCAACAAAGCAGGG	1086	4181	CCCUGCUUUUGCUGGGCAG	1410
4177	GAGUCUGUGGCAUCUGAAG	1087	4177	GAGUCUGUGGCAUCUGAAG	1087	4199	CUUCAGAUGCCACAGACUC	1411
4195	GGCUCAAACACAGACAGCG	1088	4195	GGCUCAAACACAGACAGCG	1088	4217	CGCUUGCUGUGGUUUGAGCC	1412
4213	GGCUACCAAGUCCGGAUUC	1089	4213	GGCUACCAAGUCCGGAUUC	1089	4235	GAUAUCCGGACUGGUAGCC	1413
4231	CACUCCGAUGACACAGACA	1090	4231	CACUCCGAUGACACAGACA	1090	4253	UGUCUGUGUACUCCGAGUG	1414
4249	ACCACCGUGUACUCCAGUG	1091	4249	ACCACCGUGUACUCCAGUG	1091	4271	CACUGGAGUACACGGUGGU	1415
4267	GAGGAAGCAGAACUUUUA	1092	4267	GAGGAAGCAGAACUUUUA	1092	4289	UUAAAAGUUCUCCUCCUCC	1416
4285	AAGCUGAUAGAGAUUUGGAG	1093	4285	AAGCUGAUAGAGAUUUGGAG	1093	4307	CUCCAUCUUAUCAGCUU	1417
4303	GUAGCAACCGGUAGCAGAG	1094	4303	GUAGCAACCGGUAGCAGAG	1094	4325	CUUGUCUACCGGUUUGCAC	1418
4321	GCCAGAUUUCUCCAGCCUG	1095	4321	GCCAGAUUUCUCCAGCCUG	1095	4343	CAGGCUUGGAGAAUUGGGC	1419
4339	GACUCGGGGACACACUGA	1096	4339	GACUCGGGGACACACUGA	1096	4361	UCAGUGUGGUCCCGAGUUC	1420
4357	AGCUCUCCUCCUGUUUAAA	1097	4357	AGCUCUCCUCCUGUUUAAA	1097	4379	UUUAAACAGGAGGAGAGCU	1421
4375	AAGGAAGAUCCACACCCC	1098	4375	AAGGAAGAUCCACACCCC	1098	4397	GGGUGUGGAGUCCUCCUU	1422
4393	CAACUCCGGACAUCAU	1099	4393	CAACUCCGGACAUCAU	1099	4415	AUGUGAUGUCCGGAGUUG	1423
4411	UGAGAGGUCUGCUGAGAUU	1100	4411	UGAGAGGUCUGCUGAGAUU	1100	4433	AAUCUGAGCAGACCUCUCA	1424

4429	UUUGAAGUGUUGUUCUUC	1101	4429	UUUGAAGUGUUGUUCUUC	1101	4451	GAAAGAACACACUUCAAA	1425
4447	CCACCAGCAGGAAGUAGCC	1102	4447	CCACCAGCAGGAAGUAGCC	1102	4469	GGCUACUCCUGCUGGUGG	1426
4465	CGCAUUGAUUUUUAUUC	1103	4465	CGCAUUGAUUUUUAUUC	1103	4487	GAAUGAAAAUCAAUUGCG	1427
4483	CGACAAACAGAAAAAGGACC	1104	4483	CGACAAACAGAAAAAGGACC	1104	4505	GGUCCUUUUUCUGUUGUG	1428
4501	CUCGGACUGCAGGGAGCCA	1105	4501	CUCGGACUGCAGGGAGCCA	1105	4523	UGGCUCCUGCAGUGCCGAG	1429
4519	AGUCUUCUAGGCAUAUCCU	1106	4519	AGUCUUCUAGGCAUAUCCU	1106	4541	AGGAUUGCCUAGAAGACU	1430
4537	UGGAAGAGGCUUGUGACCC	1107	4537	UGGAAGAGGCUUGUGACCC	1107	4559	GGGUCACAAGCCUUCUCCA	1431
4555	CAAGAAUGGUCUGUAGCCG	1108	4555	CAAGAAUGGUCUGUAGCCG	1108	4577	AGACACAGACACAUUCUUG	1432
4573	UUCUCCAGUGUUGACCCUG	1109	4573	UUCUCCAGUGUUGACCCUG	1109	4595	CAGGUCACACUGGGAGAA	1433
4591	GAUCCUCUUUUUAUUA	1110	4591	GAUCCUCUUUUUAUUA	1110	4613	UGAAUGAAAAAGAGGAUC	1434
4609	AUUUAAAAAGCAUUUAU	1111	4609	AUUUAAAAAGCAUUUAU	1111	4631	AUGAUUUGCUUUUUUAU	1435
4627	UGCCCUUGCUGCGGGUCUC	1112	4627	UGCCCUUGCUGCGGGUCUC	1112	4649	GAGACCCGACGAGGGGCA	1436
4645	CACCAUGGGUUUAGAACAA	1113	4645	CACCAUGGGUUUAGAACAA	1113	4667	UUUUCUUAACCCAUUGUG	1437
4663	AAGAGCUUACAAGCAUUGG	1114	4663	AAGAGCUUACAAGCAUUGG	1114	4685	GCCAUUGCUUGAAGCUCU	1438
4681	CCCCAUCCUCAAAGAGUA	1115	4681	CCCCAUCCUCAAAGAGUA	1115	4703	UACUUCUUGAGGAUGGG	1439
4699	AGCAGUACCUUGGGAGCUG	1116	4699	AGCAGUACCUUGGGAGCUG	1116	4721	CAGCUCUCCAGGUACUCU	1440
4717	GACACUUCUGUAAAACUAG	1117	4717	GACACUUCUGUAAAACUAG	1117	4739	CUAGUUUUUACAGAAUGUC	1441
4735	GAAGAUAAACAGGCAACG	1118	4735	GAAGAUAAACAGGCAACG	1118	4757	CGUUGCCUGGUUUUAUCU	1442
4753	GUAGUGUUCGAGGUUUG	1119	4753	GUAGUGUUCGAGGUUUG	1119	4775	CAACACUUGAACACACUAC	1443
4771	GAAGUUGGAAAGGAUUUG	1120	4771	GAAGUUGGAAAGGAUUUG	1120	4793	GCAAUUCCUUCUCCAUUC	1444
4789	CAGGCUUGAGUUAUCCAA	1121	4789	CAGGCUUGAGUUAUCCAA	1121	4811	UUUGAUAGACUCAGCCUG	1445
4807	AGAGGUUUGUUUAGGACG	1122	4807	AGAGGUUUGUUUAGGACG	1122	4829	CGUCCUAAACAAAGCCUCU	1446
4825	GUGGGUCCCAAGCCAAGCC	1123	4825	GUGGGUCCCAAGCCAAGCC	1123	4847	GGCUUGGCUUGGGACCCAC	1447
4843	CUUAAAGUGUGGAUUUCGA	1124	4843	CUUAAAGUGUGGAUUUCGA	1124	4865	UCCGAUUUCCACACUUAAG	1448
4861	AUUGAUAGAAAGGAAGACU	1125	4861	AUUGAUAGAAAGGAAGACU	1125	4883	AGUCUUCUUCUUAUCAAU	1449
4879	UAACGUUACCUUGCUUUG	1126	4879	UAACGUUACCUUGCUUUG	1126	4901	CCAAAGCAAGUAACGUUA	1450
4897	GAGAGUACUGGAGCCUGCA	1127	4897	GAGAGUACUGGAGCCUGCA	1127	4919	UGCAGGCUCCAGUACUCUC	1451
4915	AAUUGCAUUGUUUUGCUC	1128	4915	AAUUGCAUUGUUUUGCUC	1128	4937	GAGCAACACAAUGCAUUU	1452
4933	CUGGUGGAGGUGGGCAUGG	1129	4933	CUGGUGGAGGUGGGCAUGG	1129	4955	CCAUGCCACCUCCACCAG	1453
4951	GGUCUGUUCUGAAAUUGUA	1130	4951	GGUCUGUUCUGAAAUUGUA	1130	4973	UACAUUUGAGAAACAGACCC	1454
4969	AAAGGUUUCAGACGGGUU	1131	4969	AAAGGUUUCAGACGGGUU	1131	4991	AACCCGUCUGAACCCUUCU	1455
4987	UUCUGGUUUUAGAAAGUUG	1132	4987	UUCUGGUUUUAGAAAGUUG	1132	5009	CAACCUUCUAAAACCAGAA	1456
5005	GCGUGUUCUUCGAGUUGG	1133	5005	GCGUGUUCUUCGAGUUGG	1133	5027	CCCAACUJGSAAGAACACGC	1457
5023	GCUAAAGUAGAGUUCGUUG	1134	5023	GCUAAAGUAGAGUUCGUUG	1134	5045	CAACGAACUUAUUCUUAAGC	1458
5041	GUGCUGUUUCGACUCCUA	1135	5041	GUGCUGUUUCGACUCCUA	1135	5063	UAGGAGUCAGAAACAGCAC	1459
5059	AAUGAGAGUUCUCCUCCAGA	1136	5059	AAUGAGAGUUCUCCUCCAGA	1136	5081	UCUGGAAGGAACUUCUUAU	1460

5077	ACCGUAGCUGUCUCCUUG	1137	5077	ACCGUAGCUGUCUCCUUG	1137	5099	CAAGGAGACAGCUAACGGU	1481
5095	GCCAAAGCCCGAGGAGAA	1138	5095	GCCAAAGCCCGAGGAGAA	1138	5117	UUUCUUCUUGGGGCUUGGC	1482
5113	AAUGAGCAGCUCUGGCUC	1139	5113	AAUGAGCAGCUCUGGCUC	1139	5135	GAGCCAGAGCUGCAUCAUU	1483
5131	CCUUGUCUCCAGGCGUAG	1140	5131	CCUUGUCUCCAGGCGUAG	1140	5153	AUCAGCCUGGAGACAAGG	1484
5149	UCCUUAUUCAGAAUACCA	1141	5149	UCCUUAUUCAGAAUACCA	1141	5171	UGGUUAUUCUGAAUAAAGGA	1485
5167	ACAAAGAAAGGACAUUCAG	1142	5167	ACAAAGAAAGGACAUUCAG	1142	5188	CUGAAUGUCCUUCUUCUUGU	1486
5185	GCUCAGGCUCCCGCCG	1143	5185	GCUCAGGCUCCCGCCG	1143	5207	ACGGCAGGAGCCUUGAGC	1487
5203	UGUUGAAGAGUUCUGACUG	1144	5203	UGUUGAAGAGUUCUGACUG	1144	5225	CAGUCAGAACUCUUCACA	1488
5221	GCACAAACAGCUUCUGGU	1145	5221	GCACAAACAGCUUCUGGU	1145	5243	ACCAGAAGCUGGUUUGUC	1489
5238	UUUCUUCUGGAAUAGAAUAC	1146	5238	UUUCUUCUGGAAUAGAAUAC	1146	5261	GUUAUUAUCCAGAAAGAA	1470
5257	CCCUCAUUUCUGUCCUGAU	1147	5257	CCCUCAUUUCUGUCCUGAU	1147	5279	AUCAGGACAGAUUAGAGGG	1471
5275	UGUGAUUUGUUGAGACUG	1148	5275	UGUGAUUUGUUGAGACUG	1148	5297	CAGUCUACAGCAUAUCACA	1472
5293	GAUUGCGGAGGUUCAUUG	1149	5293	GAUUGCGGAGGUUCAUUG	1149	5315	CAUUGAACCCUCCCGCAUUC	1473
5311	GUGAAGCUGUGUGUGUGU	1150	5311	GUGAAGCUGUGUGUGUGU	1150	5333	ACACCACACACAGCUUCAC	1474
5329	UCAAAGUUUCAGGAGGGAU	1151	5329	UCAAAGUUUCAGGAGGGAU	1151	5351	AUCCUUCUGAAACUUUGA	1475
5347	UUUUACCCUUUUGUUCUUC	1152	5347	UUUUACCCUUUUGUUCUUC	1152	5369	GAAGAACAAAAGGGUAAAA	1476
5365	CCCCUCUGCCCAACCCAC	1153	5365	CCCCUCUGCCCAACCCAC	1153	5387	GUGGGUUGGGGACAGGGGG	1477
5383	CUCUCACCCCGCAACCCAU	1154	5383	CUCUCACCCCGCAACCCAU	1154	5405	AUGGUUUGCGGGUGAGAG	1478
5401	UCAGUAUUUUAUUAUUAUUG	1155	5401	UCAGUAUUUUAUUAUUAUUG	1155	5423	CAAAUAAUAAUUAUUAUUA	1479
5419	GGCCUCUACUCCAGUAAAC	1156	5419	GGCCUCUACUCCAGUAAAC	1156	5441	GUUUAUUGGAGUAGAGGCC	1480
5437	CCUGAUUUGGUUUGUUCAC	1157	5437	CCUGAUUUGGUUUGUUCAC	1157	5459	GUGAACAAACCCCAUUCAGG	1481
5455	CUCUCUGAAUUAUUAUUAUAG	1158	5455	CUCUCUGAAUUAUUAUUAUAG	1158	5477	CUAAUUAUUAUUAUUAUUA	1482
5473	GCCAGACUUCAAAUAUUAU	1159	5473	GCCAGACUUCAAAUAUUAU	1159	5495	AAUAAUUUUGAAGUCUGGC	1483
5491	UUUAUAGCCCAAAUUAUA	1160	5491	UUUAUAGCCCAAAUUAUA	1160	5513	UUAAUUAUUAUUAUUAUUA	1484
5509	ACAUCUAUUGUAUUAUUAUUA	1161	5509	ACAUCUAUUGUAUUAUUAUUA	1161	5531	UAAAUUAUUAUUAUUAUUA	1485
5527	AGACUUUUUAACAUUAUUAUAG	1162	5527	AGACUUUUUAACAUUAUUAUAG	1162	5549	CUCUAUUAUUAUUAUUAUUA	1486
5545	GCUAUUUCUACUGAUUUUU	1163	5545	GCUAUUUCUACUGAUUUUU	1163	5567	AAAAUUCAGUAGAAUUAUAGC	1487
5563	UGCCCUUUGUUCUGUCCUUU	1164	5563	UGCCCUUUGUUCUGUCCUUU	1164	5585	AAAGGACAGAACCAAGGGCA	1488
5581	UUUUUCAAUUAUUAUUAUUAUAG	1165	5581	UUUUUCAAUUAUUAUUAUUAUAG	1165	5603	CAUUUUUUUUUUUUGAAAAA	1489
5599	GUGUUUUUUUUUUGGUUACC	1166	5599	GUGUUUUUUUUUUGGUUACC	1166	5621	GGUACCAACAAAAAACAC	1490
5617	CAUAGUGGAAUUGCUGGG	1167	5617	CAUAGUGGAAUUGCUGGG	1167	5639	CCCAGCAUUAACACAUUAG	1491
5635	GAACAAGUCACUAUAAGACA	1168	5635	GAACAAGUCACUAUAAGACA	1168	5657	UGUCUUAUUAUUAUUAUUAUUG	1492
5653	AUGCUAUGGCACAUUAUUAU	1169	5653	AUGCUAUGGCACAUUAUUAU	1169	5675	AAUUAUUAUUAUUAUUAUUAU	1493
5671	UUUAUAGUCUGUUAUUAUUAU	1170	5671	UUUAUAGUCUGUUAUUAUUAU	1170	5693	CUACAUAUUAUUAUUAUUAUUA	1494
5689	GAACAUAUUAUUAUUAUUAUUAU	1171	5689	GAACAUAUUAUUAUUAUUAUUAU	1171	5711	AAUUAUUAUUAUUAUUAUUAU	1495
5707	UAAAGCCUUAUUAUUAUUAUUAU	1172	5707	UAAAGCCUUAUUAUUAUUAUUAU	1172	5729	CAUUAUUAUUAUUAUUAUUAU	1496



5725	GAACUUUUGUACUUAUCACA	1173	5725	GAACUUUUGUACUUAUCACA	1173	5747	UGUGAAUAGUACAAAGUUC	1497
5743	AUUUUGUAUCAGUUAUUG	1174	5743	AUUUUGUAUCAGUUAUUG	1174	5765	CAUAAUACUGAUACAAAUA	1498
5761	GUAGCAUACAAAGGUCAU	1175	5761	GUAGCAUACAAAGGUCAU	1175	5783	AUGACCUUUGUUAUGCUAC	1499
5779	UAAUGCUUUCAGCAUUGA	1176	5779	UAAUGCUUUCAGCAUUGA	1176	5801	UCAAUUGCUGAAAGCAUUA	1500
5797	AUGUCAUUUUAUUAAGAA	1177	5797	AUGUCAUUUUAUUAAGAA	1177	5819	UUCUUUAUAAAUUAGACAU	1501
5812	AGAACAUUGAAAAACUUGA	1178	5812	AGAACAUUGAAAAACUUGA	1178	5834	UCAAGUUUUUCAAUGUUCU	1502

gi|4503752|ref|NM\_002020.1

Pos	Target Sequence	Seq ID	UPos	Upper seq	Seq ID	LPos	Lower seq	Seq ID
1	ACCCACGCGCAGCGGCCGG	1503	1	ACCCACGCGCAGCGGCCGG	1503	23	CCGGCCGCGUGCGGUGGGU	1750
19	GAGAUAGCAGCGGGCGCCG	1504	19	GAGAUAGCAGCGGGCGCCG	1504	41	CGGCGCCCGCUGCAUCUC	1751
37	GCGCUGUGCCUGCGACUGU	1505	37	GCGCUGUGCCUGCGACUGU	1505	59	ACAGUCGACGGCACAGCGC	1752
55	UGGCUCUGCCUGGGACUCC	1506	55	UGGCUCUGCCUGGGACUCC	1506	77	GGAGUCCCGAGGAGAGCCA	1753
73	CUGGACGGCCUGGUGAGUG	1507	73	CUGGACGGCCUGGUGAGUG	1507	95	CACUACACAGGCCGUCGAG	1754
91	GACUACUCCAUAGACCCCC	1508	91	GACUACUCCAUAGACCCCC	1508	113	GGGGGUGCAUGGAGUAGUC	1755
109	CCGACCUUGAACAUACAGG	1509	109	CCGACCUUGAACAUACAGG	1509	131	CCGUGAUGUUAAGGUCGG	1756
127	GAGGAGUACACAGCUAUCG	1510	127	GAGGAGUACACAGCUAUCG	1510	149	CGAUGACGUGUGACUCCUC	1757
145	GACACCGGUGACAGCCUGU	1511	145	GACACCGGUGACAGCCUGU	1511	167	ACAGGCUUGUACCGGUGUC	1758
163	UCCAUUCUCCUGCAGGGGAC	1512	163	UCCAUUCUCCUGCAGGGGAC	1512	185	GUCCCCUGCAGGAGUUGGA	1759
181	CAGCACCCUCCUGAGUGGG	1513	181	CAGCACCCUCCUGAGUGGG	1513	203	CCCACUCGAGGGGGUGGUG	1760
199	GCUUGGCCAGGAGCUCAGG	1514	199	GCUUGGCCAGGAGCUCAGG	1514	221	CCUGAGCUCUCCUGGCCAAGC	1761
217	GAGGCGCCAGCCACCCGAG	1515	217	GAGGCGCCAGCCACCCGAG	1515	239	CUCCGGUGGCGUGGCGCCUC	1762
235	GACAAGGACAGCGAGACA	1516	235	GACAAGGACAGCGAGACA	1516	257	UGUCCUCGCUUGUCCUUGUC	1763
253	ACGGGGUGUGGCGGAGACU	1517	253	ACGGGGUGUGGCGGAGACU	1517	275	AGUCUCGACCCACCCCGGU	1764
271	UGCGAGGGCACAGACGCCA	1518	271	UGCGAGGGCACAGACGCCA	1518	293	UGGCGUCUGUGCCCUUGCA	1765
289	AGGCCCCUACUGCAAGGUGU	1519	289	AGGCCCCUACUGCAAGGUGU	1519	311	ACACCUUGCAGUAGGGCCU	1766
307	UUGCUGCUGCACGAGGUAC	1520	307	UUGCUGCUGCACGAGGUAC	1520	329	GUACCUCCUGCAGCAGCAA	1767
325	CAUGCCAAACACACAGGCA	1521	325	CAUGCCAAACACACAGGCA	1521	347	UGCCUGUGUCGUUGGCAUG	1768
343	AGCUACGUCUGCUACUACA	1522	343	AGCUACGUCUGCUACUACA	1522	365	UGUAGUAGCAGACAGUAGCU	1769
361	AAGUACAUCAAAGGCACGCA	1523	361	AAGUACAUCAAAGGCACGCA	1523	383	UGCGUGCCUUGAUUACUUU	1770
379	AUCGAGGGCACCCACGGCCG	1524	379	AUCGAGGGCACCCACGGCCG	1524	401	CGGCGGUGGUGGCCUUGCAU	1771
397	GCCAGCUCCUACGUGUUCG	1525	397	GCCAGCUCCUACGUGUUCG	1525	419	CGAACACAGUAGGAGCUGGC	1772



415	GUGAGAGACUUUGAGCAGC	1526	415	GUGAGAGACUUUGAGCAGC	1526	437	GCUGCUCAAAGUUCUCAC	1773
433	CCAUUCAUCAACAAGCCUG	1527	433	CCAUUCAUCAACAAGCCUG	1527	455	CAGGCUUGUUGAUGAUGG	1774
451	GACACGCUUUGGUCAACA	1528	451	GACACGCUUUGGUCAACA	1528	473	UGUUGACCAAGAGCGUGUC	1775
469	AGGAAGGACGCCAUGUGGG	1529	469	AGGAAGGACGCCAUGUGGG	1529	491	CCCACAUUGGCGUCCUUCU	1776
487	GUGCCCUUGUGUGUCCA	1530	487	GUGCCCUUGUGUGUCCA	1530	509	UGGACACCAGACAGGGCAC	1777
505	AUCCCGGCCCUCAAUUGCA	1531	505	AUCCCGGCCCUCAAUUGCA	1531	527	UGACAUUGAGGCCGGGGAU	1778
523	ACGUGCGCUCGCAAGCU	1532	523	ACGUGCGCUCGCAAGCU	1532	545	AGCUUUGCGAGCGCAGCGU	1779
541	UCGGUGCUGGGCCAGAGC	1533	541	UCGGUGCUGGGCCAGAGC	1533	563	CGUCUGGCCACAGCACCGA	1780
559	GGCAGGAGGUGGUGUGGG	1534	559	GGCAGGAGGUGGUGUGGG	1534	581	CCACACACCACCUCCUGCCC	1781
577	GAUACCGGGGGCAUGC	1535	577	GAUACCGGGGGGGCAUGC	1535	599	GCAUGCCCCCGCCGGUCAUC	1782
595	CUCGUGUCCACGCCACUGC	1536	595	CUCGUGUCCACGCCACUGC	1536	617	GCAGUGGCGUGGACACGAG	1783
613	CUGCAGAUGCCUGUACC	1537	613	CUGCAGAUGCCUGUACC	1537	635	GGUACAGGGCAUCUGUGCAG	1784
631	CUGCAGUGCGAGACCACCU	1538	631	CUGCAGUGCGAGACCACCU	1538	653	AGGUGGUCUCGCACUGCAG	1785
649	UGGGGAGACCAGGACUUC	1539	649	UGGGGAGACCAGGACUUC	1539	671	GGAAGUCCUGGUUCUCCCCA	1786
667	CUUCCAAACCCUUCUGG	1540	667	CUUCCAAACCCUUCUGG	1540	689	CCAGGAAGGGGUUGGAAAG	1787
685	GUGCACAUCACAGGCAACG	1541	685	GUGCACAUCACAGGCAACG	1541	707	CGUUGCCUGUGAUGUGCAG	1788
703	GAGCUCUUGACAUCACG	1542	703	GAGCUCUUGACAUCACG	1542	725	GCUGGAUGUCAUAGAGCUC	1789
721	CUGUUGCCCAGGAAGUCG	1543	721	CUGUUGCCCAGGAAGUCG	1543	743	GCGACUUCUGGGGCAACAG	1790
739	CUGGAGCUGUGGUAGGGG	1544	739	CUGGAGCUGUGGUAGGGG	1544	761	CCCCUACCAGCAGCUCCAG	1791
757	GAGAAGCUGGUCCUACU	1545	757	GAGAAGCUGGUCCUACU	1545	779	AGUUGAGGACCAGCUUCUC	1792
775	UGCACCGUGUGGGCUGAGU	1546	775	UGCACCGUGUGGGCUGAGU	1546	797	ACUCAGCCACACACGGUGCA	1793
793	UUUAACUCAGGUGUCACCU	1547	793	UUUAACUCAGGUGUCACCU	1547	815	AGGUGACACCUAGAGUUAAA	1794
811	UUUGACUGGGACUACCCAG	1548	811	UUUGACUGGGACUACCCAG	1548	833	CUGGGUAGUCCAGUCUAAA	1795
829	GGGAAGCAGGACAGCGGG	1549	829	GGGAAGCAGGACAGCGGG	1549	851	CCCGCUCUGCCUGCUUCCC	1796
847	GGUAAAGUGGUGCCCGAGC	1550	847	GGUAAAGUGGUGCCCGAGC	1550	869	GCUCGGGCACCCACUJACC	1797
865	CGACGCUCCCAACAGACCC	1551	865	CGACGCUCCCAACAGACCC	1551	887	GGGUCUCUUGGGAGCGGUCG	1798
883	CACACAGAACUCUCCAGCA	1552	883	CACACAGAACUCUCCAGCA	1552	905	UGCUGGAGAGUUCUGUGUG	1799
901	AUCCUGACCACUCCACAACG	1553	901	AUCCUGACCACUCCACAACG	1553	923	CGUUGUGGAUGGUCAGGAU	1800
919	GUACGCCAGCAGGACCUUG	1554	919	GUACGCCAGCAGGACCUUG	1554	941	CCAGGUCGUGCUGGCCUGAC	1801
937	GGCUCGUUUGUGCAAGG	1555	937	GGCUCGUUUGUGCAAGG	1555	959	CCUUGCACACAUACGAGCC	1802
955	GCCAACAACGGCAUCCAGC	1556	955	GCCAACAACGGCAUCCAGC	1556	977	GCUGGAUGCCCGUUGUGGC	1803
973	CGAUUUCGGGAGAGCACCG	1557	973	CGAUUUCGGGAGAGCACCG	1557	995	CGGUGCUCUCCCGAAUCCG	1804
991	GAGGUCAUUGUGCAUGAAA	1558	991	GAGGUCAUUGUGCAUGAAA	1558	1013	UUUCAUGCACAAUGACCUC	1805

1009	AAUCCCUUUAUACAGCGUCG	1559	1009	AAUCCCUUUAUACAGCGUCG	1559	1031	CGACGCUGAUGAAGGGAUU	1806
1027	GAGUGGCUCAAAGGACCCA	1560	1027	GAGUGGCUCAAAGGACCCA	1560	1049	UGGGUCCUUUGAGCCACUC	1807
1045	AUCCUGGAGGCCACGGCAG	1561	1045	AUCCUGGAGGCCACGGCAG	1561	1067	CUGCCGUGGGCCUCCAGGAU	1808
1063	GGAGACGAGCUGGUGAAGC	1562	1063	GGAGACGAGCUGGUGAAGC	1562	1085	GCUUACACGAGCUCGUCUC	1809
1081	CUGCCCGUGAAGCUGGCAG	1563	1081	CUGCCCGUGAAGCUGGCAG	1563	1103	CUGCCAGCUUACACGGGCAG	1810
1099	GCGUACCCCGCCCGCCGAGU	1564	1099	GCGUACCCCGCCCGCCGAGU	1564	1121	ACUCGGCGGGGGGUAACGC	1811
1117	UUCACAGUGGUACAAGGAUG	1565	1117	UUCACAGUGGUACAAGGAUG	1565	1139	CAUCCUUGUACACACUGGAA	1812
1135	GGAAGGCACUUGUCCGGGC	1566	1135	GGAAGGCACUUGUCCGGGC	1566	1157	GCCCGACAGUGCCUUUCC	1813
1153	CGCCACAGUCCACAUGCC	1567	1153	CGCCACAGUCCACAUGCC	1567	1175	GGCAUGUGGACUUGUGGCG	1814
1171	CUGGUGCUCAAGGAGGUGA	1568	1171	CUGGUGCUCAAGGAGGUGA	1568	1193	UCACCUCCUUGAGCACCAG	1815
1189	ACAGAGGCCAGCACAGGCA	1569	1189	ACAGAGGCCAGCACAGGCA	1569	1211	UGCCUGUGCUGGCCUCUGU	1816
1207	ACCUACACCCUCGCCUUGU	1570	1207	ACCUACACCCUCGCCUUGU	1570	1229	ACAGGGCGAGGGUGUAGGU	1817
1225	UGGAACUCCGUGCUGGCC	1571	1225	UGGAACUCCGUGCUGGCC	1571	1247	GGCCAGCAGCGGAGUUCCA	1818
1243	CUGAGGCGCAACAUCAGCC	1572	1243	CUGAGGCGCAACAUCAGCC	1572	1265	GGCUGAUGUUGCGCCUCAG	1819
1261	CUGGAGCUGGUGGUGAAUG	1573	1261	CUGGAGCUGGUGGUGAAUG	1573	1283	CAUUCACACACAGCUCCAG	1820
1279	GUGCCCCCAGAUACAUG	1574	1279	GUGCCCCCAGAUACAUG	1574	1301	CAUGUAUCUGGGGGGCGAC	1821
1297	GAGAAGGAGGCCUCCUCC	1575	1297	GAGAAGGAGGCCUCCUCC	1575	1319	GGGAGGAGGCCUCCUUCUC	1822
1315	CCCAGCAUCUACUCCGUC	1576	1315	CCCAGCAUCUACUCCGUC	1576	1337	GACGCGAGUAGAUUCUGGG	1823
1333	CACAGCGCCAGGCCCUCA	1577	1333	CACAGCGCCAGGCCCUCA	1577	1355	UGAGGGCCUUGCGGCGUGG	1824
1351	ACUUGCAGCGCCUACGGGG	1578	1351	ACUUGCAGCGCCUACGGGG	1578	1373	CCCUGUAGGCCGUGCAGGU	1825
1369	GUCCCCUGCCUCUCAGCA	1579	1369	GUCCCCUGCCUCUCAGCA	1579	1391	UGCUGAGAGCGAGGGCGAC	1826
1387	AUCCAGUGGCACUUGCGGC	1580	1387	AUCCAGUGGCACUUGCGGC	1580	1409	GCCGCCAGUGCCACUUGGAU	1827
1405	CCUUGGACACCCUGCAAGA	1581	1405	CCUUGGACACCCUGCAAGA	1581	1427	UCUUGCAGGGUGUCCAGGG	1828
1423	AUGUUUGCCAGCGUAGUC	1582	1423	AUGUUUGCCAGCGUAGUC	1582	1445	GACUACGUGGGGCAACAU	1829
1441	CUCCGGCGCGGAGCAGC	1583	1441	CUCCGGCGCGGAGCAGC	1583	1463	GCUCUGCCCGCCCGCGGAG	1830
1459	CAAGACCUCAUGCCACAGU	1584	1459	CAAGACCUCAUGCCACAGU	1584	1481	ACUGUGCAUGAGGUCUUG	1831
1477	UGCCGUGACUGGAGGGCGG	1585	1477	UGCCGUGACUGGAGGGCGG	1585	1499	CCGCCUCCAGUCACGGCA	1832
1495	GUGACCACGAGGAUUGCCG	1586	1495	GUGACCACGAGGAUUGCCG	1586	1517	CGGCAUCCUGCGUGGUCAC	1833
1513	GUGAACCCCAUCGAGAGCC	1587	1513	GUGAACCCCAUCGAGAGCC	1587	1535	GGCUCUGAUGGGGUCAC	1834
1531	CUGGACACCCUGGACCGAGU	1588	1531	CUGGACACCCUGGACCGAGU	1588	1553	ACUGGGUCCAGGUGUCCAG	1835
1549	UUUGUGGAGGGAAGAAUA	1589	1549	UUUGUGGAGGGAAGAAUA	1589	1571	UAUUCUUUCCUCCACAAA	1836
1567	AAGACUGAGCAAGCUGG	1590	1567	AAGACUGAGCAAGCUGG	1590	1589	CCAGCUUGCUCACAGUCUU	1837
1585	GUGAUCCAGAAUGCCAACG	1591	1585	GUGAUCCAGAAUGCCAACG	1591	1607	CGUUGGCAUUCUGGAUCAC	1838

1603	GUUUCUGCCAUUUAAGU	1592	1603	GUUUCUGCCAUUUAAGU	1592	1625	ACUUGUACUUGGAGACAC	1839
1621	UGUGUGGUCUCCAAAGG	1593	1621	UGUGUGGUCUCCAAAGG	1593	1643	CCUUGUUGGAGACCACACA	1840
1639	GUUGGCCAGGAUGAGCGGC	1594	1639	GUUGGCCAGGAUGAGCGGC	1594	1661	GCCGCUCAUCCUGGCCAC	1841
1657	CUCAUCUACUUCUAUGUGA	1595	1657	CUCAUCUACUUCUAUGUGA	1595	1678	UCACAUAGAAGUAGAUGAG	1842
1675	ACCACCAUCCCGACGGCU	1596	1675	ACCACCAUCCCGACGGCU	1596	1697	AGCCGUCGGGAUGGUGGU	1843
1693	UUCACCAUCGAAUCCAAGC	1597	1693	UUCACCAUCGAAUCCAAGC	1597	1715	GCUUGGAUUGGAUGGUGAA	1844
1711	CCAUCGAGGAGCUACUAG	1598	1711	CCAUCGAGGAGCUACUAG	1598	1733	CUAGUAGCUCCUGGGAUGG	1845
1729	GAGGCCAGCCGGUGCUCC	1599	1729	GAGGCCAGCCGGUGCUCC	1599	1751	GGAGCACCGGUGGCCCUCC	1846
1747	CUGAGCUGCCAAGCCGACA	1600	1747	CUGAGCUGCCAAGCCGACA	1600	1769	UGUGGCUUGGAGCUCAG	1847
1765	AGCUACAAGUACGAGCAUC	1601	1765	AGCUACAAGUACGAGCAUC	1601	1787	GAUGCUGUACUUGUAGCU	1848
1783	CUGCGCUGGUACCGCCUCA	1602	1783	CUGCGCUGGUACCGCCUCA	1602	1805	UGAGCGGUACCAAGCGCAG	1849
1801	AACCUUGUCCACGUCGACG	1603	1801	AACCUUGUCCACGUCGACG	1603	1823	CGUGCAGCGUGGACAGGUU	1850
1819	GAUGCGCACGGGAACCCGC	1604	1819	GAUGCGCACGGGAACCCGC	1604	1841	GCGGUUCCCGUGCGCAUC	1851
1837	CUUCUGCUCGACUGCAAGA	1605	1837	CUUCUGCUCGACUGCAAGA	1605	1859	UCUUGCAGUCGAGCAGAAG	1852
1855	AACGUGCAUCUGUUCGCCA	1606	1855	AACGUGCAUCUGUUCGCCA	1606	1877	UGGCGAACAGAUUGCAGU	1853
1873	ACCCUCUGGCCCGCCAGCC	1607	1873	ACCCUCUGGCCCGCCAGCC	1607	1895	GGCUGGGGCCAGAGGGGU	1854
1891	CUGGAGGAGGUGGCCACCU	1608	1891	CUGGAGGAGGUGGCCACCU	1608	1913	CAGGUGCCACCUCCUCCAG	1855
1909	GGGCGCGCCACGCCACGC	1609	1909	GGGCGCGCCACGCCACGC	1609	1931	GCGUGGCGUGGCGCGCCCC	1856
1927	CUCAGCCUGAGUAUCCGCC	1610	1927	CUCAGCCUGAGUAUCCGCC	1610	1949	GGGGGAUACUAGGCGUGAG	1857
1945	CGGUGCGCGCCGAGCAGC	1611	1945	CGGUGCGCGCCGAGCAGC	1611	1967	CGUGCUCGGCGCGCAGCGG	1858
1963	GAGGCCACUAUGUGUGCG	1612	1963	GAGGCCACUAUGUGUGCG	1612	1985	CGCACAUAGUUGGCCCUCC	1859
1981	GAAGUGCAAGACCGCGCA	1613	1981	GAAGUGCAAGACCGCGCA	1613	2003	UGGCCCGGUCUUGCAGUUC	1860
1999	AGCCAUGACAAGCACUGCC	1614	1999	AGCCAUGACAAGCACUGCC	1614	2021	GGCAGUGCUUGUCAUGGCU	1861
2017	CACAAGAAGUACCUUGUCGG	1615	2017	CACAAGAAGUACCUUGUCGG	1615	2039	CCGACAGGUACUUCUUGUG	1862
2035	GUCCAGGCCUUGGAAGCCC	1616	2035	GUCCAGGCCUUGGAAGCCC	1616	2057	GGGCUUCCAGGGCCUUGCAC	1863
2053	CCUCGGCUCACGAGAAACU	1617	2053	CCUCGGCUCACGAGAAACU	1617	2075	AGUUCUGCGUGAGCCGAGG	1864
2071	UUGACCGACCUCCUGGUGA	1618	2071	UUGACCGACCUCCUGGUGA	1618	2093	UCACCAGGAGGUCCGUGCAA	1865
2089	AACGUGAGCGACUCGCUUG	1619	2089	AACGUGAGCGACUCGCUUG	1619	2111	CCAGCGAGUCGCUACGCUU	1866
2107	GAGUAGCAGUGCUUGGUGG	1620	2107	GAGUAGCAGUGCUUGGUGG	1620	2129	CCACCAAGCACUGCAUCUC	1867
2125	GCCGAGCGCACGCGCCCA	1621	2125	GCCGAGCGCACGCGCCCA	1621	2147	UGGGCGGUGGCGCUCGCGC	1868
2143	AGCAUCGUGUGGUACAAG	1622	2143	AGCAUCGUGUGGUACAAG	1622	2165	CUUUGUACACACAGUAGCU	1869
2161	GACGAGAGGCUUGGAGG	1623	2161	GACGAGAGGCUUGGAGG	1623	2183	CCUCCAGCAGCCUUCUGUC	1870
2179	GAAAGUCUGGAGUCGACU	1624	2179	GAAAGUCUGGAGUCGACU	1624	2201	AGUCGACUCCAGACUUCUUC	1871

2197	UUGCGGACUCCAAACGAGA	1625	2197	UUGCGGACUCCAAACGAGA	1625	2219	UCUGGUUGGAGUCCGCCAA	1872
2215	AAGCUGAGCAUCCAGCGCG	1626	2215	AAGCUGAGCAUCCAGCGCG	1626	2237	CGCGCUGGAUGCUCAGCUU	1873
2233	GUGCGCGAGGAGGAGUCCGG	1627	2233	GUGCGCGAGGAGGAGUCCGG	1627	2255	CCGCAUCCUCCUCCGCCAC	1874
2251	GGACCGUAUUCUGAGAGCG	1628	2251	GGACCGUAUUCUGAGAGCG	1628	2273	CGCUGCACAGAUACGGUCC	1875
2269	GUGUGCAGACCCAAAGGCGU	1629	2269	GUGUGCAGACCCAAAGGCGU	1629	2291	AGCCCUUGGGUCUCGACAC	1876
2287	UGCGUCAACUCCUCCGCCA	1630	2287	UGCGUCAACUCCUCCGCCA	1630	2309	UGGCGGAGGAGUUGACGCA	1877
2305	AGCGUGGCGGUGGAAGGCU	1631	2305	AGCGUGGCGGUGGAAGGCU	1631	2327	AGCCUCCACGCGCCACGCU	1878
2323	UCCGAGGAUAAGGGAGCA	1632	2323	UCCGAGGAUAAGGGAGCA	1632	2345	UGCUGCCCUUAUCCUCGGA	1879
2341	AUGGAGAUUGUAUCCUUG	1633	2341	AUGGAGAUUGUAUCCUUG	1633	2363	CAAGGAUACAGAUCCCAU	1880
2359	GUCGGUACCGCGUCAUCG	1634	2359	GUCGGUACCGCGUCAUCG	1634	2381	CGAUGACGCCGGUACCGAC	1881
2377	GCUGUCUUCUUCUGGGUCC	1635	2377	GCUGUCUUCUUCUGGGUCC	1635	2399	GGACCCAGAAGAAGACAGC	1882
2395	CUCCUCCUCCUCAUCUUCU	1636	2395	CUCCUCCUCCUCAUCUUCU	1636	2417	AGAAGAUAGGAGGAGGAG	1883
2413	UGUAACAUGAGGAGGCGCG	1637	2413	UGUAACAUGAGGAGGCGCG	1637	2435	CGGCGCUCCUAUGUUAUA	1884
2431	GCCACGCGAGACAUAAGA	1638	2431	GCCACGCGAGACAUAAGA	1638	2453	UCUUGAUGUCUGCGUGGGC	1885
2449	ACGGGCUACCCUGUCCAUA	1639	2449	ACGGGCUACCCUGUCCAUA	1639	2471	UGAUGGACAGGUAGCCCGU	1886
2467	AUCAUGGACCCGCGGAGG	1640	2467	AUCAUGGACCCGCGGAGG	1640	2489	CCUCCCGGGGUCCAUGAU	1887
2485	GUGCCUCUGGAGGAGCAAU	1641	2485	GUGCCUCUGGAGGAGCAAU	1641	2507	AUUGCUCCUCCAGAGGCAC	1888
2503	UGCGAAUACCUUGUCCUACG	1642	2503	UGCGAAUACCUUGUCCUACG	1642	2525	CGUAGGACAGGUUUCGCA	1889
2521	GAUGCCAGCCAGUGGGAU	1643	2521	GAUGCCAGCCAGUGGGAU	1643	2543	AUUCGCCACUGGCUGGCAUC	1890
2539	UUCGCCCGAGAGCGGCU	1644	2539	UUCGCCCGAGAGCGGCU	1644	2561	GCAGCCGCUUCGCGGGGAA	1891
2557	CACCUAGGAGAGUUGCU	1645	2557	CACCUAGGAGAGUUGCU	1645	2579	CGAGCACUCUCCCGCAGGUG	1892
2575	GGCUACGGCGCCUUCGGGA	1646	2575	GGCUACGGCGCCUUCGGGA	1646	2597	UCCCGAAGGCGCGCGUAGCC	1893
2593	AAGGUGGUGGAAGCCUCCG	1647	2593	AAGGUGGUGGAAGCCUCCG	1647	2615	CGGAGGCUUCCACACCCUU	1894
2611	GCUUUCGGCAUCCACAAGG	1648	2611	GCUUUCGGCAUCCACAAGG	1648	2633	CCUUGUGGAUCCCGAAAGC	1895
2629	GGCAGCAGCUGUACACCG	1649	2629	GGCAGCAGCUGUACACCG	1649	2651	CGGUGUCACAGCUGCUGCC	1896
2647	GUGGCCGUGAAAUGCUGA	1650	2647	GUGGCCGUGAAAUGCUGA	1650	2669	UCAGCAUUUUCACGGCCAC	1897
2665	AAAGAGGGCGCCACGGCCA	1651	2665	AAAGAGGGCGCCACGGCCA	1651	2687	UGGCCGUGGCGCCCUUUU	1898
2683	AGCAGCAGCGCGCGCUGA	1652	2683	AGCAGCAGCGCGCGCUGA	1652	2705	UCAGCGCGCGCUGCUGCU	1899
2701	AUGUCGGAGCUCAGAUC	1653	2701	AUGUCGGAGCUCAGAUC	1653	2723	GGAUUUUAGCUCGACAU	1900
2719	CUCAUACAUCCGCAACC	1654	2719	CUCAUACAUCCGCAACC	1654	2741	GGUUGCCGAUGUGAAUGAG	1901
2737	CACCUCAACGUGGUAACC	1655	2737	CACCUCAACGUGGUAACC	1655	2759	GGUUGACACAGUUGAGGUG	1902
2755	CUCCUGGGGCGUGCACCA	1656	2755	CUCCUGGGGCGUGCACCA	1656	2777	UGGUGCACGCCCGGAGGAG	1903
2773	AAGCCGACGGGCCCCCUCA	1657	2773	AAGCCGACGGGCCCCCUCA	1657	2795	UGAGGGGCGCCUGCGGCUU	1904

2791	AUGGUGAUCGUGGAGUUCU	1658	2791	AUGGUGAUCGUGGAGUUCU	1658	2813	AGAACUCCACGAUCACCAU	1905
2809	UGCAAGUACGGCAACCUUCU	1659	2809	UGCAAGUACGGCAACCUUCU	1659	2831	AGAGGUUUGCCGUACUUGCA	1906
2827	UCCAACUUCUUGCGGCCA	1660	2827	UCCAACUUCUUGCGGCCA	1660	2849	UGGCGCGCAGGAAGUUGGA	1907
2845	AAGCGGACGCCUUCAGCC	1661	2845	AAGCGGACGCCUUCAGCC	1661	2867	GGCUGAAGGCGUCCCGCUU	1908
2863	CCUUGCGGAGAGUUCUC	1662	2863	CCUUGCGGAGAGUUCUC	1662	2885	GAGACUUCUCGCGCGAGGG	1909
2881	CCCGAGCAGCGCGGACGCU	1663	2881	CCCGAGCAGCGCGGACGCU	1663	2903	AGCGUCCGCGCUGCUCGGG	1910
2899	UUCGCGCCAUUGGAGC	1664	2899	UUCGCGCCAUUGGAGC	1664	2921	GCUCACCAUUGCGCGGAA	1911
2917	CUCGCCAGGCUUGGAGGA	1665	2917	CUCGCCAGGCUUGGAGGA	1665	2939	UCCGAUCCAGCCUGGCGAG	1912
2935	AGGCGGCCGGGAGCAGCG	1666	2935	AGGCGGCCGGGAGCAGCG	1666	2957	CGCUGUCCCGCGCGGCCU	1913
2953	GACAGGCUUCUUCGCGC	1667	2953	GACAGGCUUCUUCGCGC	1667	2975	GGCGAAGAGACCCUUGUC	1914
2971	CGGUUCUGAAGACCGAGG	1668	2971	CGGUUCUGAAGACCGAGG	1668	2993	CCUCGGUUCUGGAGAACCG	1915
2989	GGCGGAGCGAGGCGGCUU	1669	2989	GGCGGAGCGAGGCGGCUU	1669	3011	AAGCCGCUUGCUCGCGC	1916
3007	UCUCAGACCAAGAACGUG	1670	3007	UCUCAGACCAAGAACGUG	1670	3029	CAGCUUCUUGGUCUGGAGA	1917
3025	GAGGACCUUGGCGUGAGCC	1671	3025	GAGGACCUUGGCGUGAGCC	1671	3047	GGCUCAGCCACAGGUCCUC	1918
3043	CCGUGACCAUGGAAGAUUC	1672	3043	CCGUGACCAUGGAAGAUUC	1672	3065	GAUCUCCAUUGGUCAGCGG	1919
3061	CUUGUGCUACAGCUUC	1673	3061	CUUGUGCUACAGCUUC	1673	3083	GGAAGCUGUAGCAGACAAAG	1920
3079	CAGGUGGCCAGAGGGAUGG	1674	3079	CAGGUGGCCAGAGGGAUGG	1674	3101	CCAUCUCCUUGGCCACCUUG	1921
3097	GAGUCCUGGCUUCCCGAA	1675	3097	GAGUCCUGGCUUCCCGAA	1675	3119	UUCGGGAAGCCAGGAACUUC	1922
3115	AAGUGCAUCCACAGAGACC	1676	3115	AAGUGCAUCCACAGAGACC	1676	3137	GGUCUCUGGGAUGCACUUC	1923
3133	CUGGCUUGCUGGAACAUUC	1677	3133	CUGGCUUGCUGGAACAUUC	1677	3155	GAUUGUCCGAGCAGCCAG	1924
3151	CUGCUGUCGGAAGCGACG	1678	3151	CUGCUGUCGGAAGCGACG	1678	3173	CGUGCUCUCCGACAGCAG	1925
3169	GUGGUGAAGAUUCUGACU	1679	3169	GUGGUGAAGAUUCUGACU	1679	3191	AGUCACAGAUUCUACCCAG	1926
3187	UUUGGCCUUGCCCGGACA	1680	3187	UUUGGCCUUGCCCGGACA	1680	3209	UGUCCCGGGCAAGGCCAAA	1927
3205	AUCUACAAAGACCCCGACU	1681	3205	AUCUACAAAGACCCCGACU	1681	3227	AGUCGGGUCUUGUAGAU	1928
3223	UACGUCGCAAGGGCAGUG	1682	3223	UACGUCGCAAGGGCAGUG	1682	3245	CACUGCCUUGCGGACGUA	1929
3241	GCCCGGCUGCCCUUGAAGU	1683	3241	GCCCGGCUGCCCUUGAAGU	1683	3263	ACUUCAGGGGCGAGCCGGGC	1930
3259	UGGAUGGCCCUUGAAAGCA	1684	3259	UGGAUGGCCCUUGAAAGCA	1684	3281	UGCUUUCAGGGGCCAUCCA	1931
3277	AUCUUCGACAAAGGUGUACA	1685	3277	AUCUUCGACAAAGGUGUACA	1685	3299	UGUACACCUUGUCGAAGAU	1932
3295	ACCACGCAGAGUGAGUGU	1686	3295	ACCACGCAGAGUGAGUGU	1686	3317	ACAGUCACUCUGCGUGGU	1933
3313	UGGUCCUUGGGGUGCUUC	1687	3313	UGGUCCUUGGGGUGCUUC	1687	3335	GAAGCACCCCAAAGGACCA	1934
3331	CUCUGGGAGAUUCUUCUC	1688	3331	CUCUGGGAGAUUCUUCUC	1688	3353	GAGAGAAGAUUCUCCAGAG	1935
3349	CUGGGGCGCUCCCGUACC	1689	3349	CUGGGGCGCUCCCGUACC	1689	3371	GGUACGGGAGGGCCCCCAG	1936
3367	CCUGGGGUGCAGAUCAUUG	1690	3367	CCUGGGGUGCAGAUCAUUG	1690	3389	CAUUGAUUCGACCCCGAGG	1937

3385	GAGGAGUUCUGCCAGCGG	1691	3385	GAGGAGUUCUGCCAGCGG	1691	3407	CGCGUUGGCAGAACUCCUC	1938
3403	GUGAGAGACGGCACAAGGA	1692	3403	GUGAGAGACGGCACAAGGA	1692	3425	UCCUUGUGCCGUCUCAC	1939
3421	AUGAGGGCCCCGGAGCUGG	1693	3421	AUGAGGGCCCCGGAGCUGG	1693	3443	CCAGUCCGGGGCCUCAU	1940
3439	GCCACUCCCGCCAUACGCC	1694	3439	GCCACUCCCGCCAUACGCC	1694	3461	GGCGUUGGGGGGAGUGGC	1941
3457	CACAUCAUCUGAACUCU	1695	3457	CACAUCAUCUGAACUCU	1695	3479	AGCAGUUCAGCAUGAUG	1942
3475	UGGUCGGAGACCCCAAGG	1696	3475	UGGUCGGAGACCCCAAGG	1696	3497	CCUUGGGUUCUCCGGACCA	1943
3493	GCGAGACCUCAUUCUGG	1697	3493	GCGAGACCUCAUUCUGG	1697	3515	CCGAGAAUGCAGGUCUCG	1944
3511	GACUUGGAGAUCCUGG	1698	3511	GACUUGGAGAUCCUGG	1698	3533	CCAGGAUUCUCCACCAAGGUC	1945
3529	GGGACCUUGUCCAGGGCA	1699	3529	GGGACCUUGUCCAGGGCA	1699	3551	UGCCUUGGAGCAGGUCCCC	1946
3547	AGGGCCUUGCAAGAGGAAG	1700	3547	AGGGCCUUGCAAGAGGAAG	1700	3569	CUUCCUUCUGCAGGGCCCU	1947
3565	GAGGAGUUCUGCAUGGCC	1701	3565	GAGGAGUUCUGCAUGGCC	1701	3587	GGGCCAUGCAGACCUCCUC	1948
3583	CCGCGCAGCUCUCAGAGCU	1702	3583	CCGCGCAGCUCUCAGAGCU	1702	3605	AGCUCUGAGAGCUCGCGG	1949
3601	UCAGAAGAGGGCAGCUUCU	1703	3601	UCAGAAGAGGGCAGCUUCU	1703	3623	AGAAUGGCCUUCUUGA	1950
3619	UCGCAGGUGUCCACCAUGG	1704	3619	UCGCAGGUGUCCACCAUGG	1704	3641	CCAUGGUGGACACCUUGCGA	1951
3637	GCCUACACAUCCGCCAGG	1705	3637	GCCUACACAUCCGCCAGG	1705	3659	CCUGGGCAUGUGUAGGGC	1952
3655	GCUGACGUCAGGACAGCC	1706	3655	GCUGACGUCAGGACAGCC	1706	3677	GGCUGUCCUCCAGCGUCAGC	1953
3673	CCGCCAAGCCUGCAGCGCC	1707	3673	CCGCCAAGCCUGCAGCGCC	1707	3695	GGCGCUGCAGGCUUGGCGG	1954
3691	CACAGCCUGGCCGCCAGGU	1708	3691	CACAGCCUGGCCGCCAGGU	1708	3713	ACUUGGGCCAGGCUUG	1955
3709	UAUUACAACUGGGUGUCCU	1709	3709	UAUUACAACUGGGUGUCCU	1709	3731	AGGACACCCAGUUGUAAUA	1956
3727	UUUCCCGGUGCCUGGCCA	1710	3727	UUUCCCGGUGCCUGGCCA	1710	3749	UGGCCAGGACCCCGGAAA	1957
3745	AGAGGGGCUAGAGACCCUG	1711	3745	AGAGGGGCUAGAGACCCUG	1711	3767	CACGGGUCUCAGCCCCUCU	1958
3763	GGUUCUCCAGGAUGAAGA	1712	3763	GGUUCUCCAGGAUGAAGA	1712	3785	UCUUAUCCUGGAGGAACC	1959
3781	ACAUUUGAGGAUUUCCCA	1713	3781	ACAUUUGAGGAUUUCCCA	1713	3803	UGGGAAUUCUCAAUUGU	1960
3799	AUGACCCCAACGACCUACA	1714	3799	AUGACCCCAACGACCUACA	1714	3821	UGUAGGUCUUGGGGUCAU	1961
3817	AAAGGCUCUGUGGACAACC	1715	3817	AAAGGCUCUGUGGACAACC	1715	3839	GGUUGUCCACAGAGCCUUU	1962
3835	CAGACAGACAGUGGGAUUG	1716	3835	CAGACAGACAGUGGGAUUG	1716	3857	CCAUCCACUGUCUGUCUG	1963
3853	GUGCUGGCCUCCGGAGGAGU	1717	3853	GUGCUGGCCUCCGGAGGAGU	1717	3875	ACUCCUCCGAGGCCAGCAC	1964
3871	UUUGAGCAGAUAGAGAGCA	1718	3871	UUUGAGCAGAUAGAGAGCA	1718	3893	UGCUCUCUUCUGCUCAAA	1965
3889	AGGCAUAGACAAGAAAGCG	1719	3889	AGGCAUAGACAAGAAAGCG	1719	3911	CGCUUUCUUGUCUAUGCCU	1966
3907	GGCUACAGGUAUCUGAAGC	1720	3907	GGCUACAGGUAUCUGAAGC	1720	3929	GCUUCAGCUACCUAGGCC	1967
3925	CAGAGAGAGAGGAGGAGC	1721	3925	CAGAGAGAGAGGAGGAGC	1721	3947	GCUGCCUUCUCUCUCUCUG	1968
3943	CAUACGUCAGCAUUUUCUU	1722	3943	CAUACGUCAGCAUUUUCUU	1722	3965	AAGAAAUGCUGACGUAUG	1969
3961	UCUCUGCACUUAUAAAGAAA	1723	3961	UCUCUGCACUUAUAAAGAAA	1723	3983	UUUCUUAUAAAGUGCAGAGA	1970

3979	AGAUCAAAGACUUUAAGAC	1724	3979	AGAUCAAAGACUUUAAGAC	1724	4001	GUCUUAAAGUCUUUGAUCU	1871
3997	CUUUCGCUAUUUUCUUCUAC	1725	3997	CUUUCGCUAUUUUCUUCUAC	1725	4019	GUAGAAGAAUAAGCGAAAG	1872
4015	CUGCUAUUCUACUACAAACU	1726	4015	CUGCUAUUCUACUACAAACU	1726	4037	AGUUUGUAGUAGAUAGCAAG	1873
4033	UUCAAAGAGGAACCCAGGAG	1727	4033	UUCAAAGAGGAACCCAGGAG	1727	4055	CUCUGGUUCCUUCUUUGAA	1874
4051	GGACAAGAGGAGCAUGAAA	1728	4051	GGACAAGAGGAGCAUGAAA	1728	4073	UUUCAUGCUCUUCUUGUCC	1875
4069	AGUGGACAAGGAGUGUGAC	1729	4069	AGUGGACAAGGAGUGUGAC	1729	4091	GUACACUCCUUGUCCACU	1876
4087	CCACUGAAGCACACAGGG	1730	4087	CCACUGAAGCACACAGGG	1730	4109	CCUGUGGUGCUUCAGUGG	1877
4105	GAGGGUUAAGCCUCCGGA	1731	4105	GAGGGUUAAGCCUCCGGA	1731	4127	UCCGAGGCCUUAACCCUUC	1878
4123	AUGACUGCGGCAGGCCUG	1732	4123	AUGACUGCGGCAGGCCUG	1732	4145	CAGGCUGCCCGCAGUCAU	1879
4141	GGAUAAUUAUCCAGCCUCC	1733	4141	GGAUAAUUAUCCAGCCUCC	1733	4163	GGGAGGCGUGGAUUAUCC	1880
4159	CACAAGAGCUGGUGGAGC	1734	4159	CACAAGAGCUGGUGGAGC	1734	4181	GCUCCACCCAGCUUCUUGUG	1881
4177	CAGAGUUAUCCUGACUCC	1735	4177	CAGAGUUAUCCUGACUCC	1735	4199	GGAGUCAGGGAACACUCUG	1882
4195	CUCCAAGGAAAGGGAGACG	1736	4195	CUCCAAGGAAAGGGAGACG	1736	4217	CGUCUCCCUUCCUUGGAG	1883
4213	GCCCUUUAUGGUGUCUG	1737	4213	GCCCUUUAUGGUGUCUG	1737	4235	CAGCAGACCAUGAAAGGC	1884
4231	GAGUAAACAGGUGCCUCC	1738	4231	GAGUAAACAGGUGCCUCC	1738	4253	GGGAAGGCACCCUGUACUC	1885
4249	CAGACACUGGCGUUAUCG	1739	4249	CAGACACUGGCGUUAUCG	1739	4271	GCAGUAAAGCCAGUGUCUG	1886
4267	CUUGACCAAGAGCCCUCA	1740	4267	CUUGACCAAGAGCCCUCA	1740	4289	UGAGGCGUUCUUGGUAAG	1887
4285	AAGCGCCCUUAUGCCAGC	1741	4285	AAGCGCCCUUAUGCCAGC	1741	4307	GCUGGCAUAAAGGCCGCUU	1888
4303	CGUGACAGAGGCGUCACCU	1742	4303	CGUGACAGAGGCGUCACCU	1742	4325	AGUGAGCCCUUCUGUCACG	1889
4321	UCUUGCCUUCUAGGUCACU	1743	4321	UCUUGCCUUCUAGGUCACU	1743	4343	AGUGACCUAAGAGGCAAGA	1890
4339	UUCUCACAUAUCCCUUCA	1744	4339	UUCUCACAUAUCCCUUCA	1744	4361	UGAAGGGACAUUGUGAGAA	1891
4357	AGCACCUAGCCUUGGCC	1745	4357	AGCACCUAGCCUUGGCC	1745	4379	GGGACAGGGUCCAGGUGCU	1892
4375	CGCCGAUUAUCCUUGGUA	1746	4375	CGCCGAUUAUCCUUGGUA	1746	4397	UACCAAGGAUAUCCGGCG	1893
4393	AAUAUGAGUAUAUACAUCAA	1747	4393	AAUAUGAGUAUAUACAUCAA	1747	4415	UUGAUGUAUUAUCUUAUU	1894
4411	AAGAGUAGUAUUAAGCU	1748	4411	AAGAGUAGUAUUAAGCU	1748	4433	AGCUUUAUAUACUACUCUU	1895
4429	UAAUUAUAUUAUUAUAA	1749	4429	UAAUUAUAUUAUUAUAA	1749	4451	UUUAUAAACAUUAUUAUUA	1896

The 3'-ends of the Upper sequence and the Lower sequence of the siNA construct can include an overhang sequence, for example about 1, 2, 3, or 4 nucleotides in length, preferably 2 nucleotides in length, wherein the overhanging sequence of the lower sequence is optionally complementary to a portion of the target sequence. The upper sequence is also referred to as the sense strand, whereas the lower sequence is also referred to as the antisense strand. The upper and lower sequences in the Table can further comprise a chemical modification having Formulae I-VII or any combination thereof.



Table III: VEGF and VEGFr Synthetic Modified siNA constructs

Target Pos	Target	Seq ID	Aliases	Sequence	Seq ID
296	GCUGUCUGCUUCUCACAGGAUCU	1997	FLT1:298U21 siRNA sense	UGUCUGCUUCUCACAGGAUTT	2020
1954	GAAGGAGAGGACCUGAAACUGUC	1998	FLT1:1956U21 siRNA sense	AGGAGAGGACCUGAAACUGTT	2021
1955	AAGGAGAGGACCUGAAACUGUCU	1999	FLT1:1957U21 siRNA sense	GGAGAGGACCUGAAACUGUTT	2022
2785	GCAUUUGGCAUUUAGAAAUACACC	2000	FLT1:2787U21 siRNA sense	AUUUGGCAUUUAGAAAUACATT	2023
296	GCUGUCUGCUUCUCACAGGAUCU	1997	FLT1:316L21 siRNA (298C) antisense	AUCCUGUGAGAAGCAGACATT	2024
1954	GAAGGAGAGGACCUGAAACUGUC	1998	FLT1:1974L21 siRNA (1956C) antisense	CAGUUUCAGGUCCUCUCUCCUTT	2025
1955	AAGGAGAGGACCUGAAACUGUCU	1999	FLT1:1975L21 siRNA (1957C) antisense	ACAGUUUCAGGUCCUCUCUCCTT	2026
2785	GCAUUUGGCAUUUAGAAAUACACC	2000	FLT1:2805L21 siRNA (2787C) antisense	UGAUUUUCUUAAUAGCCAAAUUTT	2027
296	GCUGUCUGCUUCUCACAGGAUCU	1997	FLT1:298U21 siRNA stab04 sense	B UGUUCGUUUCUcAcAGGAUTT B	2028
1954	GAAGGAGAGGACCUGAAACUGUC	1998	FLT1:1956U21 siRNA stab04 sense	B AGGAGAGGAGGACCUGAAACUGTT B	2029
1955	AAGGAGAGGACCUGAAACUGUCU	1999	FLT1:1957U21 siRNA stab04 sense	B GGAGAGGAGGACCUGAAACUGTT B	2030
2785	GCAUUUGGCAUUUAGAAAUACACC	2000	FLT1:2787U21 siRNA stab04 sense	B AUUUUGGCAUUUAGAAAUACATT B	2031
296	GCUGUCUGCUUCUCACAGGAUCU	1997	FLT1:316L21 siRNA (298C) stab05 antisense	AuccuGuGAGAAAGcAGAcATsT	2032
1954	GAAGGAGAGGACCUGAAACUGUC	1998	FLT1:1974L21 siRNA (1956C) stab05 antisense	cAGUUUCAGGUCCUCUCUCCTsT	2033
1955	AAGGAGAGGACCUGAAACUGUCU	1999	FLT1:1975L21 siRNA (1957C) stab05 antisense	AcAGUUUCAGGUCCUCUCCTsT	2034
2785	GCAUUUGGCAUUUAGAAAUACACC	2000	FLT1:2805L21 siRNA (2787C) stab05 antisense	uGAUUUUUUAAUGccAAAUtsT	2035
296	GCUGUCUGCUUCUCACAGGAUCU	1997	FLT1:298U21 siRNA stab07 sense	B UGUUCGUUUCUcAcAGGAUTT B	2036
1954	GAAGGAGAGGACCUGAAACUGUC	1998	FLT1:1956U21 siRNA stab07 sense	B AGGAGAGGAGGACCUGAAACUGTT B	2037
1955	AAGGAGAGGACCUGAAACUGUCU	1999	FLT1:1957U21 siRNA stab07 sense	B GGAGAGGAGGACCUGAAACUGTT B	2038
2785	GCAUUUGGCAUUUAGAAAUACACC	2000	FLT1:2787U21 siRNA stab07 sense	B AUUUUGGCAUUUAGAAAUACATT B	2039
296	GCUGUCUGCUUCUCACAGGAUCU	1997	FLT1:316L21 siRNA (298C) stab11 antisense	AuccuGuGAGAAAGcAGAcATsT	2040
1954	GAAGGAGAGGACCUGAAACUGUC	1998	FLT1:1974L21 siRNA (1956C) stab11 antisense	cAGUUUCAGGUCCUCUCCTsT	2041
1955	AAGGAGAGGACCUGAAACUGUCU	1999	FLT1:1975L21 siRNA (1957C) stab11 antisense	AcAGUUUCAGGUCCUCUCCTsT	2042
2785	GCAUUUGGCAUUUAGAAAUACACC	2000	FLT1:2805L21 siRNA (2787C) stab11 antisense	uGAUUUUUUAAUGccAAAUtsT	2043



## VEGFR1

Target	SeqID	RPI#	Alias	Sequence	SeqID
AACUGAGUUUAAAAGGCCACCCAG	2009	29694	FLT1:349U21 siRNA stab01 sense	CsUsGsAsGsUUUAAAAGGCCACCCCTsT	2092
AACAACCCACAAAUAACAACAAGA	2010	29695	FLT1:2340U21 siRNA stab01 sense	CsAsAsCsCsACAAAUAACAACAATsT	2093
AGCCUGGAAAGAAUCAAACCCUU	2011	29696	FLT1:3912U21 siRNA stab01 sense	CsCsUsGsGsAAAGAAUCAAACCCCTsT	2094
AAGCAAGGAGGGCCUCUGAUGGU	2012	29697	FLT1:2949U21 siRNA stab01 sense	GsCsAsAsGsGAGGGCCUCUGAUGTsT	2095
AACUGAGUUUAAAAGGCCACCCAG	2009	29698	FLT1:369L21 siRNA (349C) stab01 sense	GsGsGsUsGsCCUUUUAAAACUCAGTsT	2096
AACAACCCACAAAUAACAACAAGA	2010	29699	FLT1:2358L21 siRNA (2340C) stab01 sense	UsUsGsUsUsGUUAUUUUGUGGUUGTsT	2097
AGCCUGGAAAGAAUCAAACCCUU	2011	29700	FLT1:3932L21 siRNA (3912C) stab01 sense	GsGsUsUsUsUGAUUUCUUUCCAGGTsT	2098
AAGCAAGGAGGGCCUCUGAUGGU	2012	29701	FLT1:2969L21 siRNA (2949C) stab01 sense	CsAsUsCsAsGAGGCCUCCUUGCTsT	2099
AACUGAGUUUAAAAGGCCACCCAG	2009	29702	FLT1:349U21 siRNA stab03 sense	csUsGsAsGuuuAAAAGGcAcscscsTsT	2100
AACAACCCACAAAUAACAACAAGA	2010	29703	FLT1:2340U21 siRNA stab03 sense	csAsAsCscAcAAAuuAcaAacsAsTsT	2101
AGCCUGGAAAGAAUCAAACCCUU	2011	29704	FLT1:3912U21 siRNA stab03 sense	csCsUsCsGAAAGAAuAucAAAAsCscsTsT	2102
AAGCAAGGAGGGCCUCUGAUGGU	2012	29705	FLT1:2949U21 siRNA stab03 sense	GscsAsAsGGAGGGccucuuGAsusGsTsT	2103
AACUGAGUUUAAAAGGCCACCCAG	2009	29706	FLT1:369L21 siRNA (349C) stab02 antisense	GsGsGsUsGsCsCsUsUsUsAsAsAsCsUsCsAsGsTsT	2104
AACAACCCACAAAUAACAACAAGA	2010	29707	FLT1:2358L21 siRNA (2340C) stab02 antisense	UsUsGsUsUsGsUsAsUsUsUsGsUsGsGsUsUsGsTsT	2105
AGCCUGGAAAGAAUCAAACCCUU	2011	29708	FLT1:3932L21 siRNA (3912C) stab02 antisense	GsGsUsUsUsUsGsAsUsUsCsUsUsUsCsCsAsGsGsTsT	2106
AAGCAAGGAGGGCCUCUGAUGGU	2012	29709	FLT1:2969L21 siRNA (2949C) stab02 antisense	CsAsUsCsAsGsAsGsCsCsCsUsCsCsUsUsGsCsTsT	2107
AACAACCCACAAAUAACAACAAGA	2010	29981	FLT1:2340U21 siRNA Native sense	CAACCCACAAAUAACAACAAGA	2108
AACAACCCACAAAUAACAACAAGA	2010	29982	FLT1:2358L21 siRNA (2340C) Native antisense	UUUGUUGUAUUUUUGUGGUUGUU	2109
AACAACCCACAAAUAACAACAAGA	2010	29983	FLT1:2342U21 siRNA stab01 inv	AsAsCsAsAsCAUAAAACACCAACTsT	2110
AACAACCCACAAAUAACAACAAGA	2010	29984	FLT1:2358L21 siRNA (2340C) stab01 inv	GsUsUsGsUGUUUUUAUGUUGUUTsT	2111
AACAACCCACAAAUAACAACAAGA	2010	29985	FLT1:2342U21 siRNA stab03 inv	AsAsCsAsAcAuAAAACacccAsAscsTsT	2112
AACAACCCACAAAUAACAACAAGA	2010	29986	FLT1:2358L21 siRNA (2340C) stab02 inv	GsUsUsGsGsUsGsUsUsUsAsUsGsUsGsUsUsTsT	2113

AACAACCCACAAAAUACAACAAGA	2010	29987	FLT1:2340U21 siRNA sense	2114	AGAACAAACAUAACCAAC
AACAACCCACAAAAUACAACAAGA	2010	29988	FLT1:2358L21 siRNA (2340C) Inv Native	2115	UUUGUUGGUGUUUAUGUUUU
AACAACCCACAAAAUACAACAAGA	2010	30075	FLT1:2340U21 siRNA sense	2116	CAACCACAAAAUACAACAATT
AACAACCCACAAAAUACAACAAGA	2010	30076	FLT1:2358L21 siRNA (2340C) antisense	2117	UUUGUUAUUUUUGUGGUUGTT
AACAACCCACAAAAUACAACAAGA	2010	30077	FLT1:2342U21 siRNA Inv	2118	AGAACAAACAUAACCAACATT
AACAACCCACAAAAUACAACAAGA	2010	30078	FLT1:2358L21 siRNA (2340C) Inv	2119	UUUGUUGGUGUUUAUGUUUGTT
AACAACCCACAAAAUACAACAAGA	2010	30187	FLT1:2358L21 siRNA (2340C) 2'-F U.C antisense	2120	uuGuuGuAuuuuuGuGGuuGTT
AACAACCCACAAAAUACAACAAGA	2010	30190	FLT1:2358L21 siRNA (2340C) X = nitroindole antisense	2121	uuGuuGuAuuuuuGuGGuuGXX
AACAACCCACAAAAUACAACAAGA	2010	30193	FLT1:2340U21 siRNA (2340C) Z = nitroprole antisense	2122	uuGuuGuAuuuuuGuGGuuGZZ
AACAACCCACAAAAUACAACAAGA	2010	30196	FLT1:2340U21 siRNA sense IB caps w/2'FY's sense	2123	B cAAccAcAAAAuAcAAcAAATT B
AACAACCCACAAAAUACAACAAGA	2010	30199	FLT1:2340U21 siRNA sense IB caps sense	2124	cAAccAcAAAAuAcAAcAAATT
AACAACCCACAAAAUACAACAAGA	2010	30340	FLT1:2358L21 siRNA (2340C) X = 3'dT antisense	2125	uuGuuGuAuuuuuGuGGuuGTX
AACAACCCACAAAAUACAACAAGA	2010	30341	FLT1:2358L21 siRNA (2340C) X = glycyl antisense	2126	uuGuuGuAuuuuuGuGGuuGTX
AACAACCCACAAAAUACAACAAGA	2010	30342	FLT1:2358L21 siRNA (2340C) U = 3'OMeU antisense	2127	uuGuuGuAuuuuuGuGGuuGTU
AACAACCCACAAAAUACAACAAGA	2010	30343	FLT1:2358L21 siRNA (2340C) t = L- dT antisense	2128	uuGuuGuAuuuuuGuGGuuGTt
AACAACCCACAAAAUACAACAAGA	2010	30344	FLT1:2358L21 siRNA (2340C) u = L-rU antisense	2129	uuGuuGuAuuuuuGuGGuuGTu
AACAACCCACAAAAUACAACAAGA	2010	30345	FLT1:2358L21 siRNA (2340C) D = IdT antisense	2130	uuGuuGuAuuuuuGuGGuuGTD
AACAACCCACAAAAUACAACAAGA	2010	30346	FLT1:2358L21 siRNA (2340C) X = 3'dT antisense	2131	uuGuuGuAuuuuuGuGGuuGXT
AACAACCCACAAAAUACAACAAGA	2010	30416	FLT1:2358L21 siRNA (2340C) TsT antisense	2132	uuGuuGuAuuuuuGuGGuuGTsT
AACAACCCACAAAAUACAACAAGA	2013	30777	FLT1:184U21 siRNA stab04 sense	2133	B GuGuAAGGAGUGGAccAucTT B
UCGUGUAGGAGUGGACCAUCAU	2014	30778	FLT1:3503U21 siRNA stab04 sense	2134	B AcGGAGuAuuuGuGuGGGATT B
UUACGGAGUUAUUGCUGUGGAAA	2015	30779	FLT1:4715U21 siRNA stab04 sense	2135	B GcAGGccuuAAGAcuGuGATT B
UAGCAGGCCUUAAGACAUGUGAGG	2016	30780	FLT1:4753U21 siRNA stab04 sense	2136	B cAAAAAGcAAAGGAGAGAAAAATT B

UCGUGUAAGGAGUGGACCAUCAU	2013	30781	FLT1:1202L21 siRNA (1184C) stab05 antisense	GAUGGuccAcuccuuAcAcTsT	2137
UUACGGAGUAUUGCUGUGGAAA	2014	30782	FLT1:3521L21 siRNA (3503C) stab05 antisense	uccAcAcAGcAAuAcuccGuTsT	2138
UAGCAGGCCUUAAGACAUGUGAGG	2015	30783	FLT1:4733L21 siRNA (4715C) stab05 antisense	ucAcAuGuccuuAGGccuGcTsT	2139
AGCAAAAAGCAAGGGAGAAAAGA	2016	30784	FLT1:4771L21 siRNA (4753C) stab05 antisense	uuuuucccuuGcuuuuuGTsT	2140
AACAACACAAAAUACAACAAGA	2010	30955	FLT1:2340U21 siRNA stab07 sense	B cAAccAcAAAAuAcAAcAAATT B	2141
AACAACACAAAAUACAACAAGA	2010	30956	FLT1:2358L21 siRNA (2340C) stab08 antisense	uuGuuGuAuuuuuGuGuuGTsT	2142
AACAACACAAAAUACAACAAGA	2010	30963	FLT1:2340U21 siRNA inv	AACAACAUA AAAACACCAACTT	2143
AACAACACAAAAUACAACAAGA	2010	30984	FLT1:2358L21 siRNA (2340C) inv	GUUGGUGUUUAUGUUGUUTT	2144
AACAACACAAAAUACAACAAGA	2010	30965	FLT1:2340U21 siRNA stab04 inv	B AAcAAcAuAAAAAcAcAAcTT B	2145
AACAACACAAAAUACAACAAGA	2010	30966	FLT1:2358L21 siRNA (2340C) stab05 inv	GuuGGUGuuuuAuGuuGuuTsT	2146
AACAACACAAAAUACAACAAGA	2010	30967	FLT1:2340U21 siRNA stab07 inv	B AAcAAcAuAAAAAcAcAAcTT B	2147
AACAACACAAAAUACAACAAGA	2010	30968	FLT1:2358L21 siRNA (2340C) stab08 inv	GuuGGUGuuuuAuGuuGuuTsT	2148
AACUGAGUUUAAAAGGCACCCAG	2009	31182	FLT1:349U21 siRNA TT sense FLT1:2949U21 siRNA TT	CUGAGUUUAAAAGGCACCCCTT	2149
AAGCAAGGAGGGCCUCUGAUGGU	2012	31183	antisense	GCAAGGAGGGCCUCUGAUGTT	2150
AGCCUGGAAAGAAUCAAACCCUU	2011	31184	FLT1:3912U21 siRNA TT sense	CCUGGAAAGAAUCAAACCCCTT	2151
AACUGAGUUUAAAAGGCACCCAG	2009	31185	antisense	GGGUGCCUUUUAAAACUCAGTT	2152
AAGCAAGGAGGGCCUCUGAUGGU	2012	31186	FLT1:2987L21 siRNA (2949C) TT sense	CAUCAGAGGGCCUCUUGCTT	2153
AGCCUGGAAAGAAUCAAACCCUU	2011	31187	FLT1:3930L21 siRNA (3912C) TT antisense	GGUUUUGAUUCUUUCCAGGTT	2154
AACUGAGUUUAAAAGGCACCCAG	2009	31188	FLT1:349U21 siRNA stab04 sense	B cuGAGuuuuAAAAAGGcAcctTT B	2155
AAGCAAGGAGGGCCUCUGAUGGU	2012	31189	antisense	B GcAAGGAGGGccuGcuGauGTT B	2156
AGCCUGGAAAGAAUCAAACCCUU	2011	31190	FLT1:3912U21 siRNA stab04 sense	B ccuGGAAGAAuAcAAAAcTT B	2157
AACUGAGUUUAAAAGGCACCCAG	2009	31191	FLT1:367L21 siRNA (349C) stab05 antisense	GGGUgGccuuuuAAAAcucAGTsT	2158
AAGCAAGGAGGGCCUCUGAUGGU	2012	31192	FLT1:2987L21 siRNA (2949C) stab05 antisense	cAucAGAGGccuGcuGcTsT	2159
AGCCUGGAAAGAAUCAAACCCUU	2011	31193	FLT1:3930L21 siRNA (3912C) stab05 antisense	GGuuuuuGAuuuuuuccAGGTsT	2160

AACUGAGUUUAAAAGGCCACCCAG	2009	31194	FLT1:349U21 siRNA stab07 sense	B cuGAGUUUAAAAGGcAaccTT B	2161
AAGCAAGGAGGGCCUCUGAUGGU	2012	31195	FLT1:2949U21 siRNA stab07 sense	B GcAAGGAGGGGccuicuGauGTT B	2162
AGCCUGGAAAGAAUCAAACCCUU	2011	31196	FLT1:3912U21 siRNA stab07 sense	B ccuGGAAAAGAAucAAAaccTT B	2163
AACUGAGUUUAAAAGGCCACCCAG	2009	31197	FLT1:367L21 siRNA (349C) stab08 antisense	GGGuGccuuuuuAAAacucAGTsT	2164
AAGCAAGGAGGGCCUCUGAUGGU	2012	31198	FLT1:2967L21 siRNA (2949C) stab08 antisense	cAucAGAGGGccuccuuGcTsT	2165
AGCCUGGAAAGAAUCAAACCCUU	2011	31199	FLT1:3930L21 siRNA (3912C) stab08 antisense	GGUUuuGAAuuuuuccAGGTsT	2166
AACUGAGUUUAAAAGGCCACCCAG	2009	31200	FLT1:349U21 siRNA inv TT	CCCACGGAAAAUUUGAGUCIT	2167
AAGCAAGGAGGGCCUCUGAUGGU	2012	31201	FLT1:2949U21 siRNA inv TT	GUAGUCUCCGGGAGGAACGTT	2168
AGCCUGGAAAGAAUCAAACCCUU	2011	31202	FLT1:3912U21 siRNA inv TT	CCAAAACUAAAGAAAGGUCCIT	2169
AACUGAGUUUAAAAGGCCACCCAG	2009	31203	FLT1:367L21 siRNA (349C) inv TT	GACUCAAAUUUUCGUGGGTT	2170
AAGCAAGGAGGGCCUCUGAUGGU	2012	31204	FLT1:2967L21 siRNA (2949C) inv TT	CGUUCUCCCGGAGACUACTT	2171
AGCCUGGAAAGAAUCAAACCCUU	2011	31205	FLT1:3930L21 siRNA (3912C) inv TT	GGACCUUUUUAUUUUUGGTT	2172
AACUGAGUUUAAAAGGCCACCCAG	2009	31206	FLT1:349U21 siRNA stab04 inv	B cccAcGGAAAAuuuGAGucTT B	2173
AAGCAAGGAGGGCCUCUGAUGGU	2012	31207	FLT1:2949U21 siRNA stab04 inv	B GuAGucuccGGGAGGAACGTT B	2174
AGCCUGGAAAGAAUCAAACCCUU	2011	31208	FLT1:3912U21 siRNA stab04 inv	B ccAAAAcuuAAGAAAGGUccTT B	2175
AACUGAGUUUAAAAGGCCACCCAG	2009	31209	FLT1:367L21 siRNA (349C) stab05 inv	GAcucAAAuuuuuccGuGGGTsT	2176
AAGCAAGGAGGGCCUCUGAUGGU	2012	31210	FLT1:2967L21 siRNA (2949C) stab05 inv	cGuuuccuccGGAGAcuAcTsT	2177
AGCCUGGAAAGAAUCAAACCCUU	2011	31211	FLT1:3930L21 siRNA (3912C) stab05 inv	GGAccuuuuuuAGuuuuGGTsT	2178
AACUGAGUUUAAAAGGCCACCCAG	2009	31212	FLT1:349U21 siRNA stab07 inv	B cccAcGGAAAAuuuGAGucTT B	2179
AAGCAAGGAGGGCCUCUGAUGGU	2012	31213	FLT1:2949U21 siRNA stab07 inv	B GuAGucuccGGGAGGAACGTT B	2180
AGCCUGGAAAGAAUCAAACCCUU	2011	31214	FLT1:3912U21 siRNA stab07 inv	B ccAAAAcuuAAGAAAGGUccTT B	2181
AACUGAGUUUAAAAGGCCACCCAG	2009	31215	FLT1:367L21 siRNA (349C) stab08 inv	GAcucAAAuuuuuccGuGGGTsT	2182
AAGCAAGGAGGGCCUCUGAUGGU	2012	31216	FLT1:2967L21 siRNA (2949C) stab08 inv	cGuuuccuccGGAGAcuAcTsT	2183
AGCCUGGAAAGAAUCAAACCCUU	2011	31217	FLT1:3930L21 siRNA (3912C) stab08 inv	GGAccuuuuuuAGuuuuGGTsT	2184
AACUGAGUUUAAAAGGCCACCCAG	2009	31270	FLT1:349U21 siRNA stab09 sense	B cuGAGUUUAAAAGGCCACCCIT B	2185
AAGCAAGGAGGGCCUCUGAUGGU	2012	31271	FLT1:2949U21 siRNA stab09	B GCAAGGAGGGCCUCUGAUGTT B	2186

AGCCUGGAAAGAAUCAAACCCUU	2011	31272	sense FLT1:3912U21 siRNA stab09 sense	B CCUGGAAAGAAUCAAACCTT B	2187
AACUGAGUUUAAAAGGCACCCAG	2009	31273	FLT1:367L21 siRNA (349C) stab10 antisense	GGGUGCCUUUUAAAACUCAGTsT	2188
AAGCAAGGAGGGCCUCUGAUGGU	2012	31274	FLT1:2967L21 siRNA (2949C) stab10 antisense	CAUCAGAGGCCCCUCCUUGCTsT	2189
AGCCUGGAAAGAAUCAAACCCUU	2011	31275	FLT1:3930L21 siRNA (3912C) stab10 antisense	GGUUUUUGAUUCUUUCCAGGTsT	2190
AACUGAGUUUAAAAGGCACCCAG	2009	31276	FLT1:349U21 siRNA stab09 inv	B CCCACGGAUUUUUGAGUCTT B	2191
AAGCAAGGAGGGCCUCUGAUGGU	2012	31277	FLT1:2949U21 siRNA stab08 inv	B GUAGUCUCCGGGAGGAACGTT B	2192
AGCCUGGAAAGAAUCAAACCCUU	2011	31278	FLT1:3912U21 siRNA stab09 inv	B CCAAAACUAAAGAAAGGUCCTT B	2193
AACUGAGUUUAAAAGGCACCCAG	2009	31279	FLT1:367L21 siRNA (349C) stab10 inv	GACUCAAAUUUUUCCGUGGTsT	2194
AAGCAAGGAGGGCCUCUGAUGGU	2012	31280	FLT1:2967L21 siRNA (2949C) stab10 inv	CGUUCCUCCCCGGAGACUACTsT	2195
AGCCUGGAAAGAAUCAAACCCUU	2011	31281	FLT1:3930L21 siRNA (3912C) stab10 inv	GGACCUUUUCUUAGUUUUUGGTsT	2196
AACAACCCACAAAUAACAACAAGA	2010	31424	FLT1:2358L21 siRNA (2340C) stab11 X = 3'-BrdU antisense	uuGuuGuAuuuuuuGuGGuuGXsX	2197
AAGCAAGGAGGGCCUCUGAUGGU	2012	31425	FLT1:2967L21 siRNA (2949C) stab11 X = 3'-BrdU sense	cAucAGAGGGccuccuuuGcXsX	2198
AACAACCCACAAAUAACAACAAGA	2010	31442	FLT1:2358L21 siRNA (2340C) stab11 X = 3'-BrdU antisense	uuGuuGuAuuuuuuGuGGuuGXsT	2199
AAGCAAGGAGGGCCUCUGAUGGU	2012	31443	FLT1:2967L21 siRNA (2949C) stab11 X = 3'-BrdU sense	cAucAGAGGGccuccuuuGcXsT	2200
AACAACCCACAAAUAACAACAAGA	2010	31448	FLT1:2340U21 siRNA stab09 sense	B CAACCCACAAAUAACAACAATT B	2201
AACAACCCACAAAUAACAACAAGA	2010	31450	FLT1:2340U21 siRNA inv stab09 sense	B AACACAUAAAACACCAACTT B	2202
AACAACCCACAAAUAACAACAAGA	2010	31451	FLT1:2358L21 siRNA (2340C) stab10 antisense	UUUGUUUAUUUUUGUGGUUGTsT	2203
AACAACCCACAAAUAACAACAAGA	2010	31452	FLT1:2358L21 siRNA (2340C) inv stab10 antisense	GUUGGUGUUUUUAUGUUGUUTsT	2204

## VEGFR2

Target Pos	Target	Seq ID	Aliases	Sequence	Seq ID
3302	UGACCUUGGAGCAUCUCAUCUGU	2001	KDR:3304U21 siRNA sense	ACCUUGGAGCAUCUCAUCUTT	2044
3852	UUUGAGCAUGGAAGAGGAUUUCUG	2002	KDR:3854U21 siRNA sense	UGAGCAUGGAAGAGGAUUCUTT	2045
3892	UCACCUUUUCCUGUAUGGAGGA	2003	KDR:3894U21 siRNA sense	ACCUUUUCCUGUAUGGAGTT	2046
3946	GACAACACAGCAGGAUUCAGUCA	2004	KDR:3948U21 siRNA sense	CAACACAGCAGGAUUCAGUTT	2047
3302	UGACCUUGGAGCAUCUCAUCUGU	2001	KDR:3322L21 siRNA (3304C) antisense	AGAUGAGAUUGCUCCAAGGUTT	2048
3852	UUUGAGCAUGGAAGAGGAUUUCUG	2002	KDR:3872L21 siRNA (3854C) antisense	GAAUCCUUCUCCAUUGCUCATT	2049
3892	UCACCUUUUCCUGUAUGGAGGA	2003	KDR:3912L21 siRNA (3894C) antisense	CUCCAUACAGGAAACAGGUTT	2050
3946	GACAACACAGCAGGAUUCAGUCA	2004	KDR:3966L21 siRNA (3948C) antisense	ACUGAUUCCUGCUGUGUUGTT	2051
3302	UGACCUUGGAGCAUCUCAUCUGU	2001	KDR:3304U21 siRNA stab04 sense	B AccuuGGAGcAucucAucUTT B	2052
3852	UUUGAGCAUGGAAGAGGAUUUCUG	2002	KDR:3854U21 siRNA stab04 sense	B uGAGcAuGGAAAGAGGAuucTT B	2053
3892	UCACCUUUUCCUGUAUGGAGGA	2003	KDR:3894U21 siRNA stab04 sense	B AccuGuuuuccuGuuGGAGTT B	2054
3946	GACAACACAGCAGGAUUCAGUCA	2004	KDR:3948U21 siRNA stab04 sense	B cAAcAcAGcAGGAuucAGuTT B	2055
3302	UGACCUUGGAGCAUCUCAUCUGU	2001	KDR:3322L21 siRNA (3304C) stab05 antisense	AGAUAGAGAUGcuccAAAGGuTsT	2056
3852	UUUGAGCAUGGAAGAGGAUUUCUG	2002	KDR:3872L21 siRNA (3854C) stab05 antisense	GAAuccuuccAuGcuATsT	2057
3892	UCACCUUUUCCUGUAUGGAGGA	2003	KDR:3912L21 siRNA (3894C) stab05 antisense	cuccAuAcAGGAAAcAGGuTsT	2058
3946	GACAACACAGCAGGAUUCAGUCA	2004	KDR:3966L21 siRNA (3948C) stab05 antisense	AcuGAuuuccuGcuGuuGTsT	2059
3302	UGACCUUGGAGCAUCUCAUCUGU	2001	KDR:3304U21 siRNA stab07 sense	B AccuuGGAGcAucucAucUTT B	2060
3852	UUUGAGCAUGGAAGAGGAUUUCUG	2002	KDR:3854U21 siRNA stab07 sense	B uGAGcAuGGAAAGAGGAuucTT B	2061
3892	UCACCUUUUCCUGUAUGGAGGA	2003	KDR:3894U21 siRNA stab07 sense	B AccuGuuuuccuGuuGGAGTT B	2062
3946	GACAACACAGCAGGAUUCAGUCA	2004	KDR:3948U21 siRNA stab07 sense	B cAAcAcAGcAGGAuucAGuTT B	2063
3302	UGACCUUGGAGCAUCUCAUCUGU	2001	KDR:3322L21 siRNA (3304C) stab11 antisense	AGAUAGAGAUGcuccAAAGGuTsT	2064
3852	UUUGAGCAUGGAAGAGGAUUUCUG	2002	KDR:3872L21 siRNA (3854C) stab11 antisense	GAAuccuuccAuGcuATsT	2065
3892	UCACCUUUUCCUGUAUGGAGGA	2003	KDR:3912L21 siRNA (3894C) stab11 antisense	cuccAuAcAGGAAAcAGGuTsT	2066
3946	GACAACACAGCAGGAUUCAGUCA	2004	KDR:3966L21 siRNA (3948C) stab11 antisense	AcuGAuuuccuGcuGuuGTsT	2067

## VEGFR2

Target	SeqID	RPI#	Alias	Sequence	SeqID
UGUCCACUUAACCUAGGAGCAAG	2017	30785	KDR:3076U21 siRNA stab04 sense	B uccAcuuAccuGAGGAGcATT B	2205
UUUGAGCAUGGAAGAGGAUUCUG	2002	30786	KDR:3854U21 siRNA stab04 sense	B uGAGcAuGGAAAGAGGAuucTT B	2053
AUGGUUCUUGCCUCAGAGAGCU	2018	30787	KDR:4089U21 siRNA stab04 sense	B GGUucuuGccucAGAAAGAGTT B	2206
UCUGAAGGCUCAAACCAGACAAG	2019	30788	KDR:4191U21 siRNA stab04 sense	B uGAAGGcucAAAccAGAcATT B	2207
UGUCCACUUAACCUAGGAGCAAG	2017	30789	KDR:3094L21 siRNA (3076C) stab05 antisense	uGcuccucAGGuAAGuGGATsT	2208
UUUGAGCAUGGAAGAGGAUUCUG	2002	30790	KDR:3872L21 siRNA (3854C) stab05 antisense	GAAucccuuuccAuGcucATsT	2057
AUGGUUCUUGCCUCAGAGAGCU	2018	30791	KDR:4107L21 siRNA (4089C) stab05 antisense	cucuuucuGAGGcAAAGAAccTsT	2209
UCUGAAGGCUCAAACCAGACAAG	2019	30792	KDR:4209L21 siRNA (4191C) stab05 antisense	uGucuuGGuuuGAGccuucATsT	2210
UGUCCACUUAACCUAGGAGCAAG	2017	31426	KDR:3076U21 siRNA sense	UCCACUUUACCUGAGGAGGcATT	2211
UUUGAGCAUGGAAGAGGAUUCUG	2002	31427	KDR:3854U21 siRNA sense	UGAGCAUGGAAGAGGAUUCCTT	2045
AUGGUUCUUGCCUCAGAGAGCU	2018	31428	KDR:4089U21 siRNA sense	GGUUCUUGCCUCAGAAAGAGTT	2212
UCUGAAGGCUCAAACCAGACAAG	2019	31429	KDR:4191U21 siRNA sense	UGAAGGCUCAAACCAGAGcATT	2213
UGUCCACUUAACCUAGGAGCAAG	2017	31430	KDR:3094L21 siRNA (3076C) antisense	UGCUCUCACAGGUAAAGUGGATT	2214
UUUGAGCAUGGAAGAGGAUUCUG	2002	31431	KDR:3872L21 siRNA (3854C) antisense	GAAUCCUCUUCUCCAUUGCUCATT	2049
AUGGUUCUUGCCUCAGAGAGCU	2018	31432	KDR:4107L21 siRNA (4089C) antisense	CUCUUCUGAGGCAAGAAcCTT	2215
UCUGAAGGCUCAAACCAGACAAG	2019	31433	KDR:4209L21 siRNA (4191C) antisense	UGUCUGGUUUUGAGCCUUCATT	2216
UGACCUUGGAGCAUCUCAUCUGU	2001	31434	KDR:3304U21 siRNA sense	ACCUUGGAGCAUCUCAUCUUTT	2044
UUUGAGCAUGGAAGAGGAUUCUG	2002	31435	KDR:3854U21 siRNA sense	UGAGCAUGGAAGAGGAUUCCTT	2045
UCACCUUUUCCUGUAGGAGGA	2003	31436	KDR:3894U21 siRNA sense	ACCUUUUCCUGUAGGAGTT	2046
GACAACACAGCAGGAUUCAGUCA	2004	31437	KDR:3948U21 siRNA sense	CAACACAGCAGGAUUCAGUTT	2047
UGACCUUGGAGCAUCUCAUCUGU	2001	31438	KDR:3322L21 siRNA (3304C) antisense	AGAUGAGAUUGCUCUCCAGGUTT	2048
UUUGAGCAUGGAAGAGGAUUCUG	2002	31439	KDR:3872L21 siRNA (3854C) antisense	GAAUCCUCUUCUCCAUUGCUCATT	2049
UCACCUUUUCCUGUAGGAGGA	2003	31440	KDR:3912L21 siRNA (3894C) antisense	CUCCAUACAGGAAACAGGUTT	2050
GACAACACAGCAGGAUUCAGUCA	2004	31441	KDR:3966L21 siRNA (3948C) antisense	ACUGAUUCCUGCUGUGUUGTT	2051

(400/102)

## VEGFR3

Target Pos	Target	Seq ID	Aliases	Sequence	Seq ID
2009	AGCACUGCCACAAAGAAGUACCUG	2005	FLT4:2011U21 siRNA sense	CACUGCCACAAAGAAGUACCTT	2088
3919	CUGAAGCAGAGAGAGAAAGGCA	2006	FLT4:3921U21 siRNA sense	GAAGCAGAGAGAGAGAAAGGTT	2089
4036	AAAGAGGAACCAAGAGGACAAGA	2007	FLT4:4038U21 siRNA sense	AGAGGAACCAAGAGGACAATT	2070
4052	GACAAGAGGAGCAUGAAAGUGGA	2008	FLT4:4054U21 siRNA sense	CAAGAGGAGCAUGAAAGUGTT	2071
2009	AGCACUGCCACAAAGAAGUACCUG	2005	antisense	GGUACUUCUUGUGGCAGUGTT	2072
3919	CUGAAGCAGAGAGAGAAAGGCA	2006	FLT4:3939L21 siRNA (3921C)	CCUUCUCUCUCUCUGCUUCCTT	2073
4036	AAAGAGGAACCAAGAGGACAAGA	2007	antisense	UUGUCCUCCUUGGUUCCUCUTT	2074
4052	GACAAGAGGAGCAUGAAAGUGGA	2008	antisense	CACUUUCAUGCUCUCCUUCUUGTT	2075
2009	AGCACUGCCACAAAGAAGUACCUG	2005	FLT4:2011U21 siRNA stab04 sense	B cAcuGccAcAAGAAAGuAccTT B	2076
3919	CUGAAGCAGAGAGAGAAAGGCA	2006	FLT4:3921U21 siRNA stab04 sense	B GAAAGCAGAGAGAGAGAAAGGTT B	2077
4036	AAAGAGGAACCAAGAGGACAAGA	2007	FLT4:4038U21 siRNA stab04 sense	B AGAGGAACCAAGAGGACAATT B	2078
4052	GACAAGAGGAGCAUGAAAGUGGA	2008	FLT4:4054U21 siRNA stab04 sense	B cAAGAGGAGCAUGAAAGuGTT B	2079
2009	AGCACUGCCACAAAGAAGUACCUG	2005	antisense	GGuAcuucuuGuGGcAGuGTsT	2080
3919	CUGAAGCAGAGAGAGAAAGGCA	2006	FLT4:3939L21 siRNA (3921C) stab05	ccuucucucucucucGcuucTsT	2081
4036	AAAGAGGAACCAAGAGGACAAGA	2007	antisense	uuGuuccuccuGGuuccucucTsT	2082
4052	GACAAGAGGAGCAUGAAAGUGGA	2008	antisense	cAcuuucAuGcuuccuucuuGTsT	2083
2009	AGCACUGCCACAAAGAAGUACCUG	2005	FLT4:2011U21 siRNA stab07 sense	B cAcuGccAcAAGAAAGuAccTT B	2084
3919	CUGAAGCAGAGAGAGAAAGGCA	2006	FLT4:3921U21 siRNA stab07 sense	B GAAAGCAGAGAGAGAGAAAGGTT B	2085
4036	AAAGAGGAACCAAGAGGACAAGA	2007	FLT4:4038U21 siRNA stab07 sense	B AGAGGAACCAAGAGGACAATT B	2086
4052	GACAAGAGGAGCAUGAAAGUGGA	2008	FLT4:4054U21 siRNA stab07 sense	B cAAGAGGAGCAUGAAAGuGTT B	2087
2009	AGCACUGCCACAAAGAAGUACCUG	2005	antisense	GGuAcuucuuGuGGcAGuGTsT	2088
3919	CUGAAGCAGAGAGAGAAAGGCA	2006	antisense	ccuucucucucucucGcuucTsT	2089



4036	AAAGAGGAACCAAGGAGGACAAGA	2007	FLT4:4056L21 siRNA (4038C) stab11 antisense	uuGuccuccuGGuccucuTstT	2090
4052	GACAAGAGGAGCAUGAAAGUGGA	2008	FLT4:4072L21 siRNA (4054C) stab11 antisense	cAuuuucAuGcuccucuGTstT	2091

Uppercase = ribonucleotide  
u,c = 2'-deoxy-2'-fluoro U,C  
T = thymidine  
B = inverted deoxy abasic  
s = phosphorothioate linkage  
A = deoxy Adenosine  
G = deoxy Guanosine

Table IV

Non-limiting examples of Stabilization Chemistries for chemically modified siNA constructs

Chemistry	pyrimidine	Purine	cap	p=S	Strand
"Stab 1"	Ribo	Ribo	-	5 at 5'-end 1 at 3'-end	S/AS
"Stab 2"	Ribo	Ribo	-	All linkages	Usually AS
"Stab 3"	2'-fluoro	Ribo	-	4 at 5'-end 4 at 3'-end	Usually S
"Stab 4"	2'-fluoro	Ribo	5' and 3'-ends	-	Usually S
"Stab 5"	2'-fluoro	Ribo	-	1 at 3'-end	Usually AS
"Stab 6"	2'-O-Methyl	Ribo	5' and 3'-ends	-	Usually S
"Stab 7"	2'-fluoro	2'-deoxy	5' and 3'-ends	-	Usually S
"Stab 8"	2'-fluoro	2'-O-Methyl	-	1 at 3'-end	Usually AS
"Stab 9"	Ribo	Ribo	5' and 3'-ends	-	Usually S
"Stab 10"	Ribo	Ribo	-	1 at 3'-end	Usually AS
"Stab 11"	2'-fluoro	2'-deoxy	-	1 at 3'-end	Usually AS

CAP = any terminal cap, see for example Figure 10.

All Stab 1-11 chemistries can comprise 3'-terminal thymidine (TT) residues

All Stab 1-11 chemistries typically comprise 21 nucleotides, but can vary as described herein.

S = sense strand

AS = antisense strand

Table V

A. 2.5  $\mu$ mol Synthesis Cycle ABI 394 Instrument

Reagent	Equivalents	Amount	Wait Time* DNA	Wait Time* 2'-O-methyl	Wait Time*RNA
Phosphoramidites	6.5	163 $\mu$ L	45 sec	2.5 min	7.5 min
S-Ethyl Tetrazole	23.8	238 $\mu$ L	45 sec	2.5 min	7.5 min
Acetic Anhydride	100	233 $\mu$ L	5 sec	5 sec	5 sec
N-Methyl Imidazole	186	233 $\mu$ L	5 sec	5 sec	5 sec
TCA	176	2.3 mL	21 sec	21 sec	21 sec
Iodine	11.2	1.7 mL	45 sec	45 sec	45 sec
Beaucage	12.9	645 $\mu$ L	100 sec	300 sec	300 sec
Acetonitrile	NA	6.67 mL	NA	NA	NA

B. 0.2  $\mu$ mol Synthesis Cycle ABI 394 Instrument

Reagent	Equivalents	Amount	Wait Time* DNA	Wait Time* 2'-O-methyl	Wait Time*RNA
Phosphoramidites	15	31 $\mu$ L	45 sec	233 sec	465 sec
S-Ethyl Tetrazole	38.7	31 $\mu$ L	45 sec	233 min	465 sec
Acetic Anhydride	655	124 $\mu$ L	5 sec	5 sec	5 sec
N-Methyl Imidazole	1245	124 $\mu$ L	5 sec	5 sec	5 sec
TCA	700	732 $\mu$ L	10 sec	10 sec	10 sec
Iodine	20.6	244 $\mu$ L	15 sec	15 sec	15 sec
Beaucage	7.7	232 $\mu$ L	100 sec	300 sec	300 sec
Acetonitrile	NA	2.64 mL	NA	NA	NA

C. 0.2  $\mu$ mol Synthesis Cycle 96 well Instrument

Reagent	Equivalents: DNA/ 2'-O-methyl/Ribo	Amount: DNA/2'-O- methyl/Ribo	Wait Time* DNA	Wait Time* 2'-O- methyl	Wait Time* Ribo
Phosphoramidites	22/33/66	40/60/120 $\mu$ L	60 sec	180 sec	360sec
S-Ethyl Tetrazole	70/105/210	40/60/120 $\mu$ L	60 sec	180 min	360 sec
Acetic Anhydride	265/265/265	50/50/50 $\mu$ L	10 sec	10 sec	10 sec
N-Methyl Imidazole	502/502/502	50/50/50 $\mu$ L	10 sec	10 sec	10 sec
TCA	238/475/475	250/500/500 $\mu$ L	15 sec	15 sec	15 sec
Iodine	6.8/6.8/6.8	80/80/80 $\mu$ L	30 sec	30 sec	30 sec
Beaucage	34/51/51	80/120/120	100 sec	200 sec	200 sec
Acetonitrile	NA	1150/1150/1150 $\mu$ L	NA	NA	NA

- 5 • Wait time does not include contact time during delivery.
- Tandem synthesis utilizes double coupling of linker molecule

CLAIMS

What we claim is:

1. A double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a vascular endothelial growth factor receptor (VEGFr) gene, wherein said siNA molecule comprises about 21 nucleotides.
2. The siNA molecule of claim 1, wherein said siNA molecule comprises no ribonucleotides.
3. The siNA molecule of claim 1, wherein said siNA molecule comprises ribonucleotides.
4. The siNA molecule of claim 1, wherein one of the strands of said double-stranded siNA molecule comprises a nucleotide sequence that is complementary to a nucleotide sequence or a portion thereof of a VEGFr gene, and wherein the second strand of said double-stranded siNA molecule comprises a nucleotide sequence substantially similar to the nucleotide sequence or a portion thereof of said VEGFr gene.
5. The siNA molecule of claim 4, wherein each said strand of the siNA molecule comprises about 19 to about 23 nucleotides, and wherein each said strand comprises at least about 19 nucleotides that are complementary to the nucleotides of the other strand.
6. The siNA molecule of claim 1, wherein said siNA molecule comprises an antisense region comprising a nucleotide sequence that is complementary to a nucleotide sequence or a portion thereof of a VEGFr gene, and wherein said siNA further comprises a sense region, wherein said sense region comprises a nucleotide sequence substantially similar to the nucleotide sequence or a portion thereof of said VEGFr gene.
7. The siNA molecule of claim 6, wherein said antisense region and said sense region each comprise about 19 to about 23 nucleotides, and wherein said antisense region comprises at least about 19 nucleotides that are complementary to nucleotides of the sense region.
8. The siNA molecule of claim 1, wherein said siNA molecule comprises a sense region and an antisense region and wherein said antisense region comprises a nucleotide sequence that is complementary to a nucleotide sequence or a portion thereof of RNA


encoded by a VEGFr gene and said sense region comprises a nucleotide sequence that is complementary to said antisense region.

9. The siNA molecule of claim 6, wherein said siNA molecule is assembled from two separate oligonucleotide fragments wherein one fragment comprises the sense region and the second fragment comprises the antisense region of said siNA molecule.
10. The siNA molecule of claim claim 6, wherein said sense region is connected to the antisense region via a linker molecule.
11. The siNA molecule of claim 10, wherein said linker molecule is a polynucleotide linker.
12. The siNA molecule of claim 10, wherein said linker molecule is a non-nucleotide linker.
13. The siNA molecule of claim 6, wherein pyrimidine nucleotides in the sense region are 2'-O-methyl pyrimidine nucleotides.
14. The siNA molecule of claim 6, wherein purine nucleotides in the sense region are 2'-deoxy purine nucleotides.
15. The siNA molecule of claim 6, wherein the pyrimidine nucleotides present in the sense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides.
16. The siNA molecule of claim 9, wherein the fragment comprising said sense region includes a terminal cap moiety at the 5'-end, the 3'-end, or both of the 5' and 3' ends of the fragment comprising said sense region.
17. The siNA molecule of claim 16, wherein said terminal cap moiety is an inverted deoxy abasic moiety.
18. The siNA molecule of claim 6, wherein the pyrimidine nucleotides of said antisense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides.
19. The siNA molecule of claim 6, wherein the the purine nucleotides of said antisense region are 2'-O-methyl purine nucleotides.
20. The siNA molecule of claim 6, wherein the purine nucleotides present in said antisense region comprise 2'-deoxy- purine nucleotides.
21. The siNA molecule of claim 18, wherein said antisense region comprises a phosphorothioate internucleotide linkage at the 3' end of said antisense region.

22. The siNA molecule of claim 6, wherein said antisense region comprises a glyceryl modification at the 3' end of said antisense region.
23. The siNA molecule of claim 9, wherein each of the two fragments of said siNA molecule comprise 21 nucleotides.
- 5 24. The siNA molecule of claim 23, wherein about 19 nucleotides of each fragment of the siNA molecule are base-paired to the complementary nucleotides of the other fragment of the siNA molecule and wherein at least two 3' terminal nucleotides of each fragment of the siNA molecule are not base-paired to the nucleotides of the other fragment of the siNA molecule.
- 10 25. The siNA molecule of claim 24, wherein each of the two 3' terminal nucleotides of each fragment of the siNA molecule are 2'-deoxy-pyrimidines.
26. The siNA molecule of claim 25, wherein said 2'-deoxy-pyrimidine is 2'-deoxy-thymidine.
- 15 27. The siNA molecule of claim 23, wherein all 21 nucleotides of each fragment of the siNA molecule are base-paired to the complementary nucleotides of the other fragment of the siNA molecule.
28. The siNA molecule of claim 23, wherein about 19 nucleotides of the antisense region are base-paired to the nucleotide sequence or a portion thereof of the RNA encoded by a VEGFr gene.
- 20 29. The siNA molecule of claim 23, wherein 21 nucleotides of the antisense region are base-paired to the nucleotide sequence or a portion thereof of the RNA encoded by a VEGFr gene.
30. The siNA molecule of claim 9, wherein the 5'-end of the fragment comprising said antisense region optionally includes a phosphate group.
- 25 31. The siNA molecule of claim 1, wherein said VEGFr gene is VEGFr1.
32. The siNA molecule of claim 1, wherein said VEGFr gene is VEGFr2.
33. The siNA molecule of claim 1, wherein said VEGFr gene is VEGFr3.
34. A double-stranded short interfering nucleic acid (siNA) molecule that inhibits the expression of a VEGFr gene, wherein said siNA molecule comprises no

ribonucleotides and wherein each strand of said double-stranded siNA molecule comprises about 21 nucleotides.

35. The siNA molecule of claim 34, wherein said VEGFr gene is VEGFr1.
36. The siNA molecule of claim 34, wherein said VEGFr gene is VEGFr2.
- 5 37. The siNA molecule of claim 34, wherein said VEGFr gene is VEGFr3.
38. A double-stranded short interfering nucleic acid (siNA) molecule that inhibits the expression of a VEGFr gene, wherein said siNA molecule does not require the presence of a ribonucleotide within the siNA molecule for said inhibition of expression of the VEGFr gene and wherein each strand of said double-stranded siNA molecule comprises about 21 nucleotides.
- 10 39. The siNA molecule of claim 38, wherein said VEGFr gene is VEGFr1.
40. The siNA molecule of claim 38, wherein said VEGFr gene is VEGFr2.
41. The siNA molecule of claim 38, wherein said VEGFr gene is VEGFr3.
42. A pharmaceutical composition comprising the siNA molecule of claim 1 in an acceptable carrier or diluent.
- 15 43. Medicament comprising the siNA molecule of claim 1.
44. Active ingredient comprising the siNA molecule of claim 1.
45. Use of a double-stranded short interfering nucleic acid (siNA) molecule to down-regulate expression of a VEGFr gene, wherein said siNA molecule comprises one or more chemical modifications and each strand of said double-stranded siNA comprises about 21 nucleotides.
- 20


**Figure 1**

= SOLID SUPPORT


R = TERMINAL PROTECTING GROUP

FOR EXAMPLE:



DIMETHOXYTRITYL (DMT)



(1) = CLEAVABLE LINKER


(FOR EXAMPLE: NUCLEOTIDE SUCCINATE OR  
INVERTED DEOXYABASIC SUCCINATE)

(2) = CLEAVABLE LINKER


(FOR EXAMPLE: NUCLEOTIDE SUCCINATE OR  
INVERTED DEOXYABASIC SUCCINATE)INVERTED DEOXYABASIC SUCCINATE  
LINKAGE


GLYCERYL SUCCINATE LINKAGE




*Figure 2*

**Figure 3**



**Figure 4**

POSITIONS (NN) CAN COMPRISE ANY NUCLEOTIDE, SUCH AS DEOXYNUCLEOTIDES (eg. THYMIDINE) OR UNIVERSAL BASES  
B = ABASIC, INVERTED ABASIC, INVERTED NUCLEOTIDE OR OTHER TERMINAL CAP THAT IS OPTIONALLY PRESENT  
L = GLYCERYL MOIETY THAT IS OPTIONALLY PRESENT  
S = PHOSPHOROTHIOATE OR PHOSPHORODITHIOATE

**Figure 5**

lower case = 2'-O-Methyl or 2'-deoxy-2'-fluoro

*italic lower case* = 2'-deoxy-2'-fluorounderline = 2'-O-methyl*ITALIC UPPER CASE* = DEOXY


B = INVERTED DEOXYABASIC

L = GLYCERYL MOIETY OPTIONALLY PRESENT


S = PHOSPHOROTHIOATE OR

PHOSPHORODITHIOATE


Figure 6




**Figure 7**










**Figure 10**

R = O, S, N, alkyl, substituted alkyl, O-alkyl, S-alkyl, alkaryl, or aralkyl  
B = Independently any nucleotide base, either naturally occurring or chemically modified, or optionally H (abasic).

*Figure 11: Modification Strategy*



**Figure 12: Inhibition of VEGF-Induced Angiogenesis  
by siRNAs**





Figure 13: A375 24h 36B4 VEGFR1 mRNA Expression



**This Page Blank (uspto)**

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☒ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**